

# Evidence-Based Interventions: Guidance for CCGs

**Published by NHS England in partnership with NHS Clinical Commissioners, the Academy of Medical Royal Colleges, NHS Improvement and the National Institute for Health and Care Excellence**

## **Evidence-Based Interventions: Guidance for CCGs**

Version number: 2.0

First published: 28 November 2018

Updated: 11 January 2019

Prepared by: NHS England Medical directorate and Strategy and Innovation directorate

Classification: OFFICIAL

NHS England Publications Gateway Reference: 08659

This information can be made available in alternative formats, such as easy read or large print, and may be available in alternative languages, upon request. Please contact the Evidence-Based Interventions team at [\*\*england.EBInterventions@nhs.net\*\*](mailto:england.EBInterventions@nhs.net).

# Contents

Contents .....	3
Background .....	4
1.1 Who is this guidance for? .....	4
1.2 Why have we developed this guidance? .....	4
1.3 How have the recommendations in this guidance been developed? .....	5
1.4 How have the recommendations in this guidance been developed following the results of the consultation? .....	6
1.5 Updating the guidance.....	7
Delivery actions .....	8
1.6 The NHS Standard Contract and Category 1 and 2 interventions .....	8
1.7 Introduce zero payment for Category 1 interventions without IFRs .....	9
1.8 Implement an IFR process for Category 1 interventions and a prior approval process for Category 2 interventions .....	9
1.9 National and local activity goals .....	10
1.10 Integrated monthly dashboard to monitor delivery .....	11
1.11 Local system audits to review compliance .....	12
1.12 STP and CCG Improvement and Assessment Framework .....	12
1.13 Aligning CQC inspection with the policy .....	12
Clinical criteria .....	13
Activity goals.....	48
Appendix 1: Equality and Health Inequalities Analysis Form .....	136
Appendix 2: Clinical glossary .....	172
Appendix 3: Technical glossary .....	177
Appendix 4: Procedure and diagnostic codes.....	178

# Background

## 1.1 Who is this guidance for?

1. This guidance is addressed to CCGs, to assist them in fulfilling their duties relating to securing continuous improvements in the quality of services for patients and in outcomes, particularly regarding appropriate clinical interventions. This guidance is issued as general guidance under S14Z8 of the NHS Act 2006. The objective of this guidance is to support CCGs in their decision-making, to address unwarranted variation, and to provide national advice to make local clinical decision-making more appropriate.
2. We expect CCGs to have regard to this guidance in formulating local policies and for clinicians to reflect this guidance in their clinical practice. This guidance does not remove the clinical discretion of clinicians in accordance with their professional duties.

## 1.2 Why have we developed this guidance?

3. We want to reduce the number of inappropriate interventions provided on the NHS. The primary goals of the Evidence-Based Interventions programme are to avoid needless harm to patients and free-up scarce professional time for performing other interventions - including creating headroom for proven innovations. The time saved will be reinvested in patient care.
4. Last year, the 17 interventions listed in this guidance were provided over 335,000 times. We know that across England there is substantial variation in the rate at which these interventions are performed, to an extent that cannot be explained by differences in population demand. This has an impact on clinical outcomes and contributes to perceptions of a so-called 'postcode lottery' by members of the public.
5. There is also widespread clinical consensus that NHS resources could be more appropriately targeted towards more clinically appropriate interventions. At a time when demand is exceeding the capacity available, effective use of resources is both a national and local priority.
6. NHS England has partnered with NHS Clinical Commissioners, the National Institute for Health and Care Excellence (NICE), the Academy of Royal Medical Colleges and NHS Improvement to develop this guidance.
7. The guidance, and the Evidence-Based Interventions programme as a whole, is guided by the following five goals:

- **Reduce avoidable harm to patients.** With surgical interventions, there is always a risk of complications. Weighing the risks and benefits of appropriate treatments should be co-produced with patients.
  - **Save precious professional time**, when the NHS is severely short of staff, professionals should offer appropriate and effective treatment to patients.
  - **Help clinicians maintain their professional practice** and keep up to date with the changing evidence base and best practice.
  - **Create headroom for innovation.** If we want to accelerate the adoption of new, proven innovations, we need to reduce the number of inappropriate interventions. This allows innovation in healthcare, prescribing and technology to improve patients' ability to self-care and live with long term conditions.
  - **Maximise value and avoid waste.** Inappropriate care is poor value for the taxpayer. Resources should be focused on effective and appropriate NHS services
8. Our ambition is to support systems to improve clinical outcomes for their population by ensuring that patients only receive interventions for which there is an established, high-quality evidence base. Similarly, we also hope that this national guidance will lead to more standardised local commissioning policies.

### 1.3 How have the recommendations in this guidance been developed?

9. The Evidence-based Interventions Programme has been shaped by six design principles that outline our commitment to: clinical research and evidence (as contained in NICE and/or NICE accredited and specialist society guidelines); best practice; patient and public involvement and engagement; and successful implementation that involves a range of activities.
10. With regard to the final set of 17 interventions listed in this guidance, we initially identified a large number of recommendations from clinical evidence including NICE guidance, national and international Choosing Wisely recommendations, academic studies and local CCGs' work.<sup>1</sup>
11. Taking these as a starting point, we shortlisted them by:
- Prioritising interventions that we could test our approach on and implement relatively quickly on a large scale. We focused on interventions commissioned by CCGs, where there was high variability in the application of clinical guidelines.
  - Working with the Royal Colleges, clinicians, clinical commissioners and professional leaders to refine the list, ensuring clinical consensus and speciality buy-in.

---

<sup>1</sup> This includes NICE ['do not do' recommendations](#); NICE [Cost Saving Guidance](#); <http://www.choosingwisely.co.uk/i-am-a-clinician/recommendations/>; <https://choosingwiselycanada.org/recommendations/>; <http://www.choosingwisely.org.au/recommendations/>; <http://www.choosingwisely.org/clinician-lists/>. See Section 3 for full references.

- Working with NHS Clinical Commissioners as the representative organisation for CCGs.
  - Initially liaising with a number of patients and patient representative groups to test the proposals and understand their priorities, including Healthwatch.
  - Aligning our approach with national programmes like the NHS RightCare programme and NHS Improvement's Getting It Right First Time programme.
12. In addition, we carried out an initial equality impact assessment on the proposals to determine any differential impacts across groups with protected characteristics. The result of this initial assessment can be found in Appendix 4 of the original consultation document. A final assessment, conducted following a review of the consultation evidence, can be found in Appendix 1.
13. Finally, we segmented the seventeen interventions into two groups:
- Interventions that should not be routinely commissioned, with patients only able to access such treatments where they successfully make an individual funding request (referred to as Category 1 interventions).
  - Interventions that should be commissioned or performed when specific criteria are met (referred to as Category 2 interventions).
14. Each individual intervention was reviewed by one or more appropriate clinical groups. The NHS England Medical Advisory Group, comprising national clinical directors, supported the final shortlist. We sought feedback from patients throughout the design process (see Appendix 1 of the full consultation response document for further details<sup>2</sup>).
15. Note that all of the clinical criteria consulted on were developed directly from existing NICE, NICE-accredited or specialist society guidance and local CCG policies, and the final set of wording used has been checked by the relevant Royal Colleges, specialist societies, individual specialists, as well as clinical experts from within NHS England.
16. A full public consultation on the proposals, announced at the NHS England Board Meeting in July 2018, was conducted between the 4<sup>th</sup> of July and the 28<sup>th</sup> September 2018. Section 1.4 details the main findings from the consultation and the changes that have been made as a result of what we heard. A more detailed report on the consultation can be found in 'Evidence-Based Interventions Policy: Response to the public consultation and next steps'<sup>3</sup>.

#### **1.4 How have the recommendations in this guidance been developed following the results of the consultation?**

17. Whilst the overall number of interventions remains unchanged from those listed in the consultation document, we have made important refinements and

---

<sup>2 2</sup> Please visit: <https://www.england.nhs.uk/evidence-based-interventions/>

<sup>3 3</sup> Please visit: <https://www.england.nhs.uk/evidence-based-interventions/>

clarifications to the clinical criteria in response to feedback during the consultation period (see Section 3 for the final set of criteria and Appendix 2 for a Clinical glossary). These changes include:

- Expanding the recommendation wording for carpal tunnel syndrome, Dupuytren's contracture release, ganglion excision and trigger finger release to align with proposals from the British Society of the Hand.
- Excluding children from the criteria related to Dupuytren's contracture release, trigger finger release and snoring surgery as these conditions present differently in children and may indicate more serious underlying conditions.
- Clarifying that children who cannot undergo normal assessments are still able to access specialist advice for glue ear.
- Accounting for a wide range of views on coding methodology, testing this with coding experts, clinicians, and demonstrator sites, and updating the activity projections accordingly.

18. We have also refined the delivery actions proposed, clarifying for example, our expectations regarding waiting lists and referrals for category 1 interventions and the interaction with proposed Tariff changes that could take effect from 1 April 2019 (see Section 2 for the final set of delivery actions, Section 4 for the local activity goals by CCG, STP and provider, Appendix 3 for a Technical glossary and Appendix 4 for Procedure and diagnostic codes).

19. We have also established a new national steering group which includes patient and clinical representatives. This group has helped shape our response, and will guide our implementation. We have also established a demonstrator community of local geographies to test implementation of the Evidence-Based Intervention programme as well as ideas for future phases of the programme.

## **1.5 Updating the guidance**

20. The Evidence-Based Interventions programme will monitor progress of this programme ahead of considering further expansion.

21. In the full consultation response document, we reference the intention to continue with future phases of the Evidence-Based Interventions programme. These phases represent appropriate points to reconsider the evidence base related to these and other interventions, and we intend to update the guidance through this mechanism as appropriate.

# Delivery actions

## 1.6 The NHS Standard Contract and Category 1 and 2 interventions

22. In addition to publishing statutory guidance under Section 14Z8 of the NHS Act 2006, we are mandating compliance to the Evidence-Based Interventions programme through the NHS Standard Contract (see Section 1.8.1 of the full consultation response document). The new wording will be added to Service Condition 29, as set out below. (See Appendix 3 for the Technical glossary. Note that the definition of Prior Approval Schemes in The NHS Standard Contract includes both Individual Funding Requests and Prior Approval).

### ***Evidence-Based Interventions Policy***

29.28 The Parties must comply with their respective obligations under the Evidence-Based Interventions Policy.

29.29 The Commissioners must use all reasonable endeavours to procure that, when making Referrals, Referrers comply with the Evidence-Based Interventions Policy.

29.30 The Provider must manage Referrals and provide the Services in accordance with the Evidence-Based Interventions Policy.

29.31 If the Provider carries out

29.31.1 a Category 1 Intervention without evidence of appropriate Prior Approval having been granted by the relevant Commissioner; or

29.31.2 a Category 2 Intervention other than in accordance with the Evidence-Based Interventions Policy, the relevant Commissioner will not be liable to pay for that Intervention.

29.32 For the avoidance of doubt, any Commissioner may, at its absolute discretion, impose by means of a Prior Approval Scheme notified to the Provider in accordance with SC29.24 (Prior Approval Scheme) preconditions in relation to any Category 1 Intervention or Category 2 Intervention more stringent than those set out or referred to in SC29.28 – SC29.31 and/or the Evidence-Based Interventions Policy.

23. For reference, the associated definitions within the Contract will be:

- **Evidence-Based Interventions Policy** - the national policy relating to the commissioning of interventions which are clinically inappropriate or which are appropriate only when performed in specific circumstances, published by NHS England.

- **Category 1 Interventions** - interventions which should not be routinely commissioned or performed, described as Category 1 Interventions in Evidence-Based Interventions Policy.
- **Category 2 Interventions** - interventions which should only be routinely commissioned or performed when specific criteria are met, described as Category 2 Interventions in Evidence-Based Interventions Policy.

## **1.7 Introduce zero payment for Category 1 interventions without IFRs**

24. To ensure that Category 1 interventions as described in the Evidence-Based Interventions Policy are no longer funded, unless accompanied by an IFR, we will work with NHS Improvement to create a new National Variation to the Tariff, subject to the statutory consultation on the 2019/20 National Tariff Payment System.
25. We propose that the National Variation sets out that only activity that meets the IFR criteria will be paid for. Any activity that does not meet this threshold will be reimbursed at £0. This option means that activity undertaken without an IFR will not be reimbursed and will avoid local differentiation in pricing.

## **1.8 An IFR process for Category 1 interventions and a prior approval process for Category 2 interventions**

26. The four interventions we have classified as Category 1 are interventions that should not be routinely offered to patients unless there is a clinical exception as per the Evidence-Based Interventions Policy. The best route to ensure access in exceptional circumstances is through the IFR process.
27. We therefore ask local commissioners and providers to design IFR processes which utilise relevant available expertise and respond to applicants according to the urgency of their needs. With regard to who should be responsible for submitting the IFR, we will leave it to local areas to decide but suggest that it should be the treating clinician. This should, where appropriate, enable referrals to a specialist for an opinion.
28. For the 13 Category 2 interventions, clinicians will need to demonstrate that the patient meets the criteria set out in this guidance. CCGs will need to ensure compliance. Where there are concerns about achieving the desired clinical change and proposed activity reduction goals, we encourage the use of measures such as a prior approval process. In considering the use of prior approval, we propose local areas also consider category 2 interventions be monitored through regular audits and engagement with clinicians and, if needed, be reinforced through financial levers.

29. With regard to who should be responsible for submitting the prior approval, we will leave it to local areas to decide but suggest that it could be the treating clinician. This should, where appropriate, enable referrals to a specialist for an opinion.
30. While changes to the NHS Standard Contract and National Tariff Payment System will not be in place until 1 April 2019, set out below is the transitional period between the date this document is published and 1 April 2019. The implications and rationale for this period are:
- **National clinical criteria:** CCGs, providers and clinicians will be expected to start to implement the criteria for Category 1 and 2 interventions from the date that this document is published (28 November 2018). The rationale for this is that we want to ensure patients have access to the most appropriate intervention as soon as possible and to minimise avoidable harm to patients.
  - **Waiting lists:** Changes to reimbursement for Category 1 and 2 interventions will come into effect on 1 April and will apply to all patients added to waiting lists after the statutory consultation on the National Tariff Payment System for 2019/20 is published (expected on 17 January 2019), unless an IFR is in place. This will give NHS trusts and GPs time to change clinical practice before 17 January 2019 while also ensuring that we act quickly so that patients are not subject to inappropriate interventions.

## 1.9 National and local activity goals

29. By April 1<sup>st</sup> 2019, we expect that:

- No Category 1 interventions be performed unless accompanied by an IFR and therefore the numbers of activity for Category 1 without IFRs to reduce to near zero.
- Category 2 interventions should be reduced to the 25<sup>th</sup> percentile of the age-sex standardised rate of CCGs.

30. The national inpatient activity goals are set out in Table 1 below. We have also set out activity goals by CCG and STP for 2019/20 and 2020/21 in Section 4 of this guidance (and see Appendix 4 for Procedure and diagnostic codes). These activity goals will form part of CCGs' overall assessment.

**Table 1: National activity figures<sup>4</sup>**

	Intervention	No of spells (2017/18) Total CCG activity	CCG Variation (n fold variation) <sup>5</sup>	CCG Activity reduction opportunity	CCG Remaining Activity
<b>A</b>	Intervention for snoring (not OSA)	812	-. <sup>6</sup>	812	0
<b>B</b>	Dilatation & curettage for heavy menstrual bleeding	236	-. <sup>7</sup>	236	0
<b>C</b>	Knee arthroscopy with osteoarthritis	3,437	11.3	3,437	0
<b>D</b>	Injection for nonspecific low back pain without sciatica	13,165	31.4	13,165	0
<b>Total: category 1</b>		<b>17,650</b>	<b>-</b>	<b>17,650</b>	<b>-</b>
<b>E</b>	Breast reduction	2,388	8.4	829	1,559
<b>F</b>	Removal of benign skin lesions	116,255	4.1	45,589	70,666
<b>G</b>	Grommets	8,669	6.2	3,259	5,410
<b>H</b>	Tonsillectomy	32,238	3.0	7,454	24,784
<b>I</b>	Haemorrhoid surgery	8,474	4.3	2,801	5,673
<b>J</b>	Hysterectomy for heavy bleeding	27,660	3.3	6,536	21,124
<b>K</b>	Chalazia removal	6,026	29.7	4,326	1,700
<b>L</b>	Shoulder decompression	13,930	9.1	6,807	7,123
<b>M</b>	Carpal tunnel syndrome release	44,497	5.3	14,950	29,547
<b>N</b>	Dupuytren's contracture release	14,376	4.1	4,113	10,263
<b>O</b>	Ganglion excision	6,219	6.4	2,509	3,710
<b>P</b>	Trigger finger release	7,789	5.7	2,582	5,207
<b>Q</b>	Varicose vein surgery	28,846	8.0	8,633	20,213
<b>Total: category 2</b>		<b>317,367</b>	<b>-</b>	<b>110,388</b>	
<b>Grand Total</b>		<b>335,017</b>	<b>-</b>	<b>128,038</b>	

31. The Evidence-Based Interventions Programme and the clinical criteria for the 17 interventions apply in all care settings. However, the 2017/18 activity and activity goals set out in the data tables are necessarily based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. This is due to limitations in what we are reliably able to measure nationally. Outpatient activity is therefore not included. We will work with our demonstrator community to improve data for both in and outpatient settings.

## 1.10 Integrated monthly dashboard to monitor delivery

<sup>4</sup> National activity figures are based on CCG activity figures excluding activity that could not be attributed to a CCG.

<sup>5</sup> The variation is the ratio between the 10th highest and 10th lowest age-sex standardised rate between CCGs

<sup>6</sup> 20 CCGs have no activity recorded for this intervention. It has therefore not been possible to calculate an age-sex standardised variation rate for this intervention.

<sup>7</sup> 89 CCGs have no activity recorded for this intervention. It has therefore not been possible to calculate an age-sex standardised variation rate for this intervention.

32. We plan to rollout a joint dashboard which provides a quarterly update on CCG and provider activity for each of the 17 interventions and the 18 items that should no longer be routinely prescribed in primary care.<sup>8</sup> It will include trend analysis and comparative figures as well as target figures. NHS Business Services Authority will support the production of the dashboard which will be available to all CCGs by the end of 2018.

### **1.11 Local system audits to review compliance**

33. With guidance from specialist and primary care clinicians, we will work with NHS Improvement and NHS Clinical Commissioners to review local compliance to the programme. We will include the Evidence-Based Interventions programme in the upcoming planning guidance and will work with our regional and local colleagues to ensure that these plans are understood and implemented. We are working with regional teams in both NHS England and NHS Improvement to ensure we can support adherence to the guidance presented in this report as part of local assurance processes.

### **1.12 STP and CCG Improvement and Assessment Framework**

34. Following the consultation, we will seek to develop an indicator that could be considered for inclusion in the next iteration of the CCG IAF. We have tested this proposal with our demonstrator community and will work with them and NICE to develop the indicator. The indicator would measure performance of local areas against the Evidence-Based Interventions guidance and would be calculated using activity data.

### **1.13 Aligning CQC inspection with the policy**

35. We will continue to work with CQC to incorporate the clinical guidance on the 17 interventions in to their inspection framework from the roll out of this programme on April 1st 2019. To deliver this we will agree a data sharing protocol with the Care Quality Commission in order to populate their data analysis tool known as 'Insight' with provider activity figures which will inform their inspection programme for NHS trusts.

36. We are also aware that some patients may seek to get access to these treatments privately even if they are not appropriate. To limit the 17 interventions set out in this document from being offered inappropriately, we do not expect NHS providers to offer these interventions privately. We have agreed with CQC that this will be monitored through regional assurance processes and CQC inspections.

---

<sup>8</sup> Further details of the programme can be found here: <https://www.england.nhs.uk/medicines/items-which-should-not-be-routinely-prescribed/>

# Clinical criteria

## A. Adult Snoring Surgery (in the absence of OSA)

### Updated description of the intervention

In two systematic reviews of 72 primary research studies, there was no evidence that surgery to the palate to improve snoring provides any additional benefit compared to non-surgical treatments. The surgery has up to 16% risk of severe complications (bleeding, airway compromise, death). Therefore it is no longer commissioned. A number of alternatives to surgery can improve snoring. These include lifestyle changes (weight loss, smoking cessation and reducing alcohol intake) and medical treatment of nasal congestion.

### Updated clinical criteria

#### Summary of intervention

Snoring is a noise that occurs during sleep that can be caused by vibration of tissues of the throat and palate. It is very common and as many as one in four adults snore, as long as it is not complicated by periods of apnoea (temporarily stopping breathing) it is not usually harmful to health, but can be disruptive, especially to a person's partner.

This guidance relates to surgical procedures in adults to remove, refashion or stiffen the tissues of the soft palate (Uvulopalatopharyngoplasty, Laser assisted Uvulopalatoplasty & Radiofrequency ablation of the palate) in an attempt to improve the symptom of snoring. Please note this guidance only relates to patients with snoring in the absence of Obstructive Sleep Apnoea (OSA) and should not be applied to the surgical treatment of patients who snore and have proven OSA who may benefit from surgical intervention as part of the treatment of the OSA.

It is important to note that snoring can be associated with multiple other causes such as being overweight, smoking, alcohol or blockage elsewhere in the upper airways (e.g. nose or tonsils) and often these other causes can contribute to the noise alongside vibration of the tissues of the throat and palate.

#### Number of CCG interventions in 2017/18

812

#### Recommendation

It is on the basis of limited clinical evidence of effectiveness, and the significant risks that patients could be exposed to, this procedure should no longer be routinely commissioned in the management of simple snoring.

#### Alternative Treatments

There are a number of alternatives to surgery that can improve the symptom of snoring. These include:

- Weight loss

- Stopping smoking
- Reducing alcohol intake
- Medical treatment of nasal congestion (rhinitis)
- Mouth splints (to move jaw forward when sleeping)

### Rationale for recommendation

In two systematic reviews of 72 primary research studies there is no evidence that surgery to the palate to improve snoring provides any additional benefit compared to other treatments. While some studies demonstrate improvements in subjective loudness of snoring at 6-8 weeks after surgery; this is not longstanding (> 2years) and there is no long-term evidence of health benefit. This intervention has limited to no clinical effectiveness and surgery carries a 0-16% risk of severe complications (including bleeding, airway compromise and death). There is also evidence from systematic reviews that up to 58-59% of patients suffer persistent side effects (swallowing problems, voice change, globus, taste disturbance & nasal regurgitation). It is on this basis the interventions should no longer be routinely commissioned.

### References

1. Franklin KA, Anttila H, Axelsson S, Gislason T, Maasilta P, Myhre KI, Rehnqvist N. Effects and side-effects of surgery for snoring and obstructive sleep apnoea- a systematic review. *Sleep*. 2009 Jan. 32(1):27-36
2. Main C, Liu Z, Welch K, Weiner G, Jones SQ, Stein K. Surgical procedures and non-surgical devices for the management of non-apnoeic snoring: a systematic review of clinical effects and associated treatment costs. *Health Technol Assess* 2009;13(3). <https://www.ncbi.nlm.nih.gov/pubmed/19091167>
3. Jones TM, Earis JE, Calverley PM, De S, Swift AC. Snoring surgery: A retrospective review. *Laryngoscope*. 2005 Nov 115(11): 2015-20. <https://www.ncbi.nlm.nih.gov/pubmed/16319615>

## B. Dilatation and curettage (D&C) for heavy menstrual bleeding in women

### Updated description of the intervention

NICE guidelines recommend that D&C is not offered as a diagnostic or treatment option for heavy menstrual bleeding, as there is very little evidence to suggest that it works to investigate or treat heavy periods<sup>9</sup>.

Ultrasound scans and camera tests, with sampling of the lining of the womb (hysteroscopy and biopsy), can be used to investigate heavy periods. Medication and intrauterine systems (IUS), as well as weight loss (if appropriate) can treat heavy periods.

### Updated clinical criteria

#### Summary of intervention

<sup>9</sup> <https://www.nice.org.uk/guidance/ng88> and <https://www.nhs.uk/conditions/hysteroscopy/#alternatives-to-hysteroscopy>

Dilation and curettage (D&C) is a minor surgical procedure where the opening of the womb (cervix) is widened (dilatation) and the lining of the womb is scraped out (curettage).

#### Number of CCG interventions in 2017/18

236

#### Recommendation

D&C should not be used for diagnosis or treatment for heavy menstrual bleeding in women because it is clinically ineffective.

Ultrasound scans and camera tests with sampling of the lining of the womb (hysteroscopy and biopsy) can be used to investigate heavy periods.

Medication and intrauterine systems (IUS) can be used to treat heavy periods.

For further information, please see:

- <https://www.nice.org.uk/guidance/ng88>
- <https://www.nhs.uk/conditions/hysteroscopy/#alternatives-to-hysteroscopy>

#### Rationale for Recommendation

NICE guidelines recommend that D&C is not offered as a treatment option for heavy menstrual bleeding. There is very little evidence to suggest that D&C works to treat heavy periods and the one study identified by NICE showed the effects were only temporary. D&C should not be used to investigate heavy menstrual bleeding as hysteroscopy and biopsy work better. Complications following D&C are rare but include uterine perforation, infection, adhesions (scar tissue) inside the uterus and damage to the cervix.

#### References

1. NICE guidance: <https://www.nice.org.uk/guidance/ng88>
2. NHS advice: <https://www.nhs.uk/conditions/hysteroscopy/#alternatives-to-hysteroscopy>
3. MacKenzie IZ, Bibby JG. Critical assessment of dilatation and curettage in 1029 women. *Lancet* 1978;2(8089):566–8.
4. Ben-Baruch G, Seidman DS, Schiff E, et al. Outpatient endometrial sampling with the Pipelle curette. *Gynecologic and Obstetric Investigation* 1994;37(4):260–2.
5. Gimpelson RJ, Rappold HO. A comparative study between panoramic hysteroscopy with directed biopsies and dilatation and curettage. A review of 276 cases. *American Journal of Obstetrics and Gynecology* 1988;158(3 Pt 1):489–92.
6. Haynes PJ, Hodgson H, Anderson AB, et al. Measurement of menstrual blood loss in patients complaining of menorrhagia. *British Journal of Obstetrics and Gynaecology* 1977;84(10):763–8.

### C. Knee arthroscopy for patients with osteoarthritis

#### Updated description of the intervention

NICE recommends that arthroscopic knee washout should not be used as a treatment for patients with osteoarthritis, unless the knee locks (in which case it may be considered). More effective treatments include physiotherapy, exercise programmes like [ESCAPE pain](#), losing weight (if necessary) and pain management.<sup>10</sup> If symptoms do not resolve, knee replacement or osteotomy are effective procedures that should be considered.

### Updated clinical criteria

<p><b>Summary of intervention</b></p> <p>Arthroscopic washout of the knee is an operation where an arthroscope (camera) is inserted in to the knee along with fluid. Occasionally loose debris drains out with the fluid, or debridement, (surgical removal of damaged cartilage) is performed, but the procedure does not improve symptoms or function of the knee joint.</p>
<p><b>Number of CCG interventions in 2017/18</b></p> <p>3,437</p>
<p><b>Recommendation</b></p> <p>Arthroscopic knee washout (lavage and debridement) should not be used as a treatment for osteoarthritis because it is clinically ineffective.</p> <p>Referral for arthroscopic lavage and debridement should not be offered as part of treatment for osteoarthritis, unless the person has knee osteoarthritis with a clear history of mechanical locking.</p> <p>More effective treatment includes exercise programmes (e.g. <a href="#">ESCAPE pain</a>), losing weight (if necessary) and managing pain. Osteoarthritis is relatively common in older age groups. Where symptoms do not resolve after non-operative treatment, referral for consideration of knee replacement, or joint preserving surgery such as osteotomy is appropriate.</p> <p>For further information, please see:</p> <ul style="list-style-type: none"> <li>• <a href="https://www.nice.org.uk/guidance/ipg230/evidence/overview-pdf-492463117">https://www.nice.org.uk/guidance/ipg230/evidence/overview-pdf-492463117</a></li> <li>• <a href="https://www.nice.org.uk/guidance/ipg230/chapter/1-Guidance">https://www.nice.org.uk/guidance/ipg230/chapter/1-Guidance</a></li> <li>• <a href="https://www.nice.org.uk/donotdo/referral-for-arthroscopic-lavage-and-debridement-should-not-be-offered-as-part-of-treatment-for-osteoarthritis-unless-the-person-has-knee-osteoarthritis-with-a-clear-history-of-mechanical-locking-not">https://www.nice.org.uk/donotdo/referral-for-arthroscopic-lavage-and-debridement-should-not-be-offered-as-part-of-treatment-for-osteoarthritis-unless-the-person-has-knee-osteoarthritis-with-a-clear-history-of-mechanical-locking-not</a></li> <li>• <a href="http://www.escape-pain.org/">http://www.escape-pain.org/</a></li> </ul>
<p><b>Rationale for recommendation</b></p> <p>NICE has reviewed the evidence for how well knee washout works for people with osteoarthritis. Seven clinical trials and three case studies have shown</p>

<sup>10</sup> <https://www.nice.org.uk/guidance/ipg230/evidence/overview-pdf-492463117>; <https://www.nice.org.uk/guidance/ipg230/chapter/1-Guidance>; <https://www.nice.org.uk/donotdo/referral-for-arthroscopic-lavage-and-debridement-should-not-be-offered-as-part-of-treatment-for-osteoarthritis-unless-the-person-has-knee-osteoarthritis-with-a-clear-history-of-mechanical-locking-not>

that knee wash out for people with osteoarthritis did not reduce pain nor improve how well their knees worked. There was a small increased risk of bleeding inside the knee joint (haemarthrosis) (2%) or blood clot in the leg (deep vein thrombosis) (0.5%).

### References

1. NICE guidance: <https://www.nice.org.uk/guidance/ipg230/evidence/overview-pdf-492463117>
2. NICE guidance: <https://www.nice.org.uk/guidance/ipg230/chapter/1-Guidance>
3. NICE guidance: <https://www.nice.org.uk/donotdo/referral-for-arthroscopic-lavage-and-debridement-should-not-be-offered-as-part-of-treatment-for-osteoarthritis-unless-the-person-has-knee-osteoarthritis-with-a-clear-history-of-mechanical-locking-not>
4. British Orthopaedic Association and the Royal College of Surgeons: <https://www.rcseng.ac.uk/-/media/files/rcs/standards-and-research/commissioning/boa--painful-oa-knee-guide-final-2017.pdf>
5. Siemieniuk Reed A C, Harris Ian A, Agoritsas Thomas, Poolman Rudolf W, Brignardello-Petersen Romina, Van de Velde Stijn et al. Arthroscopic surgery for degenerative knee arthritis and meniscal tears: a clinical practice guideline BMJ 2017; 357 :j1982
6. Brignardello-Petersen R, Guyatt GH, Buchbinder R, et al Knee arthroscopy versus conservative management in patients with degenerative knee disease: a systematic review BMJ Open 2017;7:e016114. doi: 10.1136/bmjopen-2017-016114`
7. Moseley JB, O'Malley K, Petersen NJ et al. (2002) A controlled trial of arthroscopic surgery for osteoarthritis of the knee. The New England Journal of Medicine 347: 81–8.
8. Hubbard MJS. (1996) Articular debridement versus washout for degeneration of the medial femoral condyle. Journal of Bone and Joint Surgery (British) 78-B: 217–19.
9. Kalunian KC, Moreland LW, Klashman DJ et al. (2000) Visually-guided irrigation in patients with early knee osteoarthritis: a multicentre randomized controlled trial. Osteoarthritis and Cartilage 8: 412–18.
10. Chang RW, Falconer J, Stulberg SD et al. (1993) A randomized, controlled trial of arthroscopic surgery versus closed-needle joint lavage for patients with osteoarthritis of the knee. Arthritis & Rheumatism 36: 289–96.
11. Forster MC, Straw R. (2003) A prospective randomised trial comparing intra-articular Hyalgan injection and arthroscopic washout for knee osteoarthritis. The Knee 10: 291–3.
12. Jackson RW, Dieterichs C. (2003) The results of arthroscopic lavage and debridement of osteoarthritic knees based on the severity of degeneration: a 4- to 6-year symptomatic follow-up. Arthroscopy: The Journal of Arthroscopic and Related Surgery 19: 13–20.
13. Bernard J, Lemon M, Patterson MH. (2004) Arthroscopic washout of the knee – a 5-year survival analysis. The Knee 11: 233–5.

14. Harwin SF. (1999) Arthroscopic debridement for osteoarthritis of the knee: predictors of patient satisfaction. *Arthroscopy: The Journal of Arthroscopic and Related Surgery* 15: 142–6.

## D. Injections for nonspecific low back pain without sciatica

### Updated description of the intervention

NICE recommends that spinal injections should not be offered for non-specific low back pain. Alternative options like pain management and physiotherapy have been shown to work<sup>11</sup>.

### Updated clinical criteria

Summary of intervention
Spinal injections of local anaesthetic and steroid in people with non-specific low back pain without sciatica.
Number of CCG interventions in 2017/18
13,165
Recommendation
<p>Spinal injections of local anaesthetic and steroid should not be offered for patients with non-specific low back pain.</p> <p>For people with non-specific low back pain the following injections should not be offered:</p> <ul style="list-style-type: none"> <li>• facet joint injections</li> <li>• therapeutic medial branch blocks</li> <li>• intradiscal therapy</li> <li>• prolotherapy</li> <li>• Trigger point injections with any agent, including botulinum toxin</li> <li>• Epidural steroid injections for chronic low back pain or for neurogenic claudication in patients with central spinal canal stenosis</li> <li>• Any other spinal injections not specifically covered above</li> </ul> <p>Radiofrequency denervation can be offered according to NICE guideline (NG59) if all non-surgical and alternative treatments have been tried and there is moderate to severe chronic pain that has improved in response to diagnostic medical branch block.</p> <p>Epidurals (local anaesthetic and steroid) should be considered in patients who have acute and severe lumbar radiculopathy at time of referral.</p> <p>Alternative and less invasive options have been shown to work e.g. exercise programmes, behavioural therapy, and attending a specialised pain clinic. Alternative options are suggested in line with the National Back Pain Pathway.</p>

<sup>11</sup> <https://www.nice.org.uk/guidance/ng59>

For further information, please see: <https://www.nice.org.uk/guidance/ng59>

### Rationale for recommendation

NICE guidelines recommend that spinal injections should not be offered for non-specific low back pain.

Radiofrequency denervation (to destroy the nerves that supply the painful facet joint in the spine) can be considered in some cases as per NICE guidance.

Exclusion criteria for the NICE (NG59) include:

Conditions of a non-mechanical nature, including;

- Inflammatory causes of back pain (for example, ankylosing spondylitis or diseases of the viscera)
- Serious spinal pathology (for example, neoplasms, infections or osteoporotic collapse)

Neurological disorders (including cauda equina syndrome or mononeuritis)

Adolescent scoliosis

Not covered were conditions with a select and uniform pathology of a mechanical nature (e.g. spondylolisthesis, scoliosis, vertebral fracture or congenital disease)

Other agreed exclusions by the GDG are: Pregnancy-related back pain, Sacroiliac joint dysfunction, Adjacent-segment disease, Failed back surgery syndrome, Spondylolisthesis and Osteoarthritis.

NICE recommends the following approach for non-surgical invasive treatments for low back pain and sciatica in over 16s

### Spinal injections

1.3.1 Do not offer spinal injections for managing nonspecific low back pain.

### Radiofrequency denervation

1.3.2 Consider referral for assessment for radiofrequency denervation for people with non-specific low back pain when:

non-surgical treatment has not worked for them and the main source of pain is thought to come from structures supplied by the medial branch nerve and they have moderate or severe levels of localised back pain (rated as 5 or more on a visual analogue scale, or equivalent) at the time of referral.

1.3.3 Only perform radiofrequency denervation in people with non-specific low back pain after a positive response to a diagnostic medial branch block.

1.3.4 Do not offer imaging for people with non-specific low back pain with specific facet joint pain as a prerequisite for radiofrequency denervation.

### References

1. NICE guidance: <https://www.nice.org.uk/guidance/ng59>,
2. United Kingdom Spine Societies Board: <https://www.ukssb.com/improving-spinal-care-project>
3. Benyamin RM, Manchikanti L, Parr AT, Diwan S, Singh V, Falco FJ, et al. The effectiveness of lumbar interlaminar epidural injections in managing chronic low back and lower extremity pain. *Pain Physician*. 2012 Jul-Aug;15(4):E363-404.

4. Choi HJ, Hahn S, Kim CH, Jang BH, Park S, Lee SM, et al. Epidural steroid injection therapy for low back pain: a meta-analysis. *Int J Technol Assess Health Care*. 2013 Jul;29(3):244-53.
5. Cohen SP, Bicket MC, Jamison D, Wilkinson I, Rathmell JP. Epidural steroids: a comprehensive, evidence-based review. *Reg Anesth Pain Med*. 2013 May- Jun;38(3):175-200.
6. Royal College of Anaesthetists: <https://www.rcoa.ac.uk/document-store/core-standards-pain-management-services-the-uk>

## E. Breast reduction

### Updated description of the intervention

The evidence highlights that breast reduction is only successful in specific circumstances and the procedure can lead to complications - for example not being able to breast feed permanently. However in some cases breast reduction surgery is necessary where large breasts impact on day to day life, for example ability to drive a car. Therefore, breast reduction should only be undertaken under specific criteria. Wearing a professionally fitted bra, losing weight (if necessary), managing pain and physiotherapy often work well to help with symptoms like back pain from large breasts.

### Updated clinical criteria

#### Summary of intervention

Breast reduction surgery is a procedure used to treat women with breast hyperplasia (enlargement), where breasts are large enough to cause problems like shoulder girdle dysfunction, intertrigo and adverse effects to quality of life.

#### Number of CCG interventions in 2017/18

2,388

#### Recommendation

The NHS will only provide breast reduction for women if **all** the following criteria are met:

- The woman has received a full package of supportive care from their GP such as advice on weight loss and managing pain.
- In cases of thoracic/ shoulder girdle discomfort, a physiotherapy assessment has been provided
- Breast size results in functional symptoms that require other treatments/interventions (e.g. intractable candidal intertrigo; thoracic backache/kyphosis where a professionally fitted bra has not helped with backache, soft tissue indentations at site of bra straps).
- Breast reduction planned to be 500gms or more per breast or at least 4 cup sizes.
- Body mass index (BMI) is <27 and stable for at least twelve months.
- Woman must be provided with written information to allow her to balance the risks and benefits of breast surgery.

- Women should be informed that smoking increases complications following breast reduction surgery and should be advised to stop smoking.
- Women should be informed that breast surgery for hypermastia can cause permanent loss of lactation.

Unilateral breast reduction is considered for asymmetric breasts as opposed to breast augmentation if there is an impact on health as per the criteria above. Surgery will not be funded for cosmetic reasons. Surgery can be approved for a difference of 150 - 200gms size as measured by a specialist. The BMI needs to be <27 and stable for at least twelve months.

Resection weights, for bilateral or unilateral (both breasts or one breast) breast reduction should be recorded for audit purposes.

This recommendation does not apply to therapeutic mammoplasty for breast cancer treatment or contralateral (other side) surgery following breast cancer surgery, and local policies should be adhered to. The Association of Breast Surgery support contralateral surgery to improve cosmesis as part of the reconstruction process following breast cancer treatment.

**Gynaecomastia:** Surgery for gynaecomastia is not routinely funded by the NHS. This recommendation does not cover surgery for gynaecomastia caused by medical treatments such as treatment for prostate cancer.

### Rationale for recommendation

One systematic review and three non-randomized studies regarding breast reduction surgery for hypermastia were identified and showed that surgery is beneficial in patients with specific symptoms. Physical and psychological improvements, such as reduced pain, increased quality of life and less anxiety and depression were found for women with hypermastia following breast reduction surgery.

Breast reduction surgery for hypermastia can cause permanent loss of lactation function of breasts, as well as decreased areolar sensation, bleeding, bruising, and scarring and often alternative approaches (e.g. weight loss or a professionally fitted bra) work just as well as surgery to reduce symptoms. For women who are severely affected by complications of hypermastia and for whom alternative approaches have not helped, surgery can be offered. The aim of surgery is not cosmetic, it is to reduce symptoms (e.g. back ache).

### References

1. An investigation into the relationship between breast size, bra size and mechanical back pain. British School of Osteopathy (2010). Pages 13 & 14
2. Royal College of Surgeons – <https://www.rcseng.ac.uk/-/media/files/rcs/library-and-publications/non-journal-publications/breast-reduction--commissioning-guide.pdf>
3. Greenbaum, a. R., Heslop, T., Morris, J., & Dunn, K. W. (2003). An investigation of the suitability of bra fit in women referred for reduction mammoplasty. British Journal of Plastic Surgery, 56(3), 230–236.

4. Wood, K., Cameron, M., & Fitzgerald, K. (2008). Breast size, bra fit and thoracic pain in young women: a correlational study. *Chiropractic & Osteopathy*, 16(1), 1-7.
5. Singh KA, Losken A. Additional benefits of reduction mammoplasty: a systematic review of the literature. *Plast Reconstr Surg*. 2012 Mar;129(3):562-70. PubMed: PM22090252
7. Strong B, Hall-Findlay EJ. How Does Volume of Resection Relate to Symptom Relief for Reduction Mammoplasty Patients? *Ann Plast Surg*. 2014 Apr 10. PubMed: PM24727444
8. Valtonen JP, Setala LP, Mustonen PK, Blom M. Can the efficacy of reduction mammoplasty be predicted? The applicability and predictive value of breast-related symptoms questionnaire in measuring breast-related symptoms pre- and postoperatively. *J Plast Reconstr Aesthet Surg*. 2014 May;67(5):676-81. PubMed: PM24508223
9. Foreman KB, Dibble LE, Droge J, Carson R, Rockwell WB. The impact of breast reduction surgery on low-back compressive forces and function in individuals with macromastia. *Plast Reconstr Surg*. 2009 Nov;124(5):1393-9. PubMed: PM20009823
10. Shah R, Al-Ajam Y, Stott D, Kang N. Obesity in mammoplasty: a study of complications following breast reduction. *J Plast Reconstr Aesthet Surg*. 2011 Apr;64(4):508-14. doi: 10.1016/j.bjps.2010.07.001. Epub 2010 Aug 3. PubMed PMID: 20682461.
11. Oo M, Wang Z, Sakakibara T, Kasai Y. Relationship Between Brassiere Cup Size and Shoulder-Neck Pain in Women. *The Open Orthopaedics Journal*. 2012;6:140-142. doi:10.2174/1874325001206010140.
12. <https://www.nhs.uk/conditions/breast-reduction-on-the-nhs/>
13. [Plast Reconstr Surg. 2011 Nov;128\(5\):395e-402e. doi:10.1097/PRS.0b013e3182284c05.The impact of obesity on breast surgery complications.Chen CL\(1\), Shore AD, Johns R, Clark JM, Manahan M, Makary MA](#)

## F. Removal of benign skin lesions

### Updated description of the intervention

Removal of benign skin lesions cannot be offered for cosmetic reasons. It should only be offered in situations where the lesion is causing symptoms according to the criteria outlined below. Risks from the procedure can include bleeding, pain, infection, and scarring.

### Updated clinical criteria

#### Summary of intervention

Removal of benign skin lesions means treating asymptomatic lumps, bumps or tags on the skin that are not suspicious of cancer. Treatment carries a small risk of infection, bleeding or scarring and is not usually offered by the NHS if it is just to improve appearance. In certain cases, treatment (surgical excision or cryotherapy) may be offered if certain criteria are met. A patient with a skin or subcutaneous lesion that has features suspicious of malignancy must be treated or referred

according to NICE skin cancer guidelines. This policy does not refer to pre-malignant lesions and other lesions with potential to cause harm.

#### Number of CCG interventions in 2017/18

116,255

#### Recommendation

This policy refers to the following benign lesions when there is diagnostic certainty and they do not meet the criteria listed below:

- benign moles (excluding large congenital naevi)
- solar comedones
- corn/callous
- dermatofibroma
- lipomas
- milia
- molluscum contagiosum (non-genital)
- epidermoid & pilar cysts (sometimes incorrectly called sebaceous cysts)
- seborrhoeic keratoses (basal cell papillomata)
- skin tags (fibroepithelial polyps) including anal tags
- spider naevi (telangiectasia)
- non-genital viral warts in immunocompetent patients
- xanthelasmata
- neurofibromata

The benign skin lesions, which are listed above, must meet at least ONE of the following criteria to be removed:

- The lesion is unavoidably and significantly traumatised on a regular basis with evidence of this causing regular bleeding or resulting in infections such that the patient requires 2 or more courses of antibiotics (oral or intravenous) per year
- There is repeated infection requiring 2 or more antibiotics per year
- The lesion bleeds in the course of normal everyday activity
- The lesion causes regular pain
- The lesion is obstructing an orifice or impairing field vision
- The lesion significantly impacts on function e.g. restricts joint movement
- The lesion causes pressure symptoms e.g. on nerve or tissue
- If left untreated, more invasive intervention would be required for removal
- Facial viral warts
- Facial spider naevi in children causing significant psychological impact
- Lipomas on the body > 5cms, or in a sub-facial position, with rapid growth and/or pain. These should be referred to Sarcoma clinic.

The following are *outside* the scope of this policy recommendation:

- Lesions that are suspicious of malignancy should be treated or referred according to NICE skin cancer guidelines.
- Any lesion where there is diagnostic uncertainty, pre-malignant lesions (actinic keratoses, Bowen disease) or lesions with pre-malignant potential should be referred or, where appropriate, treated in primary care.
- Removal of lesions other than those listed above.

Referral to appropriate speciality service (eg dermatology or plastic surgery):

- The decision as to whether a patient meets the criteria is primarily with the referring clinician. If such lesions are referred, then the referrer should state that this policy has been considered and why the patient meets the criteria.
- This policy applies to all providers, including general practitioners (GPs), GPs with enhanced role (GPwer), independent providers, and community or intermediate services.

For further information, please see:

- <https://www.nice.org.uk/guidance/csg8>
- <https://www.nice.org.uk/guidance/ng12>

### Rationale for recommendation

There is little evidence to suggest that removing benign skin lesions to improve appearance is beneficial. Risks of this procedure include bleeding, pain, infection and scarring. Though in certain specific cases as outlined by the criteria above, there are benefits for removing skin lesions, for example, avoidance of pain and allowing normal functioning.

### References

1. Higgins JC, Maher MH, Douglas MS. Diagnosing Common Benign Skin Tumors. *Am Fam Physician*. 2015 Oct 1;92(7):601-7. PubMed PMID: 26447443.
2. Tan E, Levell NJ, Garioch JJ. The effect of a dermatology restricted-referral list upon the volume of referrals. *Clin Exp Dermatol*. 2007 Jan;32(1):114-5. PubMed PMID: 17305918.

## G. Grommets for Glue Ear in Children

### Updated description of the intervention

Evidence suggests that grommets only offer a short-term hearing improvement in children with glue ear who have no other serious medical problems or disabilities. They should be offered in cases that have a history of persistent (at least 3 months) bilateral, hearing loss as defined by the NICE guidance. Hearing aids can also be offered as an alternative to surgery<sup>12</sup>.

### Updated clinical criteria

#### Summary of intervention

This is a surgical procedure to insert tiny tubes (grommets) into the eardrum as a treatment for fluid build up (glue ear) when it is affecting hearing in children.

Glue ear is a very common childhood problem (4 out of 5 children will have had an episode by age 10), and in most cases it clears up without treatment within a few weeks. Common symptoms can include earache and a reduction in hearing. Often, when the hearing loss is affecting both ears it can cause language, educational and behavioural problems.

<sup>12</sup> <https://www.nice.org.uk/Guidance/CG60>

Please note this guidance only relates to children with Glue Ear (Otitis Media with Effusion) and SHOULD NOT be applied to other clinical conditions where grommet insertion should continue to be normally funded, these include:

- Recurrent acute otitis media
- Atrophic tympanic membranes
- Access to middle ear for transtympanic instillation of medication

Investigation of unilateral glue ear in adults

#### Number of CCG interventions in 2017/18

8,669

#### Recommendation

The NHS should only commission this surgery for the treatment of glue ear in children when the criteria set out by the NICE guidelines are met:

- All children must have had specialist audiology and ENT assessment.
- Persistent bilateral otitis media with effusion over a period of 3 months.
- Hearing level in the better ear of 25-30dbHL or worse averaged at 0.5, 1, 2, & 4kHz
- Exceptionally, healthcare professionals should consider surgical intervention in children with persistent bilateral OME with a hearing loss less than 25-30dbHL where the impact of the hearing loss on a child's developmental, social or educational status is judged to be significant.
- Healthcare professionals should also consider surgical intervention in children who cannot undergo standard assessment of hearing thresholds where there is clinical and tympanographic evidence of persistent glue ear and where the impact of the hearing loss on a child's developmental, social or educational status is judged to be significant.
- The guidance is different for children with Down's Syndrome and Cleft Palate, these children may be offered grommets after a specialist MDT assessment in line with NICE guidance.
- It is also good practice to ensure glue ear has not resolved once a date of surgery has been agreed, with tympanometry as a minimum.

For further information, please see: <https://www.nice.org.uk/Guidance/CG60>.

The risks to surgery are generally low, but the most common is persistent ear discharge (10-20%) and this can require treatment with antibiotic eardrops and water precautions. In rare cases (1-2%) a persistent hole in the eardrum may remain, and if this causes problems with recurrent infection, surgical repair may be required (however this is not normally done until around 8-10 years of age).

#### Rationale for recommendation

In most cases glue ear will improve by itself without surgery. During a period of monitoring of the condition a balloon device (e.g. Otovent) can be used by the child if tolerated, this is designed to improve the function of the ventilation tube that connects the ear to the nose. In children with persistent glue ear, a hearing aid is another suitable alternative to surgery. Evidence suggests that grommets only offer

a short-term hearing improvement in children with no other serious medical problems or disabilities.

The NHS should only commission this surgery when the NICE criteria are met, as performing the surgery outside of these criteria is unlikely to derive any clinical benefit.

#### References

1. NICE guidance: <https://www.nice.org.uk/Guidance/CG60>
2. Browning, G; Rovers, M; Williamson, I; Lous, J; Burton, MJ. Grommets (ventilation tubes) for hearing loss associated with otitis media with effusion in children. Cochrane Database of Systematic Reviews 2010, Issue 10. Art. No.: CD001801. DOI: 10.1002/14651858.CD001801.pub3

## H. Tonsillectomy for Recurrent Tonsillitis

### Updated description of the intervention

Recurrent sore throats are a very common condition that present a considerable health burden. In most cases they can be treated with conservative measures. In some cases, where there are recurrent, documented episodes of acute tonsillitis that are disabling to normal function, then tonsillectomy is beneficial, but it should only be offered when the frequency of episodes set out by the Scottish Intercollegiate Guidelines Network criteria are met.

### Updated clinical criteria

#### Summary of intervention

This guidance relates to surgical procedures to remove the tonsils as a treatment for recurrent sore throats in adults and children.

Recurring sore throats are a very common condition that presents a large burden on healthcare; they can also impact on a person's ability to work or attend school. It must be recognised however, that not all sore throats are due to tonsillitis and they can be caused by other infections of the throat. In these cases, removing the tonsils will not improve symptoms.

#### Number of CCG interventions in 2017/18

32,238

#### Recommendation

The NHS should only commission this surgery for treatment of recurrent severe episodes of sore throat when the following criteria are met, as set out by the SIGN guidance and supported by ENT UK commissioning guidance:

- Sore throats are due to acute tonsillitis AND
- The episodes are disabling and prevent normal functioning AND
- Seven or more, documented, clinically significant, adequately treated sore throats in the preceding year OR
- Five or more such episodes in each of the preceding two years OR

- Three or more such episodes in each of the preceding three years.

There are a number of medical conditions where episodes of tonsillitis can be damaging to health or tonsillectomy is required as part of the on-going management. In these instances tonsillectomy may be considered beneficial at a lower threshold than this guidance after specialist assessment:

- Acute and chronic renal disease resulting from acute bacterial tonsillitis.
- As part of the treatment of severe guttate psoriasis.
- Metabolic disorders where periods of reduced oral intake could be dangerous to health.
- PFAPA (Periodic fever, Aphthous stomatitis, Pharyngitis, Cervical adenitis)
- Severe immune deficiency that would make episodes of recurrent tonsillitis dangerous

Further information on the Scottish Intercollegiate Guidelines Network guidance can be found here: <http://www.sign.ac.uk/assets/sign117.pdf>

Please note this guidance only relates to patients with recurrent tonsillitis. This guidance should not be applied to other conditions where tonsillectomy should continue to be funded, these include:

- Obstructive Sleep Apnoea / Sleep disordered breathing in Children
- Suspected Cancer (e.g. asymmetry of tonsils)
- Recurrent Quinsy (abscess next to tonsil)
- Emergency Presentations (e.g. treatment of parapharyngeal abscess)

It is important to note that national randomised control trial is underway comparing surgery versus conservative management for recurrent tonsillitis in adults in underway which may warrant review of this guidance in the near future.

### **Rationale for recommendation**

Recurrent sore throats are a very common condition that presents a considerable health burden. In most cases they can be treated with conservative measures. In some cases, where there are recurrent, documented episodes of acute tonsillitis that are disabling to normal function, then tonsillectomy is beneficial, but it should only be offered when the frequency of episodes set out by the Scottish Intercollegiate Guidelines Network criteria are met.

The surgery carries a small risk of bleeding requiring readmission to hospital (3.5%). A previous national audit quoted a 0.9% risk of requiring emergency surgery to treat bleeding after surgery but in a more recent study of 267, 159 tonsillectomies, 1.88% of patients required a return to theatre. Pain after surgery can be severe (especially in adults) for up to two weeks after surgery; this requires regular painkillers and can cause temporary difficulty swallowing. In addition to bleeding; pain or infection after surgery can require readmission to hospital for treatment. The Getting it Right First Time ENT report is due late 2018 and will present updated figures on readmission rates in relation to tonsillectomy.

There is no alternative treatment for recurrent sore throats that is known to be beneficial, however sometimes symptoms improve with a period of observation.

**References**

1. Rubie I, Haighton C, O'Hara J, Rousseau N, Steen N, Stocken DD, Sullivan F, Vale L, Wilkes S, Wilson J. The National randomised controlled Trial of Tonsillectomy IN Adults (NATTINA): a clinical and cost-effectiveness study: study protocol for a randomised control trial. *Trials*. 2015 Jun 6;16:263. <https://www.ncbi.nlm.nih.gov/pubmed/26047934>
2. <http://www.sign.ac.uk/assets/sign117.pdf>
3. Osbourne MS, Clark MPA. The surgical arrest of post-tonsillectomy haemorrhage: Hospital Episode Statistics 12 years on. *Annals RCS*. 2018. May (100) 5: 406-408

**I. Haemorrhoid surgery****Updated description of the intervention**

Numerous interventions exist for the management of haemorrhoids (piles). The evidence recommends that surgical treatment should only be considered for haemorrhoids that keep coming back after treatment or for haemorrhoids that are significantly affecting daily life. Changes to the diet like eating more fibre and drinking more water can often help with haemorrhoids. Treatments that can be done in clinic like rubber band ligation, may be effective especially for less severe haemorrhoids.

**Updated clinical criteria****Summary of intervention**

This procedure involves surgery for haemorrhoids (piles).

**Number of CCG interventions in 2017/18**

8,474

**Recommendation**

Often haemorrhoids (especially early stage haemorrhoids) can be treated by simple measures such as eating more fibre or drinking more water. If these treatments are unsuccessful many patients will respond to outpatient treatment in the form of banding or perhaps injection.

Surgical treatment should only be considered for those that do not respond to these non-operative measures or if the haemorrhoids are more severe, specifically:

- Recurrent grade 3 or grade 4 combined internal/external haemorrhoids with persistent pain or bleeding; or
- Irreducible and large external haemorrhoids

In cases where there is significant rectal bleeding the patient should be examined internally by a specialist.

**Rationale for recommendation**

Surgery should be performed, according to patient choice and only in cases of persistent grade 1 (rare) or 2 haemorrhoids that have not improved with dietary

changes, banding or perhaps in certain cases injection, and recurrent grade 3 and 4 haemorrhoids and those with a symptomatic external component.

Haemorrhoid surgery can lead to complications. Pain and bleeding are common and pain may persist for several weeks. Urinary retention can occasionally occur and may require catheter insertion. Infection, iatrogenic fissuring (tear or cut in the anus), stenosis and incontinence (lack of control over bowel motions) occur more infrequently.

### References

1. Watson AJM, Bruhn H, MacLeod K, et al. A pragmatic, multicentre, randomised controlled trial comparing stapled haemorrhoidopexy to traditional excisional surgery for haemorrhoidal disease (eTHoS): study protocol for a randomised controlled trial. *Trials*. 2014;15:439. doi:10.1186/1745-6215-15-439.
2. Watson AJM, Hudson J, Wood J, et al. Comparison of stapled haemorrhoidopexy with traditional excisional surgery for haemorrhoidal disease (eTHoS): a pragmatic, multicentre, randomised controlled trial. *Lancet (London, England)*. 2016;388(10058):2375-2385. doi:10.1016/S0140-6736(16)31803-7.
3. Brown SR. Haemorrhoids: an update on management. *Therapeutic Advances in Chronic Disease*. 2017;8(10):141-147. doi:10.1177/2040622317713957.
4. NHS website: <https://www.nhs.uk/conditions/piles-haemorrhoids/>
5. Royal College of Surgeons: [https://www.rcseng.ac.uk/-/media/files/rcs/standards-and-research/commissioning/rcsacpqbirectalbleeding2017documentfinal\\_jan18.pdf](https://www.rcseng.ac.uk/-/media/files/rcs/standards-and-research/commissioning/rcsacpqbirectalbleeding2017documentfinal_jan18.pdf)
6. Health Technol Assess. 2016 Nov;20(88):1-150. The HubBLLe Trial: haemorrhoidal artery ligation (HAL) versus rubber band ligation (RBL) for symptomatic second- and third-degree haemorrhoids: a multicentre randomised controlled trial and health-economic evaluation. Brown S et al.

## **J. Hysterectomy for heavy menstrual bleeding**

### **Updated description of the intervention**

NICE recommends that hysterectomy should not be used as a first-line treatment solely for heavy menstrual bleeding (HMB).<sup>13</sup> Heavy periods can be reduced by using medicines or intrauterine systems (IUS) or losing weight (if necessary).

### **Updated clinical criteria**

#### **Summary of intervention**

Hysterectomy is the surgical removal of the uterus.

#### **Number of CCG interventions in 2017/18**

<sup>13</sup> <https://www.nice.org.uk/guidance/ng88>

27,660

### Recommendation

Based on NICE guidelines [[Heavy menstrual bleeding: assessment and management \[NG88\] Published date: March 2018](#)], hysterectomy should not be used as a first-line treatment solely for heavy menstrual bleeding.

It is important that healthcare professionals understand what matters most to each woman and support her personal priorities and choices.

Hysterectomy should be considered only when: other treatment options have failed, are contradicted; there is a wish for amenorrhoea (no periods); the woman (who has been fully informed) requests it; the woman no longer wishes to retain her uterus and fertility.

#### 1.13.1.1.1 *NICE guideline NG88 1.5 Management of HMB*

1.5.1 When agreeing treatment options for HMB with women, take into account: the woman's preferences, any comorbidities, the presence or absence of fibroids (including size, number and location), polyps, endometrial pathology or adenomyosis, other symptoms such as pressure and pain.

#### 1.13.1.1.2 *Treatments for women with no identified pathology, fibroids less than 3 cm in diameter, or suspected or diagnosed adenomyosis*

1.5.2 Consider an LNG-IUS (levonorgestrel-releasing intrauterine system) as the first treatment for HMB in women with: no identified pathology or fibroids less than 3 cm in diameter, which are not causing distortion of the uterine cavity or suspected or diagnosed adenomyosis.

1.5.3 If a woman with HMB declines an LNG-IUS or it is not suitable, consider the following pharmacological treatments: non-hormonal: tranexamic acid, NSAIDs (non-steroidal anti-inflammatory drugs), hormonal: combined hormonal contraception, cyclical oral progestogens.

1.5.4 Be aware that progestogen-only contraception may suppress menstruation, which could be beneficial to women with HMB.

1.5.5 If treatment is unsuccessful, the woman declines pharmacological treatment, or symptoms are severe, consider referral to specialist care for: investigations to diagnose the cause of HMB, if needed, taking into account any investigations the woman has already had and alternative treatment choices, including: pharmacological options not already tried (see recommendations 1.5.2 and 1.5.3), surgical options: second-generation endometrial ablation, hysterectomy.

1.5.6 For women with submucosal fibroids, consider hysteroscopic removal.

**1.13.1.1.3** *Treatments for women with fibroids of 3 cm or more in diameter*

1.5.7 Consider referring women to specialist care to undertake additional investigations and discuss treatment options for fibroids of 3 cm or more in diameter.

1.5.8 If pharmacological treatment is needed while investigations and definitive treatment are being organised, offer tranexamic acid and/or NSAIDs.

1.5.9 Advise women to continue using NSAIDs and/or tranexamic acid for as long as they are found to be beneficial.

1.5.10 For women with fibroids of 3 cm or more in diameter, take into account the size, location and number of fibroids, and the severity of the symptoms and consider the following treatments: pharmacological: non-hormonal: tranexamic acid, NSAIDs, hormonal: LNG-IUS, combined hormonal contraception, cyclical oral progestogens, uterine artery embolization, surgical: myomectomy, hysterectomy.

1.5.12 Be aware that the effectiveness of pharmacological treatments for HMB may be limited in women with fibroids that are substantially greater than 3 cm in diameter.

1.5.13 Prior to scheduling of uterine artery embolisation or myomectomy, the woman's uterus and fibroid(s) should be assessed by ultrasound. If further information about fibroid position, size, number and vascularity is needed, MRI should be considered. [2007]

1.5.14 Consider second-generation endometrial ablation as a treatment option for women with HMB and fibroids of 3 cm or more in diameter who meet the criteria specified in the manufacturers' instructions.

1.5.15 If treatment is unsuccessful: consider further investigations to reassess the cause of HMB, taking into account the results of previous investigations and offer alternative treatment with a choice of the options described in recommendation 1.5.10.

1.5.16 Pretreatment with a gonadotrophin-releasing hormone analogue before hysterectomy and myomectomy should be considered if uterine fibroids are causing an enlarged or distorted uterus.

For further information, please see:

- <https://www.nice.org.uk/guidance/ng88>.
- <https://www.nhs.uk/conditions/heavy-periods/#Causes>

### Rationale for recommendation

NICE's Guideline Development Group considered the evidence (including 2 reviews, four randomised control trials and one cohort study comparing hysterectomy with other treatments) as well as the views of patients and the public and concluded that hysterectomy should not routinely be offered as first line treatment for heavy menstrual bleeding. The Group placed a high value on the need for education and information provision for women with heavy menstrual bleeding.

Complications following hysterectomy are usually rare but infection occurs commonly. Less common complications include: intra-operative haemorrhage; damage to other abdominal organs, such as the urinary tract or bowel; urinary dysfunction –frequent passing of urine and incontinence. Rare complications include thrombosis (DVT and clot on the lung) and very rare complications include death. Complications are more likely when hysterectomy is performed in the presence of fibroids (non-cancerous growths in the uterus). There is a risk of possible loss of ovarian function and its consequences, even if their ovaries are retained during hysterectomy. If oophorectomy (removal of the ovaries) is performed at the time of hysterectomy, menopausal-like symptoms occur.

#### References

1. NICE guidance: <https://www.nice.org.uk/guidance/ng88>.
2. NHS website: <https://www.nhs.uk/conditions/heavy-periods/#Causes>
3. Hurskainen R, Teperi J, Rissanen P, et al. Clinical outcomes and costs with the levonorgestrel-releasing intrauterine system or hysterectomy for treatment of menorrhagia: randomized trial 5-year follow-up. *JAMA: the journal of the American Medical Association* 2004;291(12):1456–63.
4. Learman LA, Summitt Jr RL, Varner RE, et al. Hysterectomy versus expanded medical treatment for abnormal uterine bleeding: Clinical outcomes in the medicine or surgery trial. *Obstetrics and Gynecology* 2004;103(5 1):824–33.
5. Zupi E, Zullo F, Marconi D, et al. Hysteroscopic endometrial resection versus laparoscopic supracervical hysterectomy for menorrhagia: a prospective randomized trial. *American Journal of Obstetrics and Gynecology* 2003;188(1):7–12.
6. Lethaby A, Hickey M, Garry R. Endometrial destruction techniques for heavy menstrual bleeding. *Cochrane Database Syst Rev.* 2005 Oct 19;(4):CD001501. Review. Update in: *Cochrane Database Syst Rev.* 2009;(4):CD001501. PubMed PMID: 16235284.
7. Hehenkamp WJ, Volkers NA, Donderwinkel PF, et al. Uterine artery embolization versus hysterectomy in the treatment of symptomatic uterine fibroids (EMMY trial): peri- and postprocedural results from a randomized controlled trial. *American Journal of Obstetrics and Gynecology* 2005;193(5):1618–29.
8. Pinto I, Chimeno P, Romo A, et al. Uterine fibroids: uterine artery embolization versus abdominal hysterectomy for treatment – a prospective, randomized, and controlled clinical trial. *Radiology* 2003;226(2):425–31.

## K. Chalazia removal

### Updated description of the intervention

The evidence shows that alternative treatment options (warm compresses, drops or ointment, steroid injection) or a “watch and wait” approach will lead to resolution of many chalazia without the risks of surgery.

### Updated clinical criteria

Summary of intervention
This procedure involves incision and curettage (scraping away) of the contents of the chalazion. Chalazia (meibomian cysts) are benign lesions on the eyelids due to blockage and swelling of an oil gland that normally change size over a few weeks. Many but not all resolve within six months with regular application of warm compresses and massage.
Number of CCG interventions in 2017/18
6,026
Recommendation
<p>Incision and curettage (or triamcinolone injection for suitable candidates) of chalazia should only be undertaken if at least <b>one</b> of the following criteria have been met:</p> <ul style="list-style-type: none"> <li>• Has been present for more than 6 months and has been managed conservatively with warm compresses, lid cleaning and massage for 4 weeks</li> <li>• Interferes significantly with vision</li> <li>• Interferes with the protection of the eye by the eyelid due to altered lid closure or lid anatomy</li> <li>• Is a source of infection that has required medical attention twice or more within a six month time frame</li> <li>• Is a source of infection causing an abscess which requires drainage</li> <li>• If malignancy (cancer) is suspected eg. Madarosis/recurrence/other suspicious features in which case the lesion should be removed and sent for histology as for all suspicious lesions</li> </ul>
Rationale for recommendation
<p>NICE recommend that warm compresses and lid massage alone are sufficient first line treatment for chalazia. If infection is suspected a drop or ointment containing an antibiotic (e.g. Chloramphenicol) should be added in addition to warm compresses. Only if there is spreading lid and facial cellulitis should a short course of oral antibiotics (e.g. co-amoxiclav) be used.</p> <p>Where there is significant inflammation of the chalazion a drop or ointment containing an antibiotic and steroid can be used along with other measures such as warm compresses. However, all use of topical steroids around the eye does</p>

carry the risk of raised intraocular pressure or cataract although this is very low with courses of less than 2 weeks.

Many chalazia, especially those that present acutely, resolve within six months and will not cause any harm however there are a small number which are persistent, very large, or can cause other problems such as distortion of vision.

In these cases surgery can remove the contents from a chalazion. However all surgery carries risks. Most people will experience some discomfort, swelling and often bruising of the eyelids and the cyst can take a few weeks to disappear even after successful surgery. Surgery also carries a small risk of infection, bleeding and scarring, and there is a remote but serious risk to the eye and vision from any procedure on the eyelids. Lastly in a proportion of successful procedures the chalazion can come back. The alternative option of an injection of a steroid (triamcinolone) also carries a small risk of serious complications such as raised eye pressure, eye perforation or bleeding.

Some trials comparing the two treatments suggest that using a single triamcinolone acetonide injection followed by lid massage is almost as effective as incision and curettage in the treatment of chalazia and with similar patient satisfaction but less pain and patient inconvenience. However this is controversial and other studies show that steroid injection is less effective than surgery. Therefore both options can be considered for suitable patients.

#### References

1. NICE clinical knowledge summaries, <https://cks.nice.org.uk/meibomian-cyst-chalazion>
2. Moorfield's Eye Hospital Patient Information, <https://www.moorfields.nhs.uk/sites/default/files/chalazion-adult.pdf>
3. Wu AY, Gervasio KA, Gergoudis KN, Wei C, Oestreicher JH, Harvey JT. Conservative therapy for chalazia: is it really effective? *Acta Ophthalmol.* 2018 Jan 16. doi: 10.1111/aos.13675. [Epub ahead of print] PubMed PMID: 29338124.
4. Goawalla A, Lee V. A prospective randomized treatment study comparing three treatment options for chalazia: triamcinolone acetonide injections, incision and curettage and treatment with hot compresses. *Clin Exp Ophthalmol.* 2007 Nov;35(8):706-12. PubMed PMID: 17997772.
5. Watson P, Austin DJ. Treatment of chalazions with injection of a steroid Suspension. *British Journal of Ophthalmology*, 1984, 68, 833-835.
6. Ben Simon, G.J., Huang, L., Nakra, T. et al. Intralesional triamcinolone acetonide injection for primary and recurrent chalazia (is it really effective?) . *Ophthalmology.* 2005; 112: 913–917.
7. Papalkar D, Francis IC. Injections for Chalazia? *Ophthalmology* 2006; 113:355–356. Incision and curettage vs steroid injection for the treatment of chalazia: a metaanalysis. Aycinena A, Achrion A et al. *Ophthalmic Plastic and reconstructive surgery.* 2016;32:220-224.
9. McStay. Stye and Chalazion. *BMJ Best Practice* <https://bestpractice.bmj.com/topics/en-gb/214> (accessed 18/10/18)

## L. Arthroscopic shoulder decompression for subacromial shoulder pain

### Updated description of the intervention

Recent research has indicated that in patients with pure subacromial impingement (with no other associated diagnoses such as rotator cuff tears, calcific tendinopathy and acromio-clavicular joint pain), non-operative management with a combination of exercise and physiotherapy is effective in the majority of cases.

Patients suffering with persistent symptoms, despite appropriate non-operative management, should be given the option to choose decompression surgery.

Treating clinicians and surgeons should refer to the 2015 BESS/BOA/NICE commissioning guidelines (guideline update due in 2018/19) for details of appropriate treatment of these patients. [https://www.boa.ac.uk/wp-content/uploads/2014/08/Subacromial-Shoulder-Commissioning-Guide\\_final.pdf](https://www.boa.ac.uk/wp-content/uploads/2014/08/Subacromial-Shoulder-Commissioning-Guide_final.pdf)

In order to facilitate non-operative treatment in primary and intermediate care, BESS and Getting It Right First Time programme have produced patient exercise rehab videos and booklets for GPs and patients to use.

<http://www.bess.org.uk/index.php/public-area/shpi-videos>

### Updated clinical criteria

<p><b>Summary of procedure</b></p> <p>Arthroscopic sub-acromial decompression is a surgical procedure that involves decompressing the sub-acromial space by removing bone spurs and soft tissue arthroscopically.</p>
<p><b>Number of CCG interventions in 2017/18</b></p> <p>13,930</p>
<p><b>Recommendation</b></p> <p>Arthroscopic subacromial decompression for pure subacromial shoulder impingement should only offered in appropriate cases. To be clear, 'pure subacromial shoulder impingement' means subacromial pain not caused by associated diagnoses such as rotator cuff tears, acromio-clavicular joint pain, or calcific tendinopathy. Non-operative treatment such as physiotherapy and exercise programmes are effective and safe in many cases.</p> <p>For patients who have persistent or progressive symptoms, in spite of adequate non-operative treatment, surgery should be considered. The latest evidence for the potential benefits and risks of subacromial shoulder decompression surgery should be discussed with the patient and a shared decision reached between surgeon and patient as to whether to proceed with surgical intervention.</p>
<p><b>Rationale for recommendation</b></p> <p>Recruiting patients with pure subacromial impingement and no other associated diagnosis, a recent randomised, pragmatic, parallel group, placebo-controlled trial investigated whether subacromial decompression compared with placebo (arthroscopy only) surgery improved pain and function<sup>1</sup>. While statistically better</p>

scores were reached by patients who had both types of surgery compared to no surgery, the differences were not clinically significant, which questions the value of this type of surgery.

On the other hand, a more recent prospective randomised trial comparing the long term outcome (10 year follow up) of surgical or non-surgical treatment of sub acromial impingement showed surgery to be superior to non-surgical treatment.<sup>3</sup>

Other studies of limited quality identify certain patients with impingement syndrome that improve with surgical subacromial decompression if non-operative management fails.<sup>4,5</sup> There is also some evidence to show the benefit of surgery when used selectively and applying national clinical guidelines.<sup>6</sup>

A review of the literature identified one further systematic review that looked at the effectiveness of surgery.<sup>2</sup> The review was limited by the quality of evidence but their findings showed no difference between patients treated with surgery and those treated with non-surgical options.

Healthcare professionals treating patients with subacromial pain should be familiar with the NICE approved commissioning and treatment guidelines for the management of subacromial pain.<sup>7</sup>

Risks associated with arthroscopic sub-acromial decompression are low but include infection, frozen shoulder, ongoing pain, potential damage to blood vessels or nerves and those associated with having a general anaesthetic.

### References

1. Beard DJ, Rees JL, Cook JA, Rombach I, Cooper C, Merritt N, Shirkey BA, Donovan JL, Gwilym S, Savulescu J, Moser J, Gray A, Jepson M, Tracey I, Judge A, Wartolowska K, Carr AJ; CSAW Study Group. Arthroscopic subacromial decompression for subacromial shoulder pain (CSAW): a multicentre, pragmatic, parallel group, placebo-controlled, three-group, randomised surgical trial. *Lancet*. 2018 Jan 27;391(10118):329-338. doi: 10.1016/S0140-6736(17)32457-1. Epub 2017 Nov 20. PubMed PMID: 29169668; PubMed Central PMCID: PMC5803129.
2. Dorrestijn O, Stevens M, Winters JC, van der Meer K, Diercks RL. Conservative or surgical treatment for subacromial impingement syndrome? A systematic review. *J Shoulder Elbow Surg* 2009; 18: 652–60.
3. Farfaras S, Sernert N, Rostgard Christensen L, Hallström EK, Kartus JT. Subacromial Decompression Yields a Better Clinical Outcome Than Therapy Alone: A Prospective Randomized Study of Patients With a Minimum 10-Year Follow-up. *Am J Sports Med*. 2018 May;46(6):1397-1407
4. Holmgren T, Björnsson Hallgren H, Öberg B, Adolfsson L, Johansson K. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. *BMJ*. 2012 Feb 20;344:e787. doi: 10.1136/bmj.e787
5. Magaji SA, Singh HP, Pandey RK. Arthroscopic subacromial decompression is effective in selected patients with shoulder impingement syndrome. *J Bone Joint Surg Br*. 2012 Aug;94(8):1086-9

6. Jacobsen JR, Jensen CM, Deutch SR. Acromioplasty in patients selected for operation by national guidelines. *J Shoulder Elbow Surg.* 2017 Oct;26(10):1854-1861.
7. [https://www.boa.ac.uk/wp-content/uploads/2014/08/Subacromial-Shoulder-Commissioning-Guide\\_final.pdf](https://www.boa.ac.uk/wp-content/uploads/2014/08/Subacromial-Shoulder-Commissioning-Guide_final.pdf)

## M. Carpal tunnel syndrome release

### Updated description of the intervention

Carpal tunnel syndrome is common, and mild acute symptoms usually get better with time. Splinting at night, pain relief and corticosteroid injection should be considered. Surgery should be considered for persistent severe symptoms. Surgical treatment of carpal tunnel should only be offered under the criteria included below.

### Updated clinical criteria

Summary of intervention
Open or endoscopic surgical procedure to release median nerve from carpal tunnel.
Number of CCG interventions in 2017/18
44,497
Recommendation
<ul style="list-style-type: none"> <li>• Mild cases with intermittent symptoms causing little or no interference with sleep or activities require no treatment.</li> <li>• Cases with intermittent symptoms which interfere with activities or sleep should first be treated with:             <ol style="list-style-type: none"> <li>a. corticosteroid injection(s) (medication injected into the wrist: <i>good evidence for short (8-12 weeks) term effectiveness</i>)</li> <li style="text-align: center;"><b>or</b></li> <li>b. night splints (a support which prevents the wrist from moving during the night: <i>not as effective as steroid injections</i>)</li> </ol> </li> <li>• Surgical treatment of carpal tunnel should be considered if one of the following criteria are met:             <ol style="list-style-type: none"> <li>a. The symptoms significantly interfere with daily activities and sleep symptoms and have not settled to a manageable level with either one local corticosteroid injection and/or nocturnal splinting for a minimum of 8 weeks;</li> <li style="text-align: center;"><b>or</b></li> <li>b. There is either:                 <ol style="list-style-type: none"> <li>i. a permanent (ever-present) reduction in sensation in the median nerve distribution;</li> <li style="text-align: center;"><b>or</b></li> <li>ii. muscle wasting or weakness of thenar abduction (moving the thumb away from the hand).</li> </ol> </li> </ol> </li> </ul>

Nerve Conduction Studies if available are suggested for consideration before surgery to predict positive surgical outcome or where the diagnosis is uncertain.

### Rationale for recommendation

Carpal tunnel syndrome is very common, and mild cases may never require any treatment. Cases which interfere with activities or sleep may resolve or settle to a manageable level with non-operative treatments such as a steroid injection (good evidence of short-term benefit (8-12 weeks) but many progress to surgery within 1 year). Wrist splints worn at night (weak evidence of benefit) may also be used but are less effective than steroid injections and reported as less cost-effective than surgery.

In refractory (keeps coming back) or severe case surgery (good evidence of excellent clinical effectiveness and long term benefit) should be considered. The surgery has a high success rate (75 to 90%) in patients with intermittent symptoms who have had a good short-term benefit from a previous steroid injection. Surgery will also prevent patients with constant wooliness of their fingers from becoming worse and can restore normal sensation to patients with total loss of sensation over a period of months.

The hand is weak and sore for 3-6 weeks after carpal tunnel surgery but recovery of normal hand function is expected, significant complications are rare ( $\approx 4\%$ ) and the lifetime risk of the carpal tunnel syndrome recurring and requiring revision surgery has been estimated at between 4 and 15%.

### References

1. Atroshi I, Flondell M, Hofer M, Ranstam J. Methylprednisolone injections for the carpal tunnel syndrome: a randomized, placebo-controlled trial. *Annals of internal medicine*. 2013;159(5):309-17.
2. Chesterton LS, Blagojevic-Bucknall M, Burton C et al. The clinical and cost-effectiveness of corticosteroid injection versus night splints for carpal tunnel syndrome (instincts trial): An open-label, parallel group, randomised controlled trial. *Lancet*. 2018, 392: 1423-33.
3. Gerritsen AA, de Vet HC, Scholten RJ, Bertelsmann FW, de Krom MC, Bouter LM. Splinting vs surgery in the treatment of carpal tunnel syndrome: A randomized controlled trial. *JAMA*. 2002, 288: 1245-51.
4. Korthals-de Bos IB, Gerritsen AA, van Tulder MW et al. Surgery is more cost-effective than splinting for carpal tunnel syndrome in the Netherlands: Results of an economic evaluation alongside a randomized controlled trial. *BMC Musculoskelet Disord*. 2006, 7: 86.
5. Louie D , Earp B & Philip Blazar P Long-term outcomes of carpal tunnel release: a critical review of the literature *HAND* (2012) 7:242–246
6. Marshall S, Tardif G, Ashworth N. Local corticosteroid injection for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2007(2):CD001554.
7. Page MJ, Massy-Westropp N, O'Connor D, Pitt V. Splinting for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2012(7):CD010003.
8. Shi Q, MacDermid JC. Is surgical intervention more effective than non-surgical treatment for carpal tunnel syndrome? A systematic review. *J Orthop Surg Res*. 2011;6:17.

9. Stark H, Amirfeyz R. Cochrane corner: local corticosteroid injection for carpal tunnel syndrome. *J Hand Surg Eur Vol.* 2013;38(8):911-4.
10. Royal College of Surgeons:  
<https://publishing.rcseng.ac.uk/doi/10.1308/rcsbull.2017.28>
11. Verdugo RJ, Salinas RA, Castillo JL, Cea JG. Surgical versus non-surgical treatment for carpal tunnel syndrome. *Cochrane Database Syst Rev.* 2008(4):CD001552.

## N. Dupuytren's contracture release in adults

### Updated description of the intervention

NICE recommends no treatment is necessary for people with Dupuytren's disease who do not have contracture. Referral to hand surgery should be made for people with Dupuytren's contractures according to the criteria listed below.

### Updated clinical criteria

#### Summary of intervention

Dupuytren's contracture is caused by fibrous bands in the palm of the hand which draw the finger(s) (and sometimes the thumb) into the palm and prevent them from straightening fully. If not treated the finger(s) may bend so far into the palm that they cannot be straightened. All treatments aim to straighten the finger(s) to restore and retain hand function for the rest of the patient's life. However none cure the condition which can recur after any intervention so that further interventions are required.

Splinting and radiotherapy have not been shown be effective treatments of established Dupuytren's contractures.

Several treatments are available: collagenase injections, needle fasciotomy, fasciectomy and dermofasciectomy. None is entirely satisfactory with some having slower recovery periods, higher complication rates or higher reoperation rates (for recurrence) than others. The need for, and choice of, intervention should be made on an individual basis and should be a shared decision between the patient and a practitioner with expertise in the various treatments of Dupuytren's contractures.

No-one knows which interventions are best for restoring and maintaining hand function throughout the rest of the patient's life, and which are the cheapest and most cost-effective in the long term. Ongoing and planned National Institute for Health Research studies aim to address these questions.

#### Number of CCG interventions in 2017/18

14,376

#### Recommendation

- Treatment is not indicated in cases where there is no contracture, and in patients with a mild (less than 20°) contractures, or one which is not progressing and does not impair function.

- An intervention (collagenase injections, needle fasciotomy, fasciectomy and dermofasciectomy) should be considered for:
  - a. finger contractures causing loss of finger extension of 30° or more at the metacarpophalangeal joint or 20° at the proximal interphalangeal joint.
  - or**
  - b. severe thumb contractures which interfere with function
- NICE concluded that collagenase should only be used for:
  - a. Participants in the ongoing clinical trial (HTA-15/102/04)
  - or**
  - b. Adult patients with a palpable cord if:
    - i. there is evidence of moderate disease (functional problems and metacarpophalangeal joint contracture of 30° to 60° and proximal interphalangeal joint contracture of less than 30° or first web contracture) plus up to two affected joints;
    - and**
    - ii. needle fasciotomy is not considered appropriate, but limited fasciectomy is considered appropriate by the treating hand surgeon

#### Rationale for recommendation

Contractures left untreated usually progress and often fail to straighten fully with any treatment if allowed to progress too far. Complications causing loss, rather than improvement, in hand function occur more commonly after larger interventions, but larger interventions carry a lower risk of need for further surgery.

Common complications after collagenase injection are normally transient and include skin breaks and localised pain. Tendon injury is possible but very rare. Significant complications with lasting impact after needle fasciotomy are very unusual (about 1%) and include nerve injury. Such complications after fasciectomy are more common (about 4%) and include infection, numbness and stiffness.

#### References

1. [http://www.bssh.ac.uk/\\_userfiles/pages/files/Patients/Conditions/Elective/dupuytren's\\_disease\\_leaflet\\_2016.pdf](http://www.bssh.ac.uk/_userfiles/pages/files/Patients/Conditions/Elective/dupuytren's_disease_leaflet_2016.pdf)
2. <https://cks.nice.org.uk/dupuytren's-disease>
3. Crean SM, Gerber RA, Le Graverand MP, Boyd DM, Cappelleri JC. The efficacy and safety of fasciectomy and fasciotomy for Dupuytren's contracture in European patients: a structured review of published studies. *J Hand Surg Eur Vol.* 2011;36(5):396-407.
4. Krefter C, Marks M, Hensler S, Herren DB, Calcagni M. Complications after treating dupuytren's disease. A systematic literature review. *Hand surgery & rehabilitation.* 2017, 36: 322-9.
5. NICE 2004. Needle fasciotomy for Dupuytren's contracture. <https://www.nice.org.uk/guidance/ipg43>
6. NICE, 2017. Collagenase clostridium histolyticum for treating Dupuytren's contracture. : <https://www.nice.org.uk/guidance/ta459>,

7. Rodrigues JN, Becker GW, Ball C, Zhang W, Giele H, Hobby J, et al. Surgery for Dupuytren's contracture of the fingers. *Cochrane Database Syst Rev.* 2015(12):CD010143.
8. Scherman P, Jenmalm P, Dahlin LB. Three-year recurrence of Dupuytren's contracture after needle fasciotomy and collagenase injection: a two-centre randomized controlled trial. *J Hand Surg Eur Vol.* 2018;43(8):836-40.
9. Skov ST, Bisgaard T, Sondergaard P, Lange J. Injectable Collagenase Versus Percutaneous Needle Fasciotomy for Dupuytren Contracture in Proximal Interphalangeal Joints: A Randomized Controlled Trial. *J Hand Surg Am.* 2017;42(5):321-8 e3.
10. Stromberg J, Ibsen Sorensen A, Friden J. Percutaneous Needle Fasciotomy Versus Collagenase Treatment for Dupuytren Contracture: A Randomized Controlled Trial with a Two-Year Follow-up. *J Bone Joint Surg Am.* 2018;100(13):1079-86.
11. van Rijssen AL, Gerbrandy FS, Ter Linden H, Klip H, Werker PM. A comparison of the direct outcomes of percutaneous needle fasciotomy and limited fasciectomy for Dupuytren's disease: A 6-week follow-up study. *J Hand Surg Am.* 2006, 31: 717-25.
12. van Rijssen AL, ter Linden H, Werker PM. Five-year results of a randomized clinical trial on treatment in Dupuytren's disease: Percutaneous needle fasciotomy versus limited fasciectomy. *Plast Reconstr Surg.* 2012, 129: 469-77.

## O. Ganglion excision

### Updated description of the intervention

Most people live comfortably with ganglia and they often resolve spontaneously over time. Ganglion excision can be unnecessary, can cause complications, and recurrence is common following surgery. The complications may be similar to or worse than the original problem. Ganglion excision should only be offered under the criteria outlined below.

### Updated clinical criteria

#### Summary of intervention

Ganglia are cystic swellings containing jelly-like fluid which form around the wrists or in the hand. In most cases wrist ganglia cause only mild symptoms which do not restrict function, and many resolve without treatment within a year. Wrist ganglion rarely press on a nerve or other structure, causing pain and reduced hand function.

Ganglia in the palm of the hand (seed ganglia) can cause pain when carrying objects.

Ganglia which form just below the nail (mucous cysts) can deform the nail bed and discharge fluid, but occasionally become infected and can result in septic arthritis of the distal finger joint.

#### Number of CCG interventions in 2017/18

6,219

## Recommendation

### Wrist ganglia

- no treatment unless causing pain or tingling/numbness or concern (worried it is a cancer);
- aspiration if causing pain, tingling/numbness or concern
- surgical excision only considered if aspiration fails to resolve the pain or tingling/numbness and there is restricted hand function.

### Seed ganglia that are painful

- puncture/aspirate the ganglion using a hypodermic needle
- surgical excision only considered if ganglion persists or recurs after puncture/aspiration.

### Mucous cysts

- no surgery considered unless recurrent spontaneous discharge of fluid or significant nail deformity.

## Rationale for recommendation

Most wrist ganglia get better on their own. Surgery causes restricted wrist and hand function for 4-6 weeks, may leave an unsightly scar and be complicated by recurrent ganglion formation. Aspiration of wrist ganglia may relieve pain and restore hand function, and “cure” a minority (30%). Most ganglia reform after aspiration but they may then be painless. Aspiration also reassures the patient that the swelling is not a cancer but a benign cyst full of jelly.

Complication and recurrence are rare after aspiration and surgery for seed ganglia

## Reference

1. Head L, Gencarelli JR, Allen M, Boyd KU. Wrist ganglion treatment: Systematic review and meta-analysis. J Hand Surg Am. 2015, 40: 546-53 e8.
2. Naam NH, Carr SB, Massoud AH. Intra-neural Ganglions of the Hand and Wrist. J Hand Surg Am. 2015 Aug;40(8):1625-30. doi: 10.1016/j.jhsa.2015.05.025. PubMed PMID: 26213199.
3. [http://www.bssh.ac.uk/userfiles/pages/files/Patients/Conditions/Elective/ganglion\\_cyst\\_leaflet-2016.pdf](http://www.bssh.ac.uk/userfiles/pages/files/Patients/Conditions/Elective/ganglion_cyst_leaflet-2016.pdf)

## P. Trigger finger release in adults

### Updated description of the intervention

Trigger finger often resolves over time and is often a nuisance rather than a serious problem. If treatment is necessary steroid injection can be considered. Surgery should only be offered in specific cases according to NICE accredited guidelines by the British Society for Surgery to the Hand, where alternative measures have not been successful and persistent or recurrent triggering, or a locked finger occurs.

### Updated clinical criteria

### Summary of intervention

Trigger digit occurs when the tendons which bend the thumb/finger into the palm intermittently jam in the tight tunnel (flexor sheath) through which they run. It may occur in one or several fingers and causes the finger to “lock” in the palm of the hand. Mild triggering is a nuisance and causes infrequent locking episodes. Other cases cause pain and loss and unreliability of hand function. Mild cases require no treatment and may resolve spontaneously.

### Number of CCG interventions in 2017/18

7,789

### Recommendation

Mild cases which cause no loss of function require no treatment or avoidance of activities which precipitate triggering and may resolve spontaneously.

Cases interfering with activities or causing pain should first be treated with:

- a. one or two steroid injections which are typically successful (*strong evidence*), but the problem may recur, especially in diabetics;  
**or**
- b. splinting of the affected finger for 3-12 weeks (*weak evidence*).

Surgery should be considered if:

- a. the triggering persists or recurs after one of the above measures (particularly steroid injections);  
**or**
- b. the finger is permanently locked in the palm;  
**or**
- c. the patient has previously had 2 other trigger digits unsuccessfully treated with appropriate nonoperative methods;  
**or**
- d. diabetics.

Surgery is usually effective and requires a small skin incision in the palm, but can be done with a needle through a puncture wound (percutaneous release).

### Rationale for recommendation

Treatment with steroid injections usually resolve troublesome trigger fingers within 1 week (*strong evidence*) but sometimes the triggering keeps recurring. Surgery is normally successful (strong evidence), provides better outcomes than a single steroid injection at 1 year and usually provides a permanent cure. Recovery after surgery takes 2-4 weeks. Problems sometimes occur after surgery, but these are rare (<3%).

#### References

1. <https://www.nhs.uk/conditions/trigger-finger/treatment/>
2. Amirfeyz R, McNinch R, Watts A, Rodrigues J, Davis TRC, Glassey N, Bullock J. Evidence-based management of adult trigger digits. J Hand Surg Eur Vol. 2017 Jun;42(5):473-480. doi: 10.1177/1753193416682917. Epub 2016 Dec 21.
3. British Society for Surgery of the Hand Evidence for Surgical Treatment (BEST).  
[http://www.bssh.ac.uk/userfiles/pages/files/professionals/BEST%20Guidelines/BEST%20trigger%20finger%20PUBLISHED\(1\).pdf](http://www.bssh.ac.uk/userfiles/pages/files/professionals/BEST%20Guidelines/BEST%20trigger%20finger%20PUBLISHED(1).pdf)
4. Chang CJ, Chang SP, Kao LT, Tai TW, Jou IM. A meta-analysis of corticosteroid injection for trigger digits among patients with diabetes. Orthopedics. 2018, 41: e8-e14.
5. Everding NG, Bishop GB, Belyea CM, Soong MC. Risk factors for complications of open trigger finger release. Hand (N Y). 2015, 10: 297-300.
6. Fiorini HJ, Tamaoki MJ, Lenza M, Gomes Dos Santos JB, Faloppa F, Belloti JC. Surgery\_for\_trigger finger. Cochrane Database Syst Rev. 2018 Feb 20;2:CD009860. doi: 10.1002/14651858.CD009860.pub2. Review.
7. Hansen RL, Sondergaard M, Lange J. Open Surgery Versus Ultrasound-Guided Corticosteroid Injection for Trigger Finger: A Randomized Controlled Trial With 1-Year Follow-up. J Hand Surg Am. 2017;42(5):359-66.
8. Lunsford D, Valdes K, Hengy S. Conservative management of trigger finger: A systematic review. J Hand Ther. 2017.
9. Peters-Veluthamaningal C, Winters JC, Groenier KH, Jong BM. Corticosteroid injections effective for trigger finger in adults in general practice: a double-blinded randomised placebo controlled trial. Ann Rheum Dis. 2008 Sep;67(9):1262-6. Epub 2008 Jan 7.

## Q. Varicose vein interventions

### Updated description of the intervention

NICE has published detailed guidance on what treatment should be considered for varicose veins and when interventions for varicose veins (endothermal ablation, sclerotherapy or surgery) should be offered. Surgery is a traditional treatment that involves removal of the vein, patients can get recurrence of symptoms which may need further treatment. Treatments like endothermal ablation or ultrasound-guided foam sclerotherapy are less invasive than surgery and have replaced surgery in the management of most patients. However surgery is the most appropriate in some cases. Patients with symptomatic varicose veins should be offered treatment of their

varicose veins. Compression hosiery is not recommended if an interventional treatment is possible. <sup>14</sup>

## Updated clinical criteria

### Summary of intervention

There are various interventional procedures for treating varicose veins. These include endothermal ablation, ultrasound guided foam sclerotherapy and traditional surgery (this is a surgical procedure that involves ligation and stripping of varicose veins) all of which have been shown to be clinically and cost effective compared to no treatment or treatment with compression hosiery. Varicose veins are common and can markedly affect patients quality of life, can be associated with complications such as eczema, skin changes, thrombophlebitis, bleeding, leg ulceration, deep vein thrombosis and pulmonary embolism that can be life threatening.

### Number of CCG interventions in 2017/18

28,846

### Recommendation

1.1 Intervention in terms of, endovenous thermal (laser ablation, and radiofrequency ablation), ultrasound guided foam sclerotherapy, open surgery (ligation and stripping) are all cost effective treatments for managing symptomatic varicose veins compared to no treatment or the use of compression hosiery. For truncal ablation there is a treatment hierarchy based on the cost effectiveness and suitability, which is endothermal ablation then ultrasound guided foam, then conventional surgery.

1.2 Refer people to a vascular service if they have any of the following;-

1. Symptomatic \* primary or recurrent varicose veins.
2. Lower-limb skin changes, such as pigmentation or eczema, thought to be caused by chronic venous insufficiency.
3. Superficial vein thrombophlebitis (characterised by the appearance of hard, painful veins) and suspected venous incompetence.
4. A venous leg ulcer (a break in the skin below the knee that has not healed within 2 weeks).
5. A healed venous leg ulcer.

\*Symptomatic: "Veins found in association with troublesome lower limb symptoms (typically pain, aching, discomfort, swelling, heaviness and itching)."

For patients whose veins are purely cosmetic and are not associated with any symptoms do not refer for NHS treatment

1.3 Refer people with bleeding varicose veins to a vascular service immediately.

1.4 Do not offer compression hosiery to treat varicose veins unless interventional treatment is unsuitable.

<sup>14</sup> <https://www.nice.org.uk/guidance/qs67>

For further information, please see:

- <https://www.nice.org.uk/guidance/qs67>
- <https://www.guidelinesinpractice.co.uk/nice-referral-advice-11-varicose-veins/300594.article>
- <https://www.nice.org.uk/guidance/cg168>

### Rationale for recommendation

International guidelines, NICE guidance and NICE Quality standards provide clear evidence of the clinical and cost-effectiveness that patients with symptomatic varicose veins should be referred to a vascular service for assessment including duplex ultrasound.

Open surgery is a traditional treatment that involves surgical removal by 'stripping' out the vein or ligation (tying off the vein), this is still a valuable technique, it is still a clinically and cost-effective treatment technique for some patients but has been mainly superseded by endothermal ablation and ultrasound guided foam sclerotherapy.

Recurrence of symptoms can occur due to the development of further venous disease, that will benefit from further intervention (see above). NICE guidance states that a review of the data from the trials of interventional procedures indicates that the rate of clinical recurrence of varicose veins at 3 years after treatment is likely to be between 10–30%.

For people with confirmed varicose veins and truncal reflux NICE recommends:

- Offer endothermal ablation of the truncal vein.
- If endothermal ablation is unsuitable, offer ultrasound-guided foam sclerotherapy.
- If ultrasound-guided foam sclerotherapy is unsuitable, offer surgery.
- Consider treatment of tributaries at the same time
- Do not offer compression hosiery to treat varicose veins unless interventional treatment is unsuitable.

Complications of intervention include recurrence of varicose veins, infection, pain, bleeding, and more rarely blood clot in the leg. Complications of non-intervention include decreasing quality of life for patients, increased symptomatology, disease progression potentially to skin changes and eventual leg ulceration, deep vein thrombosis and pulmonary embolism.

### References

1. NICE Guidance: <https://www.guidelinesinpractice.co.uk/nice-referral-advice-11-varicose-veins/300594.article>
2. NICE Guidance: <https://www.nice.org.uk/guidance/cg168>
3. NICE Quality Standard: <https://www.nice.org.uk/guidance/qs67>
4. Editor's Choice - Management of Chronic Venous Disease: Clinical Practice Guidelines of the European Society for Vascular Surgery (ESVS). Wittens C, Davies AH, Bækgaard N, Broholm R, Cavezzi A, Chastanet S, de Wolf

M, Eggen C, Giannoukas A, Gohel M, Kakkos S, Lawson J, Noppeney T, Onida S, Pittaluga P, Thomis S, Toonder I, Vuylsteke M, Esvs Guidelines Committee, Kolh P, de Borst GJ, Chakfé N, Debus S, Hinchliffe R, Koncar I, Lindholt J, de Ceniga MV, Vermassen F, Verzini F, Document Reviewers, De Maeseneer MG, Blomgren L, Hartung O, Kalodiki E, Korten E, Lugli M, Naylor R, Nicolini P, Rosales A Eur J Vasc Endovasc Surg. 2015 Jun;49(6):678-737. doi: 10.1016/j.ejvs.2015.02.007. Epub 2015 Apr 25.

5. The care of patients with varicose veins and associated chronic venous diseases: clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. Gloviczki P1, Comerota AJ, Dalsing MC, Eklof BG, Gillespie DL, Gloviczki ML, Lohr JM, McLafferty RB, Meissner MH, Murad MH, Padberg FT, Pappas PJ, Passman MA, Raffetto JD, Vasquez MA, Wakefield TW; Society for Vascular Surgery; American Venous Forum. J Vasc Surg. 2011 May;53(5 Suppl):2S-48S. doi: 10.1016/j.jvs.2011.01.079..
6. A Randomized Trial of Early Endovenous Ablation in Venous Ulceration. Gohel MS1, Heatley F1, Liu X1, Bradbury A1, Bulbulia R1, Cullum N1, Epstein DM1, Nyamekye I1, Poskitt KR1, Renton S1, Warwick J1, Davies AH1; EVRA Trial Investigators. N Engl J Med. 2018 May 31;378(22):2105-2114. doi: 10.1056/NEJMoa1801214. Epub 2018 Apr 24

# Activity goals

This section sets out the 2017/18 activity baseline for STPs, CCGs and providers. The data tables for STPs and CCGs also set out the expected activity reduction in response to the national clinical criteria set out in the previous section.

We have segmented the activity and activity reduction into the following tables:

- Table 2: 2017/18 CCG activity and activity reduction for Category 1 interventions (A-D)
- Table 3: 2017/18 CCG activity and activity reduction for Category 2 interventions (E-K)
- Table 4: 2017/18 CCG activity and activity reduction for Category 2 interventions (L-Q)
- Table 5: 2017/18 STP activity and activity reduction for Category 1 interventions (A-D)
- Table 6: 2017/18 STP activity and activity reduction for Category 2 interventions (E-K)
- Table 7: 2017/18 STP activity and activity reduction for Category 2 interventions (L-Q)
- Table 8: 2017/18 provider activity baseline estimates for Category 1 interventions (A-D) and total category 1 activity
- Table 9: 2017/18 provider activity baseline estimates for Category 2 interventions (E-Q) and total Category 2 activity.

The Evidence-Based Interventions Programme and the clinical criteria for the 17 interventions apply in all care settings. However, the 2017/18 activity and activity goals set out in the data tables are necessarily based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what we are reliably able to measure nationally, outpatient activity is therefore not included. We will work with our demonstrator community to improve data for both in and outpatient settings

Note: the Category 1 activity reduction goals do not include Individual Funding Request (IFR) activity.

Note: the sum of the 2017/18 activity values in the provider table do not exactly match those in the CCG and STP tables as we have not excluded non-CCG activity from provider table.

**Table 2<sup>15</sup>: 2017/18 CCG Activity and activity reduction for Category 1 interventions (A-D) <sup>16</sup>**

CCG code	CCG name	A Surgery for snoring		B Dilatation & curettage for heavy menstrual bleeding		C Knee arthroscopy with osteoarthritis		D Injection for nonspecific low back pain without sciatica	
		No of spells		No of spells		No of spells		No of spells	
		2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
00C	NHS Darlington CCG	0	0	#	#	#	#	51	51
00D	NHS Durham Dales, Easington and Sedgefield CCG	7	7	#	#	11	11	114	114
00J	NHS North Durham CCG	7	7	0	0	15	15	38	38
00K	NHS Hartlepool and Stockton-on-Tees CCG	#	#	#	#	#	#	15	15
00L	NHS Northumberland CCG	#	#	#	#	6	6	32	32
00M	NHS South Tees CCG	#	#	0	0	7	7	35	35
00N	NHS South Tyneside CCG	5	5	0	0	11	11	74	74
00P	NHS Sunderland CCG	7	7	0	0	23	23	108	108
00Q	NHS Blackburn with Darwen CCG	8	8	#	#	18	18	24	24
00R	NHS Blackpool CCG	#	#	0	0	13	13	186	186
00T	NHS Bolton CCG	9	9	#	#	16	16	17	17
00V	NHS Bury CCG	#	#	0	0	13	13	74	74
00X	NHS Chorley and South Ribble CCG	7	7	#	#	11	11	163	163
00Y	NHS Oldham CCG	#	#	0	0	17	17	18	18
01A	NHS East Lancashire CCG	15	15	0	0	56	56	59	59
01C	NHS Eastern Cheshire CCG	#	#	0	0	#	#	20	20

<sup>15</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>16</sup> National activity figures are based on CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>01D</b>	NHS Heywood, Middleton and Rochdale CCG	#	#	#	#	14	14	63	63
<b>01E</b>	NHS Greater Preston CCG	17	17	#	#	#	#	254	254
<b>01F</b>	NHS Halton CCG	6	6	#	#	9	9	11	11
<b>01G</b>	NHS Salford CCG	#	#	0	0	22	22	55	55
<b>01H</b>	NHS Cumbria CCG	5	5	0	0	68	68	165	165
<b>01J</b>	NHS Knowsley CCG	7	7	#	#	#	#	46	46
<b>01K</b>	NHS Morecambe Bay CCG	7	7	6	6	39	39	429	429
<b>01R</b>	NHS South Cheshire CCG	#	#	0	0	15	15	50	50
<b>01T</b>	NHS South Sefton CCG	6	6	0	0	#	#	74	74
<b>01V</b>	NHS Southport and Formby CCG	#	#	0	0	#	#	24	24
<b>01W</b>	NHS Stockport CCG	#	#	0	0	9	9	15	15
<b>01X</b>	NHS St Helens CCG	#	#	0	0	#	#	25	25
<b>01Y</b>	NHS Tameside and Glossop CCG	#	#	0	0	31	31	77	77
<b>02A</b>	NHS Trafford CCG	#	#	0	0	15	15	41	41
<b>02D</b>	NHS Vale Royal CCG	0	0	0	0	10	10	17	17
<b>02E</b>	NHS Warrington CCG	#	#	5	5	16	16	10	10
<b>02F</b>	NHS West Cheshire CCG	#	#	0	0	8	8	143	143
<b>02G</b>	NHS West Lancashire CCG	#	#	#	#	#	#	19	19
<b>02H</b>	NHS Wigan Borough CCG	#	#	#	#	11	11	24	24
<b>02M</b>	NHS Fylde & Wyre CCG	6	6	0	0	15	15	225	225
<b>02N</b>	NHS Airedale, Wharfedale and Craven CCG	#	#	0	0	#	#	23	23
<b>02P</b>	NHS Barnsley CCG	5	5	#	#	53	53	163	163
<b>02Q</b>	NHS Bassetlaw CCG	5	5	0	0	44	44	53	53
<b>02R</b>	NHS Bradford Districts CCG	10	10	0	0	44	44	24	24
<b>02T</b>	NHS Calderdale CCG	5	5	#	#	20	20	54	54
<b>02W</b>	NHS Bradford City CCG	#	#	#	#	7	7	5	5
<b>02X</b>	NHS Doncaster CCG	#	#	#	#	62	62	326	326
<b>02Y</b>	NHS East Riding of Yorkshire CCG	13	13	#	#	26	26	42	42
<b>03A</b>	NHS Greater Huddersfield CCG	9	9	0	0	23	23	47	47
<b>03D</b>	NHS Hambleton, Richmondshire and Whitby CCG	#	#	0	0	#	#	8	8

OFFICIAL

<b>03E</b>	NHS Harrogate and Rural District CCG	#	#	0	0	8	8	#	#
<b>03F</b>	NHS Hull CCG	15	15	0	0	22	22	23	23
<b>03H</b>	NHS North East Lincolnshire CCG	0	0	0	0	48	48	10	10
<b>03J</b>	NHS North Kirklees CCG	#	#	#	#	10	10	9	9
<b>03K</b>	NHS North Lincolnshire CCG	#	#	0	0	26	26	28	28
<b>03L</b>	NHS Rotherham CCG	5	5	#	#	28	28	139	139
<b>03M</b>	NHS Scarborough and Ryedale CCG	#	#	0	0	9	9	#	#
<b>03N</b>	NHS Sheffield CCG	17	17	0	0	8	8	40	40
<b>03Q</b>	NHS Vale of York CCG	#	#	#	#	24	24	15	15
<b>03R</b>	NHS Wakefield CCG	#	#	#	#	70	70	33	33
<b>03T</b>	NHS Lincolnshire East CCG	#	#	#	#	39	39	174	174
<b>03V</b>	NHS Corby CCG	0	0	#	#	6	6	31	31
<b>03W</b>	NHS East Leicestershire and Rutland CCG	#	#	0	0	18	18	73	73
<b>03X</b>	NHS Erewash CCG	#	#	#	#	#	#	68	68
<b>03Y</b>	NHS Hardwick CCG	#	#	0	0	12	12	90	90
<b>04C</b>	NHS Leicester City CCG	#	#	0	0	29	29	58	58
<b>04D</b>	NHS Lincolnshire West CCG	#	#	0	0	37	37	45	45
<b>04E</b>	NHS Mansfield and Ashfield CCG	#	#	0	0	18	18	86	86
<b>04F</b>	NHS Milton Keynes CCG	#	#	#	#	19	19	48	48
<b>04G</b>	NHS Nene CCG	5	5	10	10	57	57	93	93
<b>04H</b>	NHS Newark & Sherwood CCG	#	#	#	#	6	6	101	101
<b>04J</b>	NHS North Derbyshire CCG	10	10	0	0	19	19	143	143
<b>04K</b>	NHS Nottingham City CCG	#	#	0	0	11	11	248	248
<b>04L</b>	NHS Nottingham North and East CCG	0	0	0	0	#	#	155	155
<b>04M</b>	NHS Nottingham West CCG	0	0	0	0	0	0	59	59
<b>04N</b>	NHS Rushcliffe CCG	0	0	0	0	#	#	95	95
<b>04Q</b>	NHS South West Lincolnshire CCG	#	#	#	#	14	14	102	102
<b>04R</b>	NHS Southern Derbyshire CCG	#	#	#	#	17	17	63	63
<b>04V</b>	NHS West Leicestershire CCG	#	#	0	0	18	18	91	91

OFFICIAL

<b>04Y</b>	NHS Cannock Chase CCG	0	0	#	#	11	11	12	12
<b>05A</b>	NHS Coventry and Rugby CCG	14	14	8	8	43	43	88	88
<b>05C</b>	NHS Dudley CCG	#	#	#	#	43	43	48	48
<b>05D</b>	NHS East Staffordshire CCG	0	0	0	0	16	16	5	5
<b>05F</b>	NHS Herefordshire CCG	#	#	0	0	10	10	#	#
<b>05G</b>	NHS North Staffordshire CCG	#	#	#	#	9	9	16	16
<b>05H</b>	NHS Warwickshire North CCG	#	#	0	0	37	37	9	9
<b>05J</b>	NHS Redditch and Bromsgrove CCG	0	0	#	#	9	9	27	27
<b>05L</b>	NHS Sandwell and West Birmingham CCG	#	#	#	#	44	44	71	71
<b>05N</b>	NHS Shropshire CCG	#	#	#	#	11	11	17	17
<b>05Q</b>	NHS South East Staffordshire and Seisdon Peninsula CCG	#	#	#	#	17	17	27	27
<b>05R</b>	NHS South Warwickshire CCG	10	10	#	#	12	12	24	24
<b>05T</b>	NHS South Worcestershire CCG	0	0	#	#	14	14	56	56
<b>05V</b>	NHS Stafford and Surrounds CCG	#	#	0	0	6	6	15	15
<b>05W</b>	NHS Stoke on Trent CCG	#	#	0	0	18	18	19	19
<b>05X</b>	NHS Telford and Wrekin CCG	0	0	#	#	9	9	6	6
<b>05Y</b>	NHS Walsall CCG	#	#	#	#	42	42	27	27
<b>06A</b>	NHS Wolverhampton CCG	#	#	#	#	10	10	57	57
<b>06D</b>	NHS Wyre Forest CCG	#	#	#	#	7	7	24	24
<b>06F</b>	NHS Bedfordshire CCG	#	#	#	#	28	28	67	67
<b>06H</b>	NHS Cambridgeshire and Peterborough CCG	9	9	#	#	31	31	171	171
<b>06K</b>	NHS East and North Hertfordshire CCG	7	7	#	#	18	18	59	59
<b>06L</b>	NHS Ipswich and East Suffolk CCG	#	#	#	#	14	14	211	211
<b>06M</b>	NHS Great Yarmouth and Waveney CCG	14	14	#	#	16	16	15	15
<b>06N</b>	NHS Herts Valleys CCG	19	19	#	#	34	34	80	80
<b>06P</b>	NHS Luton CCG	#	#	0	0	14	14	43	43
<b>06Q</b>	NHS Mid Essex CCG	0	0	#	#	61	61	21	21
<b>06T</b>	NHS North East Essex CCG	#	#	#	#	16	16	51	51

OFFICIAL

<b>06V</b>	NHS North Norfolk CCG	14	14	#	#	#	#	27	27
<b>06W</b>	NHS Norwich CCG	12	12	#	#	#	#	42	42
<b>06Y</b>	NHS South Norfolk CCG	9	9	#	#	8	8	75	75
<b>07G</b>	NHS Thurrock CCG	#	#	0	0	9	9	17	17
<b>07H</b>	NHS West Essex CCG	#	#	#	#	13	13	32	32
<b>07J</b>	NHS West Norfolk CCG	#	#	#	#	11	11	53	53
<b>07K</b>	NHS West Suffolk CCG	#	#	0	0	23	23	71	71
<b>07L</b>	NHS Barking and Dagenham CCG	#	#	#	#	11	11	50	50
<b>07M</b>	NHS Barnet CCG	#	#	#	#	10	10	71	71
<b>07N</b>	NHS Bexley CCG	#	#	0	0	8	8	66	66
<b>07P</b>	NHS Brent CCG	#	#	#	#	23	23	37	37
<b>07Q</b>	NHS Bromley CCG	#	#	0	0	26	26	143	143
<b>07R</b>	NHS Camden CCG	#	#	#	#	#	#	37	37
<b>07T</b>	NHS City and Hackney CCG	#	#	0	0	5	5	107	107
<b>07V</b>	NHS Croydon CCG	#	#	#	#	17	17	50	50
<b>07W</b>	NHS Ealing CCG	#	#	6	6	22	22	23	23
<b>07X</b>	NHS Enfield CCG	5	5	#	#	6	6	104	104
<b>07Y</b>	NHS Hounslow CCG	#	#	0	0	9	9	16	16
<b>08A</b>	NHS Greenwich CCG	5	5	0	0	8	8	43	43
<b>08C</b>	NHS Hammersmith and Fulham CCG	#	#	#	#	10	10	23	23
<b>08D</b>	NHS Haringey CCG	#	#	#	#	10	10	32	32
<b>08E</b>	NHS Harrow CCG	#	#	#	#	36	36	25	25
<b>08F</b>	NHS Havering CCG	6	6	#	#	18	18	78	78
<b>08G</b>	NHS Hillingdon CCG	5	5	#	#	35	35	45	45
<b>08H</b>	NHS Islington CCG	7	7	0	0	6	6	54	54
<b>08J</b>	NHS Kingston CCG	0	0	0	0	#	#	47	47
<b>08K</b>	NHS Lambeth CCG	11	11	0	0	16	16	85	85
<b>08L</b>	NHS Lewisham CCG	#	#	#	#	5	5	20	20
<b>08M</b>	NHS Newham CCG	5	5	0	0	5	5	27	27
<b>08N</b>	NHS Redbridge CCG	5	5	0	0	12	12	83	83
<b>08P</b>	NHS Richmond CCG	0	0	0	0	#	#	16	16

OFFICIAL

<b>08Q</b>	NHS Southwark CCG	6	6	0	0	9	9	83	83
<b>08R</b>	NHS Merton CCG	#	#	#	#	#	#	85	85
<b>08T</b>	NHS Sutton CCG	#	#	0	0	#	#	145	145
<b>08V</b>	NHS Tower Hamlets CCG	7	7	0	0	12	12	29	29
<b>08W</b>	NHS Waltham Forest CCG	#	#	#	#	12	12	40	40
<b>08X</b>	NHS Wandsworth CCG	#	#	0	0	#	#	105	105
<b>08Y</b>	NHS West London CCG	#	#	#	#	10	10	32	32
<b>09A</b>	NHS Central London (Westminster) CCG	#	#	#	#	#	#	28	28
<b>09C</b>	NHS Ashford CCG	#	#	0	0	#	#	81	81
<b>09D</b>	NHS Brighton and Hove CCG	6	6	#	#	6	6	118	118
<b>09E</b>	NHS Canterbury and Coastal CCG	#	#	0	0	13	13	151	151
<b>09F</b>	NHS Eastbourne, Hailsham and Seaford CCG	#	#	#	#	13	13	19	19
<b>09G</b>	NHS Coastal West Sussex CCG	12	12	#	#	31	31	205	205
<b>09H</b>	NHS Crawley CCG	0	0	0	0	11	11	159	159
<b>09J</b>	NHS Dartford, Gravesham and Swanley CCG	#	#	0	0	15	15	112	112
<b>09L</b>	NHS East Surrey CCG	0	0	0	0	12	12	141	141
<b>09N</b>	NHS Guildford and Waverley CCG	#	#	#	#	#	#	61	61
<b>09P</b>	NHS Hastings and Rother CCG	#	#	0	0	10	10	28	28
<b>09W</b>	NHS Medway CCG	5	5	#	#	37	37	38	38
<b>09X</b>	NHS Horsham and Mid Sussex CCG	0	0	0	0	7	7	115	115
<b>09Y</b>	NHS North West Surrey CCG	#	#	#	#	#	#	163	163
<b>10A</b>	NHS South Kent Coast CCG	#	#	#	#	11	11	139	139
<b>10C</b>	NHS Surrey Heath CCG	#	#	#	#	#	#	14	14
<b>10D</b>	NHS Swale CCG	#	#	#	#	13	13	34	34
<b>10E</b>	NHS Thanet CCG	#	#	#	#	13	13	174	174
<b>10J</b>	NHS North Hampshire CCG	0	0	0	0	8	8	10	10
<b>10K</b>	NHS Fareham and Gosport CCG	#	#	0	0	10	10	15	15
<b>10L</b>	NHS Isle of Wight CCG	#	#	0	0	14	14	16	16
<b>10Q</b>	NHS Oxfordshire CCG	#	#	0	0	15	15	7	7

## OFFICIAL

<b>10R</b>	NHS Portsmouth CCG	#	#	0	0	11	11	6	6
<b>10V</b>	NHS South Eastern Hampshire CCG	#	#	0	0	7	7	25	25
<b>10X</b>	NHS Southampton CCG	#	#	0	0	16	16	7	7
<b>11A</b>	NHS West Hampshire CCG	5	5	#	#	18	18	29	29
<b>11E</b>	NHS Bath and North East Somerset CCG	0	0	9	9	11	11	11	11
<b>11J</b>	NHS Dorset CCG	7	7	#	#	29	29	42	42
<b>11M</b>	NHS Gloucestershire CCG	7	7	#	#	24	24	12	12
<b>11N</b>	NHS Kernow CCG	#	#	#	#	45	45	29	29
<b>11X</b>	NHS Somerset CCG	#	#	#	#	26	26	7	7
<b>12D</b>	NHS Swindon CCG	#	#	#	#	25	25	#	#
<b>12F</b>	NHS Wirral CCG	#	#	0	0	11	11	58	58
<b>13T</b>	NHS Newcastle Gateshead CCG	10	10	#	#	40	40	161	161
<b>14L</b>	NHS Manchester CCG	#	#	#	#	24	24	95	95
<b>14Y</b>	NHS Buckinghamshire CCG	#	#	#	#	13	13	24	24
<b>15A</b>	NHS Berkshire West CCG	#	#	0	0	35	35	7	7
<b>15C</b>	NHS Bristol, North Somerset and South Gloucestershire CCG	7	7	7	7	41	41	12	12
<b>15D</b>	NHS Berkshire East CCG	#	#	0	0	18	18	37	37
<b>15E</b>	NHS Birmingham and Solihull CCG	#	#	5	5	89	89	307	307
<b>15F</b>	NHS Leeds CCG	33	33	#	#	57	57	49	49
<b>99A</b>	NHS Liverpool CCG	10	10	#	#	6	6	72	72
<b>99C</b>	NHS North Tyneside CCG	6	6	0	0	7	7	52	52
<b>99D</b>	NHS South Lincolnshire CCG	#	#	#	#	14	14	105	105
<b>99E</b>	NHS Basildon and Brentwood CCG	5	5	0	0	13	13	23	23
<b>99F</b>	NHS Castle Point and Rochford CCG	5	5	0	0	12	12	208	208
<b>99G</b>	NHS Southend CCG	#	#	0	0	11	11	193	193
<b>99H</b>	NHS Surrey Downs CCG	#	#	#	#	10	10	243	243
<b>99J</b>	NHS West Kent CCG	8	8	0	0	30	30	123	123
<b>99K</b>	NHS High Weald Lewes Havens CCG	#	#	#	#	7	7	37	37

OFFICIAL

<b>99M</b>	NHS North East Hampshire and Farnham CCG	#	#	#	#	9	9	38	38
<b>99N</b>	NHS Wiltshire CCG	8	8	8	8	67	67	28	28
<b>99P</b>	NHS Northern, Eastern and Western Devon CCG	13	13	#	#	23	23	68	68
<b>99Q</b>	NHS South Devon and Torbay CCG	6	6	0	0	13	13	178	178
<b>Total</b>		812	812	236	236	3,437	3,437	13,165	13,165

Table 3<sup>17</sup>: 2017/18 CCG Activity and activity reduction for Category 2 interventions (E-K) <sup>18</sup>

		E Breast reduction		F Removal of benign skin lesions		G Grommets		H Tonsillectomy		I Haemorrhoid surgery		J Hysterectomy for heavy bleeding		K Chalazia removal	
CCG code	CCG name	No of spells		No of spells		No of spells		No of spells		No of spells		No of spells		No of spells	
		2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
00C	NHS Darlington CCG	#	0	224	82	38	27	53	6	33	21	75	31	18	14
00D	NHS Durham Dales, Easington and Sedgefield CCG	17	8	730	343	126	99	164	42	50	19	208	88	68	59
00J	NHS North Durham CCG	7	0	792	461	96	74	136	23	42	15	169	69	18	10
00K	NHS Hartlepool and Stockton-on-Tees CCG	26	17	1141	764	56	25	131	0	33	#	175	59	40	31
00L	NHS Northumberland CCG	31	21	803	352	42	14	99	0	14	0	187	48	#	0

<sup>17</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>18</sup> National activity figures are based on CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>00M</b>	NHS South Tees CCG	17	8	966	591	30	0	134	0	59	29	161	49	25	16
<b>00N</b>	NHS South Tyneside CCG	11	6	497	291	29	14	131	65	14	0	119	57	48	43
<b>00P</b>	NHS Sunderland CCG	19	10	1149	780	82	56	179	58	33	#	203	90	87	78
<b>00Q</b>	NHS Blackburn with Darwen CCG	5	0	261	55	42	21	131	43	39	22	73	11	59	53
<b>00R</b>	NHS Blackpool CCG	11	6	787	557	29	13	131	59	10	0	77	9	9	#
<b>00T</b>	NHS Bolton CCG	7	0	414	38	51	16	212	65	30	0	135	20	14	#
<b>00V</b>	NHS Bury CCG	#	0	207	0	28	6	92	0	11	0	83	#	22	15
<b>00X</b>	NHS Chorley and South Ribble CCG	5	0	474	234	21	#	99	21	38	18	83	8	14	8
<b>00Y</b>	NHS Oldham CCG	#	0	278	0	30	0	147	21	28	#	139	47	47	39
<b>01A</b>	NHS East Lancashire CCG	9	0	689	208	84	44	284	111	77	38	191	45	92	80
<b>01C</b>	NHS Eastern Cheshire CCG	7	0	223	0	14	0	131	49	26	#	85	0	43	36
<b>01D</b>	NHS Heywood, Middleton and Rochdale CCG	10	#	339	57	14	0	168	57	27	#	109	23	25	17
<b>01E</b>	NHS Greater Preston CCG	7	#	531	276	17	0	131	32	33	12	75	0	13	6
<b>01F</b>	NHS Halton CCG	16	12	344	180	23	9	86	27	21	7	77	25	25	21
<b>01G</b>	NHS Salford CCG	9	#	376	58	63	33	201	67	34	7	117	26	38	29

OFFICIAL

<b>01H</b>	NHS Cumbria CCG	32	22	245	0	32	#	234	105	23	0	295	160	#	0
<b>01J</b>	NHS Knowsley CCG	14	9	488	284	25	8	126	51	37	20	119	55	30	25
<b>01K</b>	NHS Morecambe Bay CCG	19	8	608	135	58	29	215	71	57	20	197	57	12	#
<b>01R</b>	NHS South Cheshire CCG	#	0	413	170	15	0	108	30	18	0	88	13	42	36
<b>01T</b>	NHS South Sefton CCG	9	#	391	189	23	8	112	46	15	0	95	33	39	34
<b>01V</b>	NHS Southport and Formby CCG	9	5	311	133	15	5	50	#	8	0	73	20	10	6
<b>01W</b>	NHS Stockport CCG	9	0	572	169	58	27	224	90	51	18	143	18	19	9
<b>01X</b>	NHS St Helens CCG	8	#	531	271	40	21	134	50	37	16	154	73	49	43
<b>01Y</b>	NHS Tameside and Glossop CCG	7	0	292	0	56	30	222	110	44	18	157	60	33	25
<b>02A</b>	NHS Trafford CCG	9	#	506	202	43	17	126	17	55	30	97	#	74	66
<b>02D</b>	NHS Vale Royal CCG	#	0	165	27	12	#	61	15	10	0	44	0	18	14
<b>02E</b>	NHS Warrington CCG	10	#	426	148	21	0	106	11	43	20	87	0	28	21
<b>02F</b>	NHS West Cheshire CCG	13	5	761	411	46	22	142	32	15	0	129	22	14	6
<b>02G</b>	NHS West Lancashire CCG	#	0	383	232	11	#	60	11	12	0	55	9	31	27
<b>02H</b>	NHS Wigan Borough CCG	17	7	1012	592	44	12	190	48	44	9	237	106	61	51

OFFICIAL

<b>02M</b>	NHS Fylde & Wyre CCG	11	5	1071	825	27	13	128	65	12	0	93	20	16	11
<b>02N</b>	NHS Airedale, Wharfedale and Craven CCG	5	0	282	69	51	35	79	12	24	7	93	28	24	19
<b>02P</b>	NHS Barnsley CCG	5	0	1183	850	45	19	159	46	37	10	165	61	46	38
<b>02Q</b>	NHS Bassetlaw CCG	5	#	123	0	24	13	65	16	17	#	56	8	7	#
<b>02R</b>	NHS Bradford Districts CCG	15	6	401	0	79	40	190	27	46	13	172	49	20	9
<b>02T</b>	NHS Calderdale CCG	9	#	288	10	65	42	152	53	23	0	97	9	#	0
<b>02W</b>	NHS Bradford City CCG	#	0	129	0	27	6	65	0	14	#	34	0	20	15
<b>02X</b>	NHS Doncaster CCG	15	5	382	0	71	39	117	0	29	0	146	22	6	0
<b>02Y</b>	NHS East Riding of Yorkshire CCG	32	22	854	425	77	51	229	113	46	13	196	66	14	5
<b>03A</b>	NHS Greater Huddersfield CCG	10	#	361	51	73	48	180	65	30	5	87	0	10	#
<b>03D</b>	NHS Hambleton, Richmondshire and Whitby CCG	10	5	578	373	37	25	40	0	14	0	73	10	11	6
<b>03E</b>	NHS Harrogate and Rural District CCG	11	6	165	0	34	20	67	#	37	19	61	0	12	7
<b>03F</b>	NHS Hull CCG	18	9	688	327	110	78	325	184	59	29	120	13	12	#

OFFICIAL

<b>03H</b>	NHS North East Lincolnshire CCG	9	#	155	0	8	0	106	32	23	5	118	52	#	0
<b>03J</b>	NHS North Kirklees CCG	9	#	411	178	42	20	153	58	29	10	113	42	60	54
<b>03K</b>	NHS North Lincolnshire CCG	15	9	187	0	11	0	81	7	20	#	119	49	5	0
<b>03L</b>	NHS Rotherham CCG	15	7	500	163	40	13	140	23	63	36	178	75	119	111
<b>03M</b>	NHS Scarborough and Ryedale CCG	5	#	128	0	20	9	72	25	7	0	59	9	6	#
<b>03N</b>	NHS Sheffield CCG	33	16	2174	1,447	72	15	156	0	131	73	331	124	229	210
<b>03Q</b>	NHS Vale of York CCG	26	15	327	0	44	14	189	34	44	7	128	0	21	10
<b>03R</b>	NHS Wakefield CCG	26	15	763	288	108	71	355	192	72	33	217	69	75	63
<b>03T</b>	NHS Lincolnshire East CCG	19	11	452	98	24	#	163	67	15	0	131	27	7	0
<b>03V</b>	NHS Corby CCG	0	0	94	#	22	13	38	0	21	13	53	24	6	#
<b>03W</b>	NHS East Leicestershire and Rutland CCG	17	7	553	108	50	19	183	47	49	14	213	74	43	33
<b>03X</b>	NHS Erewash CCG	#	0	235	106	9	0	58	17	14	#	55	15	19	16
<b>03Y</b>	NHS Hardwick CCG	5	#	226	87	18	8	57	14	14	#	81	38	6	#
<b>04C</b>	NHS Leicester City CCG	12	#	521	73	59	14	269	54	54	17	154	26	103	90

OFFICIAL

<b>04D</b>	NHS Lincolnshire West CCG	15	8	395	85	57	34	169	60	21	0	114	21	#	0
<b>04E</b>	NHS Mansfield and Ashfield CCG	12	6	598	350	49	29	143	58	21	0	129	53	#	0
<b>04F</b>	NHS Milton Keynes CCG	17	9	495	152	76	41	200	57	36	6	113	#	32	22
<b>04G</b>	NHS Nene CCG	21	#	975	123	82	11	203	0	129	59	425	160	72	51
<b>04H</b>	NHS Newark & Sherwood CCG	6	#	426	246	35	22	71	15	12	0	59	#	7	#
<b>04J</b>	NHS North Derbyshire CCG	12	#	653	252	31	6	151	36	42	10	231	107	42	33
<b>04K</b>	NHS Nottingham City CCG	16	6	742	329	49	12	161	0	51	18	129	18	42	29
<b>04L</b>	NHS Nottingham North and East CCG	5	0	400	199	19	#	72	8	27	11	76	13	18	13
<b>04M</b>	NHS Nottingham West CCG	#	0	252	127	11	#	33	0	14	#	32	0	13	10
<b>04N</b>	NHS Rushcliffe CCG	8	#	367	198	21	9	45	0	11	0	46	0	10	6
<b>04Q</b>	NHS South West Lincolnshire CCG	#	0	198	16	22	9	84	29	36	21	82	25	#	0
<b>04R</b>	NHS Southern Derbyshire CCG	16	0	930	227	75	20	331	84	56	0	385	170	53	36
<b>04V</b>	NHS West Leicestershire CCG	18	6	720	218	76	40	249	79	41	0	227	74	56	44

OFFICIAL

<b>04Y</b>	NHS Cannock Chase CCG	7	#	243	69	19	6	75	19	17	#	103	48	#	0
<b>05A</b>	NHS Coventry and Rugby CCG	25	11	1483	878	45	0	426	171	43	0	209	37	102	86
<b>05C</b>	NHS Dudley CCG	13	#	1247	828	73	41	189	51	21	0	250	124	25	15
<b>05D</b>	NHS East Staffordshire CCG	#	0	155	0	21	7	96	34	13	0	79	23	#	0
<b>05F</b>	NHS Herefordshire CCG	12	6	821	559	93	76	78	5	29	8	69	0	#	0
<b>05G</b>	NHS North Staffordshire CCG	8	#	477	184	34	15	110	19	31	8	145	57	8	#
<b>05H</b>	NHS Warwickshire North CCG	8	#	721	471	25	6	144	62	14	0	106	29	18	12
<b>05J</b>	NHS Redditch and Bromsgrove CCG	7	#	185	0	35	17	61	0	15	0	126	55	5	0
<b>05L</b>	NHS Sandwell and West Birmingham CCG	15	0	870	220	52	0	316	25	79	25	241	54	89	71
<b>05N</b>	NHS Shropshire CCG	7	0	447	15	33	7	101	0	28	0	152	21	8	0
<b>05Q</b>	NHS South East Staffordshire and Seisdon Peninsula CCG	7	0	522	231	37	17	139	49	21	0	153	63	6	0
<b>05R</b>	NHS South Warwickshire CCG	17	8	439	55	49	24	173	59	19	0	155	38	13	#

OFFICIAL

<b>05T</b>	NHS South Worcestershire CCG	5	0	349	0	56	28	123	0	27	0	203	76	7	0
<b>05V</b>	NHS Stafford and Surrounds CCG	10	5	320	116	17	#	72	13	27	10	96	34	#	0
<b>05W</b>	NHS Stoke on Trent CCG	8	0	462	102	47	15	154	19	39	10	206	100	11	#
<b>05X</b>	NHS Telford and Wrekin CCG	9	#	197	0	26	6	60	0	18	0	98	26	5	0
<b>05Y</b>	NHS Walsall CCG	17	9	532	181	27	0	203	66	56	28	162	58	8	0
<b>06A</b>	NHS Wolverhampton CCG	14	6	263	0	34	#	139	7	35	7	166	64	#	0
<b>06D</b>	NHS Wyre Forest CCG	#	0	152	0	24	13	57	11	11	0	82	33	#	0
<b>06F</b>	NHS Bedfordshire CCG	11	0	1179	578	83	33	204	0	64	14	214	25	28	13
<b>06H</b>	NHS Cambridgeshire and Peterborough CCG	18	0	1276	94	155	60	557	116	72	0	389	34	85	55
<b>06K</b>	NHS East and North Hertfordshire CCG	9	0	709	0	137	76	394	124	146	85	240	6	125	106
<b>06L</b>	NHS Ipswich and East Suffolk CCG	18	6	465	0	48	10	219	50	47	#	202	39	58	46
<b>06M</b>	NHS Great Yarmouth and Waveney CCG	8	#	826	494	67	46	149	53	37	11	115	19	25	18
<b>06N</b>	NHS Herts Valleys CCG	27	9	1043	242	115	45	478	184	89	22	212	0	87	67
<b>06P</b>	NHS Luton CCG	#	0	202	0	29	0	187	64	35	13	59	0	16	8

OFFICIAL

<b>06Q</b>	NHS Mid Essex CCG	5	0	616	105	108	70	234	66	66	25	212	52	20	8
<b>06T</b>	NHS North East Essex CCG	16	6	722	256	65	32	341	191	57	21	152	14	11	0
<b>06V</b>	NHS North Norfolk CCG	#	0	633	373	20	6	71	10	20	#	93	18	17	12
<b>06W</b>	NHS Norwich CCG	#	0	690	399	35	13	161	50	26	#	99	16	12	#
<b>06Y</b>	NHS South Norfolk CCG	7	0	736	424	54	33	126	34	31	7	135	41	#	0
<b>07G</b>	NHS Thurrock CCG	#	0	176	0	35	13	144	56	44	26	88	22	40	34
<b>07H</b>	NHS West Essex CCG	10	#	312	0	51	18	262	123	86	53	125	0	21	11
<b>07J</b>	NHS West Norfolk CCG	#	0	635	387	27	12	123	55	27	8	98	25	40	34
<b>07K</b>	NHS West Suffolk CCG	9	#	981	640	49	26	198	97	31	#	130	26	#	0
<b>07L</b>	NHS Barking and Dagenham CCG	#	0	627	388	16	0	197	70	40	19	73	0	80	73
<b>07M</b>	NHS Barnet CCG	7	0	864	363	28	0	163	0	52	10	71	0	49	36
<b>07N</b>	NHS Bexley CCG	8	#	401	102	33	6	172	59	56	31	132	37	16	8
<b>07P</b>	NHS Brent CCG	15	5	477	48	19	0	106	0	60	22	91	0	65	52
<b>07Q</b>	NHS Bromley CCG	13	#	1315	872	63	26	235	80	43	6	180	38	16	5
<b>07R</b>	NHS Camden CCG	10	#	832	522	8	0	95	0	45	18	46	0	43	33
<b>07T</b>	NHS City and Hackney CCG	#	0	356	16	28	0	100	0	62	31	42	0	97	86
<b>07V</b>	NHS Croydon CCG	16	5	433	0	23	0	156	0	59	17	189	37	18	5
<b>07W</b>	NHS Ealing CCG	22	11	526	28	22	0	136	0	86	41	103	0	31	17

OFFICIAL

<b>07X</b>	NHS Enfield CCG	10	0	600	206	24	0	202	32	25	0	84	0	50	39
<b>07Y</b>	NHS Hounslow CCG	16	8	225	0	22	0	158	7	43	11	64	0	34	24
<b>08A</b>	NHS Greenwich CCG	#	0	345	14	19	0	193	42	51	22	132	26	49	39
<b>08C</b>	NHS Hammersmith and Fulham CCG	12	6	209	0	30	11	110	8	35	12	46	0	41	33
<b>08D</b>	NHS Haringey CCG	12	#	683	322	14	0	152	0	51	18	67	0	56	45
<b>08E</b>	NHS Harrow CCG	13	6	319	0	21	0	92	0	32	5	66	0	34	25
<b>08F</b>	NHS Havering CCG	10	#	1036	684	37	8	208	82	60	32	137	29	76	67
<b>08G</b>	NHS Hillingdon CCG	9	#	568	202	43	8	131	0	86	55	141	31	25	15
<b>08H</b>	NHS Islington CCG	6	0	857	575	18	0	111	0	63	37	36	0	81	72
<b>08J</b>	NHS Kingston CCG	11	5	207	0	31	9	99	0	46	24	71	0	#	0
<b>08K</b>	NHS Lambeth CCG	20	9	574	120	16	0	139	0	89	47	119	0	42	28
<b>08L</b>	NHS Lewisham CCG	17	8	935	566	14	0	168	5	60	27	130	10	38	27
<b>08M</b>	NHS Newham CCG	7	0	301	0	37	0	244	31	87	49	94	0	89	76
<b>08N</b>	NHS Redbridge CCG	6	0	567	207	36	0	215	54	46	15	94	0	121	111
<b>08P</b>	NHS Richmond CCG	19	13	216	0	19	0	106	7	36	12	42	0	11	#

OFFICIAL

<b>08Q</b>	NHS Southwark CCG	13	#	684	322	12	0	141	0	89	56	85	0	38	27
<b>08R</b>	NHS Merton CCG	12	6	196	0	13	0	115	8	45	21	70	0	14	6
<b>08T</b>	NHS Sutton CCG	16	10	152	0	20	0	118	27	32	12	125	48	6	0
<b>08V</b>	NHS Tower Hamlets CCG	#	0	305	0	31	0	140	0	86	56	48	0	96	85
<b>08W</b>	NHS Waltham Forest CCG	8	0	326	0	26	0	209	54	19	0	73	0	73	63
<b>08X</b>	NHS Wandsworth CCG	17	6	361	0	21	0	147	0	83	42	78	0	31	17
<b>08Y</b>	NHS West London CCG	9	#	299	0	15	0	88	0	54	26	53	0	27	19
<b>09A</b>	NHS Central London (Westminster) CCG	10	#	255	0	6	0	64	0	25	#	33	0	30	22
<b>09C</b>	NHS Ashford CCG	#	0	313	144	25	11	131	70	31	17	62	9	8	#
<b>09D</b>	NHS Brighton and Hove CCG	13	#	816	438	25	0	184	38	72	39	90	0	8	0
<b>09E</b>	NHS Canterbury and Coastal CCG	#	0	621	332	35	17	142	39	46	24	118	34	8	#
<b>09F</b>	NHS Eastbourne, Hailsham and Seaford CCG	7	#	566	289	17	0	141	64	29	8	140	60	25	19
<b>09G</b>	NHS Coastal West Sussex CCG	10	0	1717	993	114	70	350	150	52	0	366	152	#	0
<b>09H</b>	NHS Crawley CCG	6	#	202	43	17	#	100	36	26	12	76	27	#	0
<b>09J</b>	NHS Dartford, Gravesham	12	#	421	88	35	5	180	55	104	77	125	22	17	8

OFFICIAL

	and Swanley CCG														
<b>09L</b>	NHS East Surrey CCG	11	5	248	16	27	7	132	50	31	12	82	9	0	0
<b>09N</b>	NHS Guildford and Waverley CCG	6	0	451	162	18	0	97	0	39	16	126	38	6	0
<b>09P</b>	NHS Hastings and Rother CCG	9	#	183	0	18	#	137	63	33	13	122	43	11	5
<b>09W</b>	NHS Medway CCG	16	7	577	210	37	#	227	86	46	16	260	145	15	5
<b>09X</b>	NHS Horsham and Mid Sussex CCG	10	#	508	192	39	15	139	36	51	25	117	17	0	0
<b>09Y</b>	NHS North West Surrey CCG	24	13	517	46	42	#	229	64	76	37	146	0	46	34
<b>10A</b>	NHS South Kent Coast CCG	#	0	717	437	34	15	202	117	64	42	120	36	22	16
<b>10C</b>	NHS Surrey Heath CCG	#	0	347	222	23	13	38	0	14	#	54	14	17	14
<b>10D</b>	NHS Swale CCG	6	#	189	47	17	#	115	63	16	#	117	73	#	0
<b>10E</b>	NHS Thanet CCG	#	0	405	210	25	10	100	38	37	22	125	66	5	0
<b>10J</b>	NHS North Hampshire CCG	5	0	351	63	33	10	119	19	10	0	94	#	22	15
<b>10K</b>	NHS Fareham and Gosport CCG	7	0	207	0	25	5	69	0	21	0	144	58	#	0
<b>10L</b>	NHS Isle of Wight CCG	#	0	136	0	8	0	65	11	35	19	71	9	6	#
<b>10Q</b>	NHS Oxfordshire CCG	45	24	1910	990	86	17	299	0	40	0	164	0	11	0

OFFICIAL

<b>10R</b>	NHS Portsmouth CCG	7	0	238	0	28	5	99	0	14	0	129	50	#	0
<b>10V</b>	NHS South Eastern Hampshire CCG	10	#	251	0	19	0	74	0	21	0	151	61	6	0
<b>10X</b>	NHS Southampton CCG	5	0	344	14	21	0	158	14	12	0	121	30	5	0
<b>11A</b>	NHS West Hampshire CCG	13	0	978	212	66	15	242	11	57	0	255	22	15	0
<b>11E</b>	NHS Bath and North East Somerset CCG	12	6	268	7	11	0	50	0	22	#	73	0	#	0
<b>11J</b>	NHS Dorset CCG	31	7	3295	2,185	48	0	255	0	101	16	496	174	133	109
<b>11M</b>	NHS Gloucestershire CCG	30	11	531	0	65	#	275	0	52	0	242	0	24	#
<b>11N</b>	NHS Kernow CCG	31	13	1312	518	101	51	239	9	78	17	234	0	13	0
<b>11X</b>	NHS Somerset CCG	10	0	1170	382	94	43	234	#	63	#	301	66	13	0
<b>12D</b>	NHS Swindon CCG	11	#	243	0	38	11	127	16	43	19	103	12	6	0
<b>12F</b>	NHS Wirral CCG	35	25	1561	1,116	82	50	198	56	23	0	123	0	9	0
<b>13T</b>	NHS Newcastle Gateshead CCG	32	17	1010	374	107	60	208	0	74	23	313	133	21	#
<b>14L</b>	NHS Manchester CCG	15	0	735	45	79	9	369	27	95	36	187	0	90	69
<b>14Y</b>	NHS Buckinghamshire CCG	17	#	523	0	113	55	146	0	60	#	185	0	15	0

OFFICIAL

<b>15A</b>	NHS Berkshire West CCG	21	6	486	0	41	0	195	0	36	0	151	0	13	0
<b>15C</b>	NHS Bristol, North Somerset and South Gloucestershire CCG	28	0	2591	1,354	105	7	217	0	130	30	325	0	70	39
<b>15D</b>	NHS Berkshire East CCG	11	0	789	245	53	#	175	0	61	15	193	23	14	0
<b>15E</b>	NHS Birmingham and Solihull CCG	57	23	1858	310	198	48	660	5	112	0	616	162	138	98
<b>15F</b>	NHS Leeds CCG	54	30	1559	514	151	63	695	265	259	174	309	7	60	32
<b>99A</b>	NHS Liverpool CCG	27	12	889	255	55	6	337	84	91	39	215	30	73	56
<b>99C</b>	NHS North Tyneside CCG	22	15	563	275	42	21	76	0	23	0	156	65	8	#
<b>99D</b>	NHS South Lincolnshire CCG	13	8	438	210	24	8	88	20	19	#	99	29	#	0
<b>99E</b>	NHS Basildon and Brentwood CCG	#	0	326	0	41	12	254	128	40	11	123	14	49	40
<b>99F</b>	NHS Castle Point and Rochford CCG	6	0	350	96	27	11	165	90	35	15	92	15	23	17
<b>99G</b>	NHS Southend CCG	#	0	368	125	25	6	154	71	20	0	109	35	32	26
<b>99H</b>	NHS Surrey Downs CCG	17	8	891	487	29	0	114	0	46	13	119	0	5	0
<b>99J</b>	NHS West Kent CCG	18	#	423	0	44	0	310	93	57	6	241	40	6	0

OFFICIAL

<b>99K</b>	NHS High Weald Lewes Havens CCG	7	#	535	301	17	#	110	41	16	0	73	0	5	0
<b>99M</b>	NHS North East Hampshire and Farnham CCG	8	#	910	622	42	18	117	15	24	0	118	26	66	59
<b>99N</b>	NHS Wiltshire CCG	11	0	632	0	50	#	202	0	67	15	249	41	15	0
<b>99P</b>	NHS Northern, Eastern and Western Devon CCG	32	#	2498	1,247	232	150	419	28	79	0	550	178	28	0
<b>99Q</b>	NHS South Devon and Torbay CCG	5	0	440	24	67	42	128	16	42	10	209	84	0	0
<b>Total</b>		2,388	829	116,255	45,589	8,669	3,259	32,238	7,454	8,474	2,801	27,660	6,536	6,026	4,326

Table 4<sup>19</sup>: 2017/18 CCG Activity and activity reduction for Category 2 interventions (L-Q) <sup>20</sup>

		L Shoulder decompression		M Carpal tunnel syndrome release		N Dupuytren's contracture release		O Ganglion excision		P Trigger finger release		Q Varicose vein surgery	
CCG code	CCG name	No of spells		No of spells		No of spells		No of spells		No of spells		No of spells	
		2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
00C	NHS Darlington CCG	13	0	100	37	43	21	23	15	17	6	57	11
00D	NHS Durham Dales, Easington and Sedgefield CCG	46	#	182	7	67	#	34	13	31	0	184	57
00J	NHS North Durham CCG	27	0	162	17	51	0	39	21	31	5	210	103
00K	NHS Hartlepool and Stockton-on-Tees CCG	12	0	127	0	58	0	51	30	22	0	260	137
00L	NHS Northumberland CCG	59	10	320	111	132	54	46	23	72	34	171	22
00M	NHS South Tees CCG	28	0	356	193	104	47	38	18	49	20	188	67
00N	NHS South Tyneside CCG	29	7	91	0	47	15	20	9	32	15	190	122

<sup>19</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>20</sup> National activity figures are based on CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>00P</b>	NHS Sunderland CCG	35	0	259	96	82	24	60	40	50	21	433	312
<b>00Q</b>	NHS Blackburn with Darwen CCG	29	7	134	49	28	0	22	10	27	12	132	66
<b>00R</b>	NHS Blackpool CCG	24	0	77	0	60	23	16	#	23	5	97	22
<b>00T</b>	NHS Bolton CCG	75	35	270	111	45	0	24	#	62	34	75	0
<b>00V</b>	NHS Bury CCG	47	19	140	29	38	0	26	12	34	14	145	61
<b>00X</b>	NHS Chorley and South Ribble CCG	83	57	118	11	83	44	47	34	14	0	75	0
<b>00Y</b>	NHS Oldham CCG	73	41	46	0	55	12	27	10	36	13	200	102
<b>01A</b>	NHS East Lancashire CCG	98	47	386	178	111	37	68	42	63	26	395	239
<b>01C</b>	NHS Eastern Cheshire CCG	62	31	122	0	100	53	41	26	32	9	155	61
<b>01D</b>	NHS Heywood, Middleton and Rochdale CCG	72	42	203	84	50	9	29	13	31	10	223	131
<b>01E</b>	NHS Greater Preston CCG	76	49	153	45	83	46	43	29	31	12	110	28
<b>01F</b>	NHS Halton CCG	18	0	115	43	32	6	11	#	31	18	117	62
<b>01G</b>	NHS Salford CCG	41	9	320	193	85	43	64	46	71	49	64	0
<b>01H</b>	NHS Cumbria CCG	72	24	329	127	138	63	35	12	53	17	213	68
<b>01J</b>	NHS Knowsley CCG	45	22	129	40	31	#	12	0	32	16	93	26
<b>01K</b>	NHS Morecambe Bay CCG	59	9	412	199	165	87	47	22	67	29	313	160
<b>01R</b>	NHS South Cheshire CCG	111	85	247	139	77	38	17	#	42	23	73	0
<b>01T</b>	NHS South Sefton CCG	73	51	94	#	46	14	7	0	7	0	60	0
<b>01V</b>	NHS Southport and Formby CCG	33	14	108	25	43	13	9	0	13	0	97	39
<b>01W</b>	NHS Stockport CCG	88	45	356	178	127	66	44	23	44	13	103	0
<b>01X</b>	NHS St Helens CCG	29	#	217	101	69	27	17	#	37	16	150	65
<b>01Y</b>	NHS Tameside and Glossop CCG	122	88	198	63	67	20	25	8	24	0	83	0

OFFICIAL

<b>02A</b>	NHS Trafford CCG	47	14	190	58	85	41	39	23	32	9	103	#
<b>02D</b>	NHS Vale Royal CCG	55	40	118	56	40	18	6	0	21	10	49	#
<b>02E</b>	NHS Warrington CCG	40	9	145	22	56	14	10	0	22	0	124	32
<b>02F</b>	NHS West Cheshire CCG	48	11	186	29	81	25	22	#	37	9	126	12
<b>02G</b>	NHS West Lancashire CCG	31	15	93	25	46	21	8	0	18	6	95	46
<b>02H</b>	NHS Wigan Borough CCG	135	89	441	256	99	33	59	36	75	42	274	135
<b>02M</b>	NHS Fylde & Wyre CCG	23	0	109	0	74	31	20	8	20	0	187	107
<b>02N</b>	NHS Airedale, Wharfedale and Craven CCG	52	29	190	94	91	56	34	23	33	16	136	67
<b>02P</b>	NHS Barnsley CCG	110	74	204	57	72	20	53	35	46	20	123	14
<b>02Q</b>	NHS Bassetlaw CCG	83	66	69	0	32	6	20	11	33	20	15	0
<b>02R</b>	NHS Bradford Districts CCG	97	55	354	184	115	58	45	23	39	9	280	151
<b>02T</b>	NHS Calderdale CCG	52	21	247	125	69	26	26	11	27	5	125	33
<b>02W</b>	NHS Bradford City CCG	15	#	90	46	#	0	8	0	6	0	70	30
<b>02X</b>	NHS Doncaster CCG	106	62	226	46	99	36	75	53	40	8	53	0
<b>02Y</b>	NHS East Riding of Yorkshire CCG	141	95	391	192	141	66	35	13	93	57	205	65
<b>03A</b>	NHS Greater Huddersfield CCG	94	61	148	15	38	0	15	0	13	0	179	79
<b>03D</b>	NHS Hambleton, Richmondshire and Whitby CCG	15	0	184	87	57	20	10	0	20	#	114	46
<b>03E</b>	NHS Harrogate and Rural District CCG	53	29	141	38	63	26	11	0	25	7	125	52
<b>03F</b>	NHS Hull CCG	117	79	304	153	90	38	23	#	80	53	191	75
<b>03H</b>	NHS North East Lincolnshire CCG	70	46	167	69	57	22	6	0	11	0	61	0

OFFICIAL

<b>03J</b>	NHS North Kirklees CCG	53	29	111	13	53	20	48	35	29	12	162	87
<b>03K</b>	NHS North Lincolnshire CCG	59	34	84	0	50	12	13	#	23	#	60	0
<b>03L</b>	NHS Rotherham CCG	56	20	261	113	79	27	37	19	41	15	58	0
<b>03M</b>	NHS Scarborough and Ryedale CCG	55	37	99	22	39	10	10	#	13	0	73	18
<b>03N</b>	NHS Sheffield CCG	101	28	598	299	148	47	127	87	132	80	131	0
<b>03Q</b>	NHS Vale of York CCG	143	95	294	91	123	52	21	0	38	#	315	166
<b>03R</b>	NHS Wakefield CCG	120	69	272	64	181	108	109	83	69	32	337	181
<b>03T</b>	NHS Lincolnshire East CCG	133	96	18	0	64	#	12	0	#	0	98	0
<b>03V</b>	NHS Corby CCG	14	#	16	0	27	14	8	#	9	#	22	0
<b>03W</b>	NHS East Leicestershire and Rutland CCG	88	40	213	11	61	0	36	13	34	0	70	0
<b>03X</b>	NHS Erewash CCG	29	15	128	70	47	26	8	#	17	6	41	0
<b>03Y</b>	NHS Hardwick CCG	54	39	87	24	31	8	23	15	28	16	30	0
<b>04C</b>	NHS Leicester City CCG	36	0	231	59	27	0	16	0	42	12	69	0
<b>04D</b>	NHS Lincolnshire West CCG	64	32	25	0	62	14	10	0	5	0	110	11
<b>04E</b>	NHS Mansfield and Ashfield CCG	73	46	124	15	56	17	11	0	38	18	48	0
<b>04F</b>	NHS Milton Keynes CCG	118	81	247	106	51	6	24	#	35	10	72	0
<b>04G</b>	NHS Nene CCG	199	108	480	114	165	39	85	39	93	28	165	0
<b>04H</b>	NHS Newark & Sherwood CCG	72	52	89	8	44	14	16	6	18	#	35	0
<b>04J</b>	NHS North Derbyshire CCG	149	105	348	164	131	63	65	44	71	38	111	0
<b>04K</b>	NHS Nottingham City CCG	119	80	318	165	73	25	40	15	67	41	43	0
<b>04L</b>	NHS Nottingham North and East CCG	44	22	152	62	57	25	12	#	24	8	17	0

OFFICIAL

<b>04M</b>	NHS Nottingham West CCG	30	16	86	30	33	13	11	#	13	#	20	0
<b>04N</b>	NHS Rushcliffe CCG	44	26	131	55	49	22	16	7	21	7	18	0
<b>04Q</b>	NHS South West Lincolnshire CCG	53	33	36	0	36	5	#	0	9	0	49	0
<b>04R</b>	NHS Southern Derbyshire CCG	98	24	636	331	148	42	54	17	95	41	251	23
<b>04V</b>	NHS West Leicestershire CCG	51	0	308	89	90	11	20	0	39	0	64	0
<b>04Y</b>	NHS Cannock Chase CCG	44	25	127	49	54	26	10	0	16	#	135	77
<b>05A</b>	NHS Coventry and Rugby CCG	120	60	495	251	70	0	44	10	47	5	211	22
<b>05C</b>	NHS Dudley CCG	65	21	378	192	72	6	27	5	40	7	313	178
<b>05D</b>	NHS East Staffordshire CCG	7	0	85	#	36	7	#	0	13	0	68	8
<b>05F</b>	NHS Herefordshire CCG	22	0	303	182	62	17	25	11	32	10	100	15
<b>05G</b>	NHS North Staffordshire CCG	77	46	147	15	83	35	16	0	36	12	134	39
<b>05H</b>	NHS Warwickshire North CCG	46	19	235	124	41	#	10	0	32	12	78	0
<b>05J</b>	NHS Redditch and Bromsgrove CCG	70	45	255	153	46	9	23	10	20	#	140	64
<b>05L</b>	NHS Sandwell and West Birmingham CCG	56	0	136	0	56	0	18	0	16	0	351	146
<b>05N</b>	NHS Shropshire CCG	163	117	422	222	119	45	62	40	64	28	189	48
<b>05Q</b>	NHS South East Staffordshire and Seisdon Peninsula CCG	26	0	168	37	59	12	11	0	25	#	167	71
<b>05R</b>	NHS South Warwickshire CCG	70	29	324	152	98	37	36	16	56	26	102	0
<b>05T</b>	NHS South Worcestershire CCG	32	0	434	245	90	22	68	46	41	7	166	30
<b>05V</b>	NHS Stafford and Surrounds CCG	54	32	80	0	68	34	11	0	25	8	119	52

OFFICIAL

<b>05W</b>	NHS Stoke on Trent CCG	101	63	184	31	91	38	21	#	31	#	115	0
<b>05X</b>	NHS Telford and Wrekin CCG	93	68	232	133	61	26	31	18	29	11	93	18
<b>05Y</b>	NHS Walsall CCG	158	122	150	#	33	0	22	#	19	0	152	40
<b>06A</b>	NHS Wolverhampton CCG	76	41	346	202	47	0	19	0	49	24	231	122
<b>06D</b>	NHS Wyre Forest CCG	24	7	155	81	31	#	14	5	10	0	86	33
<b>06F</b>	NHS Bedfordshire CCG	45	0	300	38	94	5	17	0	37	0	120	0
<b>06H</b>	NHS Cambridgeshire and Peterborough CCG	101	0	812	316	205	38	85	21	119	33	143	0
<b>06K</b>	NHS East and North Hertfordshire CCG	69	0	516	191	119	12	65	25	101	45	228	0
<b>06L</b>	NHS Ipswich and East Suffolk CCG	15	0	469	225	75	0	22	0	36	0	280	104
<b>06M</b>	NHS Great Yarmouth and Waveney CCG	66	32	157	7	74	18	10	0	16	0	141	35
<b>06N</b>	NHS Herts Valleys CCG	154	69	398	54	145	34	63	20	53	0	310	49
<b>06P</b>	NHS Luton CCG	17	0	83	0	27	0	11	0	15	0	69	0
<b>06Q</b>	NHS Mid Essex CCG	69	14	156	0	76	0	26	0	47	7	131	0
<b>06T</b>	NHS North East Essex CCG	67	20	111	0	70	0	31	7	27	0	201	52
<b>06V</b>	NHS North Norfolk CCG	36	9	46	0	62	14	11	0	8	0	110	26
<b>06W</b>	NHS Norwich CCG	30	#	35	0	56	16	14	0	5	0	130	38
<b>06Y</b>	NHS South Norfolk CCG	30	0	92	0	74	22	19	#	11	0	151	50
<b>07G</b>	NHS Thurrock CCG	22	0	81	0	23	0	8	0	12	0	42	0
<b>07H</b>	NHS West Essex CCG	41	0	217	43	60	#	33	12	30	0	124	0

OFFICIAL

<b>07J</b>	NHS West Norfolk CCG	37	11	100	0	56	13	11	0	19	0	106	26
<b>07K</b>	NHS West Suffolk CCG	31	0	220	66	68	12	35	17	29	#	202	91
<b>07L</b>	NHS Barking and Dagenham CCG	30	5	144	54	25	0	44	30	22	7	116	40
<b>07M</b>	NHS Barnet CCG	26	0	150	0	35	0	9	0	35	0	138	0
<b>07N</b>	NHS Bexley CCG	27	0	133	#	42	0	16	0	19	0	113	16
<b>07P</b>	NHS Brent CCG	29	0	164	0	20	0	28	#	26	0	360	220
<b>07Q</b>	NHS Bromley CCG	64	17	206	12	91	28	39	15	70	37	211	66
<b>07R</b>	NHS Camden CCG	#	0	25	0	9	0	#	0	10	0	101	#
<b>07T</b>	NHS City and Hackney CCG	#	0	73	0	25	0	44	23	31	11	124	13
<b>07V</b>	NHS Croydon CCG	77	25	257	57	53	0	54	27	43	9	153	0
<b>07W</b>	NHS Ealing CCG	23	0	134	0	32	0	32	#	15	0	421	258
<b>07X</b>	NHS Enfield CCG	46	#	153	0	16	0	15	0	25	0	137	9
<b>07Y</b>	NHS Hounslow CCG	16	0	129	0	21	0	26	5	24	0	337	219
<b>08A</b>	NHS Greenwich CCG	33	0	172	41	38	0	26	6	22	0	124	15
<b>08C</b>	NHS Hammersmith and Fulham CCG	15	0	51	0	13	0	18	#	6	0	209	127
<b>08D</b>	NHS Haringey CCG	29	0	142	0	30	0	18	0	19	0	143	23
<b>08E</b>	NHS Harrow CCG	13	0	103	0	22	0	17	0	24	#	293	190
<b>08F</b>	NHS Havering CCG	59	23	257	104	51	0	51	32	40	14	116	#
<b>08G</b>	NHS Hillingdon CCG	48	10	145	0	58	11	39	19	25	0	190	72
<b>08H</b>	NHS Islington CCG	27	0	98	0	28	0	21	#	30	13	114	21
<b>08J</b>	NHS Kingston CCG	74	48	108	8	31	0	11	0	17	0	68	0
<b>08K</b>	NHS Lambeth CCG	33	0	175	7	32	0	42	14	32	#	196	46

OFFICIAL

<b>08L</b>	NHS Lewisham CCG	53	14	211	66	25	0	24	#	45	21	100	0
<b>08M</b>	NHS Newham CCG	12	0	82	0	6	0	24	0	21	0	215	83
<b>08N</b>	NHS Redbridge CCG	38	#	174	32	32	0	52	31	29	5	168	52
<b>08P</b>	NHS Richmond CCG	59	29	85	0	38	#	12	0	19	0	103	14
<b>08Q</b>	NHS Southwark CCG	40	#	129	0	28	0	33	11	25	#	148	28
<b>08R</b>	NHS Merton CCG	20	0	113	6	29	0	20	5	29	11	135	47
<b>08T</b>	NHS Sutton CCG	17	0	158	57	34	#	36	23	42	24	93	15
<b>08V</b>	NHS Tower Hamlets CCG	17	0	143	38	19	0	27	6	17	0	150	49
<b>08W</b>	NHS Waltham Forest CCG	34	0	76	0	23	0	36	15	33	10	130	16
<b>08X</b>	NHS Wandsworth CCG	51	6	148	0	37	0	38	11	35	8	262	115
<b>08Y</b>	NHS West London CCG	12	0	59	0	18	0	7	0	8	0	188	86
<b>09A</b>	NHS Central London (Westminster) CCG	9	0	44	0	14	0	6	0	10	0	109	26
<b>09C</b>	NHS Ashford CCG	27	9	31	0	34	8	11	#	7	0	41	0
<b>09D</b>	NHS Brighton and Hove CCG	53	13	237	84	77	29	58	36	67	41	180	57
<b>09E</b>	NHS Canterbury and Coastal CCG	47	18	93	0	63	18	52	36	19	0	34	0
<b>09F</b>	NHS Eastbourne, Hailsham and Seaford CCG	74	46	190	63	43	0	36	22	25	#	97	9
<b>09G</b>	NHS Coastal West Sussex CCG	191	117	546	214	155	36	55	19	103	45	203	0
<b>09H</b>	NHS Crawley CCG	47	30	146	80	18	0	10	#	18	6	44	0
<b>09J</b>	NHS Dartford, Gravesham and Swanley CCG	74	39	186	43	54	6	36	18	35	10	165	57
<b>09L</b>	NHS East Surrey CCG	79	54	192	91	40	6	13	0	20	#	113	37

OFFICIAL

<b>09N</b>	NHS Guildford and Waverley CCG	48	18	205	79	62	19	18	#	45	23	107	14
<b>09P</b>	NHS Hastings and Rother CCG	105	77	220	98	81	36	60	46	45	23	71	0
<b>09W</b>	NHS Medway CCG	82	43	248	90	50	0	46	26	46	18	80	0
<b>09X</b>	NHS Horsham and Mid Sussex CCG	87	53	236	94	71	23	46	29	36	11	92	0
<b>09Y</b>	NHS North West Surrey CCG	82	32	325	121	68	#	17	0	56	21	204	50
<b>10A</b>	NHS South Kent Coast CCG	64	34	92	0	76	29	51	36	25	#	72	0
<b>10C</b>	NHS Surrey Heath CCG	30	16	85	29	29	10	5	0	11	#	44	#
<b>10D</b>	NHS Swale CCG	34	19	145	83	21	0	19	11	21	10	18	0
<b>10E</b>	NHS Thanet CCG	34	14	74	0	52	20	19	9	19	#	41	0
<b>10J</b>	NHS North Hampshire CCG	49	18	202	76	41	0	19	#	26	#	11	0
<b>10K</b>	NHS Fareham and Gosport CCG	33	#	200	74	65	20	21	6	49	27	7	0
<b>10L</b>	NHS Isle of Wight CCG	21	0	148	51	33	0	12	#	18	0	#	0
<b>10Q</b>	NHS Oxfordshire CCG	189	94	681	292	183	52	50	0	105	38	256	0
<b>10R</b>	NHS Portsmouth CCG	28	0	177	67	42	6	15	0	44	25	8	0
<b>10V</b>	NHS South Eastern Hampshire CCG	38	7	230	96	75	27	20	5	43	19	13	0
<b>10X</b>	NHS Southampton CCG	50	17	213	85	26	0	48	29	34	12	18	0
<b>11A</b>	NHS West Hampshire CCG	158	78	593	246	164	40	84	45	88	27	41	0
<b>11E</b>	NHS Bath and North East Somerset CCG	76	49	119	6	58	19	13	0	18	0	89	6
<b>11J</b>	NHS Dorset CCG	493	381	1049	548	321	140	79	23	135	48	346	0
<b>11M</b>	NHS Gloucestershire CCG	68	0	706	324	187	51	52	8	96	29	146	0
<b>11N</b>	NHS Kernow CCG	225	142	677	314	246	111	62	21	103	38	483	226

OFFICIAL

<b>11X</b>	NHS Somerset CCG	135	53	715	357	216	85	18	0	158	95	210	0
<b>12D</b>	NHS Swindon CCG	12	0	126	#	50	9	27	11	31	9	101	6
<b>12F</b>	NHS Wirral CCG	126	79	524	324	113	42	22	0	34	0	303	158
<b>13T</b>	NHS Newcastle Gateshead CCG	27	0	346	84	136	48	93	58	86	40	289	88
<b>14L</b>	NHS Manchester CCG	103	35	368	115	79	#	51	10	67	24	157	0
<b>14Y</b>	NHS Buckinghamshire CCG	64	0	317	#	140	33	15	0	46	0	156	0
<b>15A</b>	NHS Berkshire West CCG	137	67	329	52	83	0	24	0	65	17	105	0
<b>15C</b>	NHS Bristol, North Somerset and South Gloucestershire CCG	342	216	823	305	343	170	64	0	141	52	115	0
<b>15D</b>	NHS Berkshire East CCG	122	64	215	0	66	0	10	0	29	0	128	0
<b>15E</b>	NHS Birmingham and Solihull CCG	246	90	894	262	224	19	84	0	121	13	1143	655
<b>15F</b>	NHS Leeds CCG	335	230	880	456	268	129	205	147	181	108	601	271
<b>99A</b>	NHS Liverpool CCG	154	88	327	66	91	#	45	9	80	34	143	0
<b>99C</b>	NHS North Tyneside CCG	22	0	190	61	70	25	43	27	31	8	98	#
<b>99D</b>	NHS South Lincolnshire CCG	93	69	48	0	36	0	15	#	16	0	52	0
<b>99E</b>	NHS Basildon and Brentwood CCG	58	21	153	0	54	#	10	0	10	0	109	0
<b>99F</b>	NHS Castle Point and Rochford CCG	69	42	170	54	64	22	29	16	17	0	164	82
<b>99G</b>	NHS Southend CCG	59	33	181	75	50	14	36	23	22	#	164	85
<b>99H</b>	NHS Surrey Downs CCG	77	34	269	89	95	34	45	24	60	29	147	16
<b>99J</b>	NHS West Kent CCG	159	92	236	0	99	#	70	37	75	26	109	0

OFFICIAL

<b>99K</b>	NHS High Weald Lewes Havens CCG	74	48	122	14	50	11	24	12	26	6	71	0
<b>99M</b>	NHS North East Hampshire and Farnham CCG	112	81	172	46	63	20	16	0	30	8	42	0
<b>99N</b>	NHS Wiltshire CCG	328	257	41	0	177	69	36	#	90	37	124	0
<b>99P</b>	NHS Northern, Eastern and Western Devon CCG	233	105	667	106	316	113	95	31	186	88	507	107
<b>99Q</b>	NHS South Devon and Torbay CCG	66	22	288	95	131	59	12	0	42	8	107	0
<b>Total</b>		13,930	6,807	44,497	14,950	14,376	4,113	6,219	2,509	7,789	2,582	28,846	8,633

Table 5<sup>21</sup>: 2017/18 STP activity and activity reduction for Category 1 interventions (A-D) <sup>22</sup>

STP code	STP name	A Surgery for snoring		B Dilatation & curettage for heavy menstrual bleeding		C Knee arthroscopy with osteoarthritis		D Injection for nonspecific low back pain without sciatica	
		2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
QE1	Healthier Lancashire and South Cumbria STP	68	68	12	12	158	158	1,359	1,359
QF7	South Yorkshire and Bassetlaw STP	35	35	#	#	195	195	721	721
QGH	Herefordshire and Worcestershire STP	#	#	6	6	40	40	109	109
QH8	Mid and South Essex STP	16	16	#	#	106	106	462	462
QHG	Bedfordshire, Luton and Milton Keynes STP	5	5	#	#	61	61	158	158
QHL	Birmingham and Solihull STP	#	#	5	5	89	89	307	307
QHM	Cumbria and North East STP	60	60	9	9	197	197	853	853
QJ2	Joined Up Care Derbyshire STP	19	19	#	#	50	50	364	364
QJG	Suffolk and North East Essex STP	9	9	#	#	53	53	333	333
QJK	Devon STP	19	19	#	#	36	36	246	246
QJM	Lincolnshire STP	8	8	#	#	104	104	426	426
QK1	Leicester, Leicestershire and Rutland STP	6	6	0	0	65	65	222	222
QKK	Our Healthier South East London STP	31	31	#	#	72	72	440	440
QKS	Kent and Medway STP	24	24	5	5	135	135	852	852

<sup>21</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>22</sup> STP activity figures are the sum of CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>QM7</b>	Hertfordshire and West Essex STP	27	27	5	5	65	65	171	171
<b>QMF</b>	East London Health & Care Partnership (STP)	28	28	5	5	75	75	414	414
<b>QMJ</b>	North London Partners in Health & Care (STP)	20	20	#	#	35	35	298	298
<b>QMM</b>	Norfolk and Waveney Health & Care Partnership (STP)	52	52	9	9	40	40	212	212
<b>QNC</b>	Staffordshire and Stoke on Trent STP	7	7	#	#	77	77	94	94
<b>QNN</b>	Frimley Health & Care ICS (STP)	5	5	#	#	30	30	89	89
<b>QNX</b>	Sussex and East Surrey STP	23	23	8	8	97	97	822	822
<b>QOC</b>	Shropshire and Telford and Wrekin STP	#	#	#	#	20	20	23	23
<b>QOP</b>	Greater Manchester Health and Social Care Partnership (STP)	36	36	8	8	172	172	479	479
<b>QOQ</b>	Humber, Coast and Vale STP	34	34	#	#	155	155	121	121
<b>QOX</b>	Bath and North East Somerset, Swindon and Wiltshire STP	9	9	20	20	103	103	40	40
<b>QPM</b>	Northamptonshire STP	5	5	13	13	63	63	124	124
<b>QR1</b>	Gloucestershire STP	7	7	#	#	24	24	12	12
<b>QRL</b>	Hampshire and the Isle of Wight STP	17	17	#	#	84	84	108	108
<b>QRV</b>	North West London Health & Care Partnership (STP)	18	18	21	21	148	148	229	229
<b>QSL</b>	Somerset STP	#	#	#	#	26	26	7	7
<b>QT1</b>	Nottingham and Nottinghamshire Health and Care STP	5	5	#	#	40	40	744	744
<b>QT6</b>	Cornwall and the Isles of Scilly Health & Social Care Partnership (STP)	#	#	#	#	45	45	29	29
<b>QU9</b>	Buckinghamshire, Oxfordshire and Berkshire West STP	9	9	#	#	63	63	38	38
<b>QUA</b>	The Black Country and West Birmingham STP	11	11	7	7	139	139	203	203
<b>QUE</b>	Cambridgeshire and Peterborough STP	9	9	#	#	31	31	171	171
<b>QUY</b>	Bristol, North Somerset and South Gloucestershire STP	7	7	7	7	41	41	12	12
<b>QVV</b>	Dorset STP	7	7	#	#	29	29	42	42

OFFICIAL

<b>QWE</b>	South West London Health & Care Partnership (STP)	13	13	#	#	33	33	448	448
<b>QWO</b>	West Yorkshire and Harrogate (Health & Care Partnership) STP	70	70	9	9	243	243	245	245
<b>QWU</b>	Coventry and Warwickshire STP	27	27	12	12	92	92	121	121
<b>QXU</b>	Surrey Heartlands Health & Care Partnership (STP)	6	6	7	7	15	15	467	467
<b>QYG</b>	Cheshire and Merseyside STP	46	46	12	12	91	91	550	550
<b>Total</b>		812	812	236	236	3,437	3,437	13,165	13,165

Table 6<sup>23</sup>: 2017/18 STP activity and activity reduction for Category 2 interventions (E-K) <sup>24</sup>

STP code	STP name	E Breast reduction		F Removal of benign skin lesions		G Grommets		H Tonsillectomy		I Haemorrhoid surgery		J Hysterectomy for heavy bleeding		K Chalazia removal	
		No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells
		2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
QE1	Healthier Lancashire and South Cumbria STP	71	20	4,804	2,522	289	124	1,179	413	278	110	844	159	159	189
QF7	South Yorkshire and Bassetlaw STP	73	29	4,362	2,460	252	99	637	85	277	123	876	290	290	362
QGH	Herefordshire and Worcestershire STP	27	7	1,507	559	208	134	319	16	82	8	480	164	164	0
QH8	Mid and South Essex STP	22	0	1,836	326	236	112	951	411	205	77	624	138	138	125
QHG	Bedfordshire, Luton and Milton Keynes STP	30	9	1,876	730	188	74	591	121	135	33	386	27	27	43
QHL	Birmingham and Solihull STP	57	23	1,858	310	198	48	660	5	112	0	616	162	162	98
QHM	Cumbria and North East STP	228	129	8,698	4,686	717	418	1,585	299	412	112	2,134	859	859	262
QJ2	Joined Up Care Derbyshire STP	36	#	2,044	672	133	34	597	151	126	16	752	330	330	87
QJG	Suffolk and North East Essex STP	43	13	2,168	896	162	68	758	338	135	29	484	79	79	46

<sup>23</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>24</sup> STP activity figures are the sum of CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>QJK</b>	Devon STP	37	#	2,938	1,271	299	192	547	44	121	10	759	262	262	0
<b>QJM</b>	Lincolnshire STP	50	27	1,483	409	127	53	504	176	91	22	426	102	102	0
<b>QK1</b>	Leicester, Leicestershire and Rutland STP	47	14	1,794	399	185	73	701	180	144	31	594	174	174	167
<b>QKK</b>	Our Healthier South East London STP	74	25	4,254	1,996	157	32	1,048	186	388	189	778	111	111	134
<b>QKS</b>	Kent and Medway STP	63	16	3,666	1,468	252	66	1,407	561	401	208	1,168	425	425	34
<b>QM7</b>	Hertfordshire and West Essex STP	46	10	2,064	242	303	139	1,134	431	321	160	577	6	6	184
<b>QMF</b>	East London Health & Care Partnership (STP)	39	#	3,518	1,295	211	8	1,313	291	400	202	561	29	29	561
<b>QMJ</b>	North London Partners in Health & Care (STP)	45	5	3,836	1,988	92	0	723	32	236	83	304	0	0	225
<b>QMM</b>	Norfolk and Waveney Health & Care Partnership (STP)	23	#	3,520	2,077	203	110	630	202	141	30	540	119	119	68
<b>QNC</b>	Staffordshire and Stoke on Trent STP	44	9	2,179	702	175	64	646	153	148	30	782	325	325	#
<b>QNQ</b>	Frimley Health & Care ICS (STP)	22	#	2,046	1,089	118	32	330	15	99	18	365	63	63	73
<b>QNX</b>	Sussex and East Surrey STP	73	19	4,775	2,272	274	96	1,293	478	310	109	1,066	308	308	24
<b>QOC</b>	Shropshire and Telford and Wrekin STP	16	#	644	15	59	13	161	0	46	0	250	47	47	0
<b>QOP</b>	Greater Manchester Health and Social Care Partnership (STP)	87	14	4,731	1,161	466	150	1,951	502	419	124	1,404	305	305	324
<b>QOQ</b>	Humber, Coast and Vale STP	105	60	2,339	752	270	152	1,002	395	199	55	740	189	189	19
<b>QOX</b>	Bath and North East Somerset, Swindon and Wiltshire STP	34	10	1,143	7	99	14	379	16	132	35	425	53	53	0

OFFICIAL

<b>QPM</b>	Northamptonshire STP	21	#	1,069	124	104	24	241	0	150	72	478	184	184	54
<b>QR1</b>	Gloucestershire STP	30	11	531	0	65	#	275	0	52	0	242	0	0	#
<b>QRL</b>	Hampshire and the Isle of Wight STP	48	#	2,505	289	200	35	826	55	170	19	965	232	232	16
<b>QRV</b>	North West London Health & Care Partnership (STP)	106	43	2,878	278	178	19	885	15	421	174	597	31	31	207
<b>QSL</b>	Somerset STP	10	0	1,170	382	94	43	234	#	63	#	301	66	66	0
<b>QT1</b>	Nottingham and Nottinghamshire Health and Care STP	50	18	2,785	1,449	184	78	525	81	136	33	471	87	87	61
<b>QT6</b>	Cornwall and the Isles of Scilly Health & Social Care Partnership (STP)	31	13	1,312	518	101	51	239	9	78	17	234	0	0	0
<b>QU9</b>	Buckinghamshire, Oxfordshire and Berkshire West STP	83	31	2,919	990	240	72	640	0	136	#	500	0	0	0
<b>QUA</b>	The Black Country and West Birmingham STP	59	19	2,912	1,229	186	44	847	149	191	60	819	300	300	86
<b>QUE</b>	Cambridgeshire and Peterborough STP	18	0	1,276	94	155	60	557	116	72	0	389	34	34	55
<b>QUY</b>	Bristol, North Somerset and South Gloucestershire STP	28	0	2,591	1,354	105	7	217	0	130	30	325	0	0	39
<b>QVV</b>	Dorset STP	31	7	3,295	2,185	48	0	255	0	101	16	496	174	174	109
<b>QWE</b>	South West London Health & Care Partnership (STP)	91	45	1,565	0	127	9	741	42	301	128	575	85	85	32
<b>QWO</b>	West Yorkshire and Harrogate (Health	142	65	4,359	1,110	630	345	1,936	674	534	263	1,183	204	204	201

OFFICIAL

	& Care Partnership) STP														
<b>QWU</b>	Coventry and Warwickshire STP	50	21	2,643	1,404	119	30	743	292	76	0	470	104	104	102
<b>QXU</b>	Surrey Heartlands Health & Care Partnership (STP)	47	21	1,859	695	89	#	440	64	161	66	391	38	38	34
<b>QYG</b>	Cheshire and Merseyside STP	151	77	6,503	3,184	371	130	1,591	454	344	105	1,289	271	271	298
<b>Total</b>		2,388	829	116,255	45,589	8,669	3,259	32,238	7,454	8,474	2,801	27,660	6,536	6,026	4,326

Table 7<sup>25</sup>: 2017/18 STP activity and activity reduction for Category 2 interventions (L-Q) <sup>26</sup>

		L Shoulder decompression		M Carpal tunnel syndrome release		N Dupuytren's contracture release		O Ganglion excision		P Trigger finger release		Q Varicose vein surgery	
		No of spells		No of spells		No of spells		No of spells		No of spells		No of spells	
STP code	STP name	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction	2017/18 activity	Activity reduction
QE1	Healthier Lancashire and South Cumbria STP	423	184	1,482	507	650	289	271	149	263	90	1,404	668
QF7	South Yorkshire and Bassetlaw STP	456	250	1,358	515	430	136	312	205	292	143	380	14
QGH	Herefordshire and Worcestershire STP	148	52	1,147	661	229	51	130	72	103	19	492	142
QH8	Mid and South Essex STP	277	110	741	129	267	39	109	39	108	10	610	167
QHG	Bedfordshire, Luton and Milton Keynes STP	180	81	630	144	172	11	52	#	87	10	261	0
QHL	Birmingham and Solihull STP	246	90	894	262	224	19	84	0	121	13	1,143	655
QHM	Cumbria and North East STP	385	45	2,646	820	985	321	492	266	494	168	2,407	1,036
QJ2	Joined Up Care Derbyshire STP	330	183	1,199	589	357	139	150	77	211	101	433	23

<sup>25</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>26</sup> STP activity figures are the sum of CCG activity figures excluding activity that could not be attributed to a CCG.

OFFICIAL

<b>QJG</b>	Suffolk and North East Essex STP	113	20	800	291	213	12	88	24	92	#	683	247
<b>QJK</b>	Devon STP	299	127	955	201	447	172	107	31	228	96	614	107
<b>QJM</b>	Lincolnshire STP	343	230	127	0	198	20	39	#	32	0	309	11
<b>QK1</b>	Leicester, Leicestershire and Rutland STP	175	40	752	159	178	11	72	13	115	12	203	0
<b>QKK</b>	Our Healthier South East London STP	250	33	1,026	129	256	28	180	48	213	64	892	171
<b>QKS</b>	Kent and Medway STP	521	268	1,105	216	449	84	304	175	247	69	560	57
<b>QM7</b>	Hertfordshire and West Essex STP	264	69	1,131	288	324	48	161	57	184	45	662	49
<b>QMF</b>	East London Health & Care Partnership (STP)	194	29	949	228	181	0	278	137	193	47	1,019	256
<b>QMJ</b>	North London Partners in Health & Care (STP)	132	#	568	0	118	0	67	#	119	13	633	56
<b>QMM</b>	Norfolk and Waveney Health & Care Partnership (STP)	199	53	430	7	322	83	65	#	59	0	638	175
<b>QNC</b>	Staffordshire and Stoke on Trent STP	309	166	791	136	391	152	73	#	146	27	738	247
<b>QNQ</b>	Frimley Health & Care ICS (STP)	264	161	472	75	158	30	31	0	70	9	214	#
<b>QNX</b>	Sussex and East Surrey STP	710	438	1,889	738	535	141	302	165	340	137	871	103
<b>QOC</b>	Shropshire and Telford and Wrekin STP	256	185	654	355	180	71	93	58	93	39	282	66
<b>QOP</b>	Greater Manchester Health and Social Care Partnership (STP)	803	417	2,532	1,087	730	226	388	184	476	208	1,427	432
<b>QOQ</b>	Humber, Coast and Vale STP	585	386	1,339	527	500	200	108	18	258	116	905	324
<b>QOX</b>	Bath and North East Somerset, Swindon and Wiltshire STP	416	306	286	9	285	97	76	13	139	46	314	12
<b>QPM</b>	Northamptonshire STP	213	112	496	114	192	53	93	41	102	30	187	0
<b>QR1</b>	Gloucestershire STP	68	0	706	324	187	51	52	8	96	29	146	0
<b>QRL</b>	Hampshire and the Isle of Wight STP	377	123	1,763	695	446	93	219	89	302	114	102	0

OFFICIAL

<b>QRV</b>	North West London Health & Care Partnership (STP)	165	10	829	0	198	11	173	33	138	#	2,107	1,198
<b>QSL</b>	Somerset STP	135	53	715	357	216	85	18	0	158	95	210	0
<b>QT1</b>	Nottingham and Nottinghamshire Health and Care STP	382	242	900	335	312	116	106	33	181	80	181	0
<b>QT6</b>	Cornwall and the Isles of Scilly Health & Social Care Partnership (STP)	225	142	677	314	246	111	62	21	103	38	483	226
<b>QU9</b>	Buckinghamshire, Oxfordshire and Berkshire West STP	390	161	1,327	347	406	85	89	0	216	55	517	0
<b>QUA</b>	The Black Country and West Birmingham STP	355	184	1,010	395	208	6	86	8	124	31	1,047	486
<b>QUE</b>	Cambridgeshire and Peterborough STP	101	0	812	316	205	38	85	21	119	33	143	0
<b>QUY</b>	Bristol, North Somerset and South Gloucestershire STP	342	216	823	305	343	170	64	0	141	52	115	0
<b>QVV</b>	Dorset STP	493	381	1,049	548	321	140	79	23	135	48	346	0
<b>QWE</b>	South West London Health & Care Partnership (STP)	298	108	869	128	222	#	171	66	185	52	814	191
<b>QWO</b>	West Yorkshire and Harrogate (Health & Care Partnership) STP	871	525	2,433	1,035	882	423	501	322	422	189	2,015	951
<b>QWU</b>	Coventry and Warwickshire STP	236	108	1,054	527	209	38	90	26	135	43	391	22
<b>QXU</b>	Surrey Heartlands Health & Care Partnership (STP)	207	84	799	289	225	54	80	26	161	73	458	80
<b>QYG</b>	Cheshire and Merseyside STP	794	431	2,332	848	779	255	219	47	388	135	1,490	458
<b>Total</b>		13,930	6,807	44,497	14,950	14,376	4,113	6,219	2,509	7,789	2,582	28,846	8,633

Table 8<sup>27</sup>: 2017/18 provider activity baseline estimate for Category 1 interventions (A-D) and total category 1 activity <sup>28</sup>

Provider code	Provider name	A Surgery for snoring	B Dilatation & curettage for heavy menstrual bleeding	C Knee arthroscopy with osteoarthritis	D Injection for nonspecific low back pain without sciatica
		No of spells	No of spells	No of spells	No of spells
		2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity
8G301	RYALLS ANN (COUNSELLOR)	0	0	0	7
8G326	THE SPENCER WING	0	0	0	27
AAH01	TETBURY HOSPITAL TRUST	#	0	0	0
ADP02	KIMS HOSPITAL (NEWNHAM COURT)	0	0	14	52
AHH01	FOSCOTE COURT (BANBURY) TRUST	0	0	0	0
AJX01	SUSSEX MSK PARTNERSHIP 2 LTD	0	0	11	#
AVQ01	ONE ASHFORD HOSPITAL	0	0	0	0
RXG00	WILTSHIRE HEATH AND CARE LLP	0	0	0	0
NAM01	PROBUS SURGICAL CENTRE	0	0	0	0
NAM02	MENEAGE STREET SURGERY	0	0	0	0
NAM04	LISKEARD COMMUNITY HOSPITAL	0	0	0	0
NAM06	MORRAB SURGERY	0	0	0	0
NEQ01	PHOENIX HEALTH SOLUTIONS LIMITED	0	0	0	0
NFH01	SOMERSET SURGICAL SERVICES	0	0	#	0
NHW03	WIMBOURNE COMMUNITY HOSPITAL (STANDARD HEALTH)	0	0	#	0
NN401	TYNESIDE SURGICAL SERVICES	0	0	#	16

<sup>27</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>28</sup> Provider activity figures include activity that could not be attributed to a CCG.

OFFICIAL

<b>NN501</b>	TRAFFORD GENERAL HOSPITAL	#	0	0	0
<b>NN801</b>	THE SPENCER WING (RAMSGATE ROAD)	0	0	0	33
<b>NN802</b>	THE SPENCER WING (WILLIAM HARVEY HOSPITAL)	0	0	0	9
<b>NNE02</b>	DORKING GENERAL HOSPITAL	0	0	#	57
<b>NQM01</b>	ORTHOPAEDICS & SPINE SPECIALIST HOSPITAL SITE	0	0	#	0
<b>NT202</b>	NUFFIELD HEALTH, BOURNEMOUTH HOSPITAL	0	0	0	0
<b>NT204</b>	NUFFIELD HEALTH, BRENTWOOD HOSPITAL	0	0	#	0
<b>NT206</b>	NUFFIELD HEALTH, BRISTOL HOSPITAL (CHESTERFIELD)	0	0	#	0
<b>NT209</b>	NUFFIELD HEALTH, CAMBRIDGE HOSPITAL	0	0	#	0
<b>NT210</b>	NUFFIELD HEALTH, THE GROSVENOR HOSPITAL, CHESTER	0	0	#	0
<b>NT211</b>	NUFFIELD HEALTH, CHELTENHAM HOSPITAL	0	0	0	0
<b>NT212</b>	NUFFIELD HEALTH, CHICHESTER HOSPITAL	0	0	#	16
<b>NT213</b>	NUFFIELD HEALTH, DERBY HOSPITAL	0	0	#	0
<b>NT214</b>	NUFFIELD HEALTH, WESSEX HOSPITAL	0	0	0	0
<b>NT215</b>	NUFFIELD HEALTH, EXETER HOSPITAL	0	0	0	0
<b>NT218</b>	NUFFIELD HEALTH, HAYWARDS HEATH HOSPITAL	0	0	0	0
<b>NT219</b>	NUFFIELD HEALTH, HEREFORD HOSPITAL	0	0	0	0
<b>NT224</b>	NUFFIELD HEALTH, WARWICKSHIRE HOSPITAL	0	0	#	0
<b>NT225</b>	NUFFIELD HEALTH, LEEDS HOSPITAL	11	0	8	0
<b>NT226</b>	NUFFIELD HEALTH, LEICESTER HOSPITAL	0	0	0	0
<b>NT229</b>	NUFFIELD HEALTH, NEWCASTLE UPON TYNE HOSPITAL	0	0	0	0
<b>NT230</b>	NUFFIELD HEALTH, NORTH STAFFORDSHIRE HOSPITAL	0	0	6	#

OFFICIAL

<b>NT233</b>	NUFFIELD HEALTH, PLYMOUTH HOSPITAL	0	0	0	0
<b>NT235</b>	NUFFIELD HEALTH, SHREWSBURY HOSPITAL	0	0	#	0
<b>NT237</b>	NUFFIELD HEALTH, TEES HOSPITAL	0	0	#	0
<b>NT238</b>	NUFFIELD HEALTH, TAUNTON HOSPITAL	#	0	#	#
<b>NT239</b>	NUFFIELD HEALTH, TUNBRIDGE WELLS HOSPITAL	0	0	0	#
<b>NT241</b>	NUFFIELD HEALTH, WOKING HOSPITAL	0	0	0	15
<b>NT242</b>	NUFFIELD HEALTH, WOLVERHAMPTON HOSPITAL	0	0	#	15
<b>NT245</b>	NUFFIELD HEALTH, YORK HOSPITAL	0	0	#	0
<b>NT301</b>	SPIRE SOUTH BANK HOSPITAL	0	0	9	0
<b>NT302</b>	SPIRE BRISTOL HOSPITAL	0	0	7	0
<b>NT304</b>	SPIRE SOUTHAMPTON HOSPITAL	0	0	16	0
<b>NT305</b>	SPIRE PORTSMOUTH HOSPITAL	0	0	#	0
<b>NT308</b>	SPIRE GATWICK PARK HOSPITAL	0	0	#	26
<b>NT309</b>	SPIRE SUSSEX HOSPITAL	0	0	#	#
<b>NT310</b>	SPIRE TUNBRIDGE WELLS HOSPITAL	0	0	#	0
<b>NT312</b>	SPIRE ALEXANDRA HOSPITAL	0	0	19	0
<b>NT313</b>	SPIRE WELLESLEY HOSPITAL	0	0	9	140
<b>NT314</b>	SPIRE LONDON EAST	#	0	17	159
<b>NT315</b>	SPIRE BUSHEY HOSPITAL	0	0	7	0
<b>NT316</b>	SPIRE HARPENDEN HOSPITAL	#	0	14	0
<b>NT317</b>	SPIRE CAMBRIDGE LEA HOSPITAL	0	0	7	0
<b>NT318</b>	SPIRE NORWICH HOSPITAL	0	0	#	0
<b>NT319</b>	SPIRE HARTSWOOD HOSPITAL	0	0	9	0
<b>NT320</b>	SPIRE PARKWAY HOSPITAL	0	0	8	#
<b>NT321</b>	SPIRE LITTLE ASTON HOSPITAL	0	0	69	0
<b>NT322</b>	SPIRE LEICESTER HOSPITAL	0	0	22	0
<b>NT324</b>	SPIRE CHESHIRE HOSPITAL	#	0	12	0
<b>NT325</b>	SPIRE MURRAYFIELD HOSPITAL	0	0	5	5

OFFICIAL

<b>NT327</b>	SPIRE MANCHESTER HOSPITAL	0	0	5	0
<b>NT332</b>	SPIRE LEEDS HOSPITAL	0	0	13	0
<b>NT333</b>	SPIRE WASHINGTON HOSPITAL	0	0	15	49
<b>NT337</b>	SPIRE LIVERPOOL HOSPITAL	#	0	#	12
<b>NT338</b>	SPIRE YALE HOSPITAL	0	0	0	0
<b>NT339</b>	SPIRE REGENCY HOSPITAL	0	0	0	9
<b>NT343</b>	SPIRE THAMES VALLEY HOSPITAL	0	0	5	0
<b>NT344</b>	SPIRE DUNEDIN HOSPITAL	0	0	#	0
<b>NT345</b>	SPIRE CLARE PARK HOSPITAL	0	0	5	13
<b>NT347</b>	SPIRE FYLDE COAST HOSPITAL	#	0	8	155
<b>NT348</b>	SPIRE ELLAND HOSPITAL	0	0	#	5
<b>NT350</b>	SPIRE METHLEY PARK HOSPITAL	0	0	43	9
<b>NT351</b>	SPIRE HULL AND EAST RIDING HOSPITAL	0	0	35	30
<b>NT364</b>	SPIRE MONTEFIORE HOSPITAL	0	0	0	47
<b>NT401</b>	BMI - THE ALEXANDRA HOSPITAL	0	0	10	9
<b>NT402</b>	BMI - BATH CLINIC	0	0	15	#
<b>NT403</b>	BMI - THE BEARDWOOD HOSPITAL	#	0	17	#
<b>NT404</b>	BMI - THE BEAUMONT HOSPITAL	#	0	6	#
<b>NT405</b>	BMI - BISHOPS WOOD	#	0	9	11
<b>NT406</b>	BMI - THE BLACKHEATH HOSPITAL	0	0	#	#
<b>NT408</b>	BMI - THE CHAUCER HOSPITAL	0	0	0	8
<b>NT409</b>	BMI - CHELSFIELD PARK HOSPITAL	0	0	#	5
<b>NT410</b>	BMI - THE CHILTERN HOSPITAL	0	0	#	0
<b>NT411</b>	BMI - THE CLEMENTINE CHURCHILL HOSPITAL	#	0	39	#
<b>NT412</b>	BMI - THE DROITWICH SPA HOSPITAL	0	0	10	0
<b>NT413</b>	BMI - THE ESPERANCE HOSPITAL	0	0	0	0
<b>NT414</b>	BMI - FAWKHAM MANOR HOSPITAL	0	0	0	#
<b>NT416</b>	BMI - HENDON HOSPITAL	0	0	#	0
<b>NT417</b>	BMI - GORING HALL HOSPITAL	0	0	#	8
<b>NT418</b>	BMI - THE HAMPSHIRE CLINIC	0	0	#	6

OFFICIAL

<b>NT419</b>	BMI - THE HARBOUR HOSPITAL	0	0	0	0
<b>NT420</b>	BMI - THE HIGHFIELD HOSPITAL	#	0	23	#
<b>NT421</b>	BMI - THE KINGS OAK HOSPITAL	#	0	0	#
<b>NT422</b>	BMI - THE LONDON INDEPENDENT HOSPITAL	0	0	7	9
<b>NT423</b>	BMI - THE MANOR HOSPITAL	0	0	0	0
<b>NT424</b>	BMI - THE MERIDEN HOSPITAL	#	0	10	0
<b>NT427</b>	BMI - THE PARK HOSPITAL	0	0	#	#
<b>NT428</b>	BMI - THE PRINCESS MARGARET HOSPITAL	0	0	#	0
<b>NT429</b>	BMI - THE PRIORY HOSPITAL	0	0	5	0
<b>NT430</b>	BMI - THE RIDGEWAY HOSPITAL	0	0	#	0
<b>NT431</b>	BMI - THE RUNNYMEDE HOSPITAL	0	0	#	7
<b>NT432</b>	BMI - THE SANDRINGHAM HOSPITAL	0	0	#	0
<b>NT433</b>	BMI - SARUM ROAD HOSPITAL	0	0	6	0
<b>NT434</b>	BMI - THE SAXON CLINIC	0	0	#	#
<b>NT435</b>	BMI - THE SHELBURNE HOSPITAL	0	0	#	0
<b>NT436</b>	BMI - SHIRLEY OAKS HOSPITAL	0	0	5	12
<b>NT437</b>	BMI - THE SLOANE HOSPITAL	0	0	#	#
<b>NT438</b>	BMI - THE SOMERFIELD HOSPITAL	0	0	#	5
<b>NT439</b>	BMI - THE SOUTH CHESHIRE PRIVATE HOSPITAL	0	#	7	0
<b>NT440</b>	BMI - THORNBURY HOSPITAL	0	0	0	0
<b>NT441</b>	BMI - THREE SHIRES HOSPITAL	0	0	9	0
<b>NT443</b>	BMI - THE WINTERBOURNE HOSPITAL	0	0	#	0
<b>NT445</b>	BMI THE EDGBASTON HOSPITAL	0	0	#	0
<b>NT446</b>	BMI ST EDMUNDS HOSPITAL	0	0	#	0
<b>NT447</b>	BMI THE DUCHY HOSPITAL	0	0	5	#
<b>NT448</b>	BMI THE HUDDERSFIELD HOSPITAL	8	0	21	0
<b>NT449</b>	BMI THE LANCASTER HOSPITAL	0	0	11	9
<b>NT450</b>	BMI THE LINCOLN HOSPITAL	0	0	22	0
<b>NT451</b>	BMI THE CAVELL HOSPITAL	0	0	#	#

OFFICIAL

<b>NT455</b>	BMI MOUNT ALVERNIA HOSPITAL	0	0	0	6
<b>NT457</b>	BMI WOODLANDS HOSPITAL	0	0	5	7
<b>NT490</b>	BMI SOUTHEND PRIVATE HOSPITAL	0	0	0	0
<b>NT497</b>	BMI GISBURNE PARK HOSPITAL	#	0	31	#
<b>NTE02</b>	ST HUGH'S HOSPITAL	0	0	54	0
<b>NTP11</b>	SOUTHAMPTON NHS TREATMENT CENTRE	#	0	12	#
<b>NTP13</b>	BARLBOROUGH NHS TREATMENT CENTRE	0	0	7	6
<b>NTP15</b>	NORTH EAST LONDON TREATMENT CENTRE CARE UK	#	0	#	0
<b>NTP16</b>	WILL ADAMS NHS TREATMENT CENTRE	0	0	#	0
<b>NTPAD</b>	ST MARY'S NHS TREATMENT CENTRE	0	0	0	0
<b>NTPAE</b>	THE CROFT SHIFA HEALTH CENTRE	0	0	0	0
<b>NTPH1</b>	SHEPTON MALLET NHS TREATMENT CENTRE	0	0	8	#
<b>NTPH2</b>	EMERSONS GREEN NHS TREATMENT CENTRE	#	#	6	0
<b>NTPH3</b>	DEVIZES NHS TREATMENT CENTRE	0	#	#	0
<b>NTPH5</b>	PENINSULA NHS TREATMENT CENTRE	0	0	#	0
<b>NTX01</b>	ONE HEALTH GROUP LTD	0	0	0	#
<b>NTX06</b>	ONE HEALTH GROUP CLINIC - OSSET	0	0	0	#
<b>NTX09</b>	ONE HEALTH GROUP CLINIC - GAINSBOROUGH	0	0	0	#
<b>NTX11</b>	ONE HEALTH GROUP CLINIC - THORNBURY	0	0	#	5
<b>NTX12</b>	ONE HEALTH GROUP CLINIC - CLAREMONT	0	0	#	0
<b>NV302</b>	CIRCLE BATH HOSPITAL	0	18	13	#
<b>NV313</b>	CIRCLE - NOTTINGHAM NHS TREATMENT CENTRE	0	#	6	398
<b>NV323</b>	CIRCLE READING HOSPITAL	0	0	#	#
<b>NVC01</b>	ASHTHEAD HOSPITAL	0	0	#	34
<b>NVC02</b>	THE BERKSHIRE INDEPENDENT HOSPITAL	0	0	0	0

OFFICIAL

<b>NVC04</b>	DUCHY HOSPITAL	0	0	33	0
<b>NVC05</b>	EUXTON HALL HOSPITAL	0	#	10	11
<b>NVC06</b>	FITZWILLIAM HOSPITAL	#	0	8	100
<b>NVC07</b>	FULWOOD HALL HOSPITAL	#	#	#	11
<b>NVC08</b>	MOUNT STUART HOSPITAL	0	0	#	0
<b>NVC09</b>	NEW HALL HOSPITAL	#	0	#	#
<b>NVC0M</b>	RAMSAY CROYDON DAY HOSPITAL	0	0	0	0
<b>NVC0R</b>	TEES VALLEY HOSPITAL	0	0	0	0
<b>NVC11</b>	NORTH DOWNS HOSPITAL	0	0	#	17
<b>NVC12</b>	OAKLANDS HOSPITAL	#	0	18	0
<b>NVC13</b>	OAKS HOSPITAL	0	0	#	35
<b>NVC14</b>	PARK HILL HOSPITAL	#	0	6	36
<b>NVC15</b>	PINEHILL HOSPITAL	0	0	#	29
<b>NVC16</b>	RENACRES HOSPITAL	0	#	#	30
<b>NVC17</b>	ROWLEY HALL HOSPITAL	0	0	5	11
<b>NVC18</b>	SPRINGFIELD HOSPITAL	0	0	30	16
<b>NVC19</b>	RIVERS HOSPITAL	0	0	0	31
<b>NVC20</b>	THE YORKSHIRE CLINIC	16	0	#	51
<b>NVC21</b>	WEST MIDLANDS HOSPITAL	#	#	19	61
<b>NVC22</b>	WINFIELD HOSPITAL	0	0	#	0
<b>NVC23</b>	WOODLAND HOSPITAL	0	8	9	#
<b>NVC25</b>	HORTON NHS TREATMENT CENTRE	0	0	15	0
<b>NVC27</b>	BOSTON WEST HOSPITAL	0	#	13	22
<b>NVC28</b>	CLIFTON PARK HOSPITAL	0	0	26	0
<b>NVC29</b>	COBALT HOSPITAL	0	0	0	0
<b>NVC31</b>	BLAKELANDS HOSPITAL	0	0	0	0
<b>NVC35</b>	TEES VALLEY TREATMENT CENTRE	0	0	#	0
<b>NVC40</b>	WOODTHORPE HOSPITAL	0	0	#	5
<b>NVC44</b>	THE WESTBOURNE CENTRE	0	0	0	0
<b>NVG01</b>	FAIRFIELD HOSPITAL	0	0	#	0
<b>NVM01</b>	COBHAM DAY SURGERY HOSPITAL	0	0	0	0

OFFICIAL

<b>NVM02</b>	EPSOM DAY SURGERY LIMITED	0	0	0	0
<b>NW612</b>	HCA - 52 ALDERLEY ROAD	0	0	#	0
<b>NWF01</b>	BENENDEN HOSPITAL	0	0	#	25
<b>NWX11</b>	BICS-COMMUNITY	0	0	#	165
<b>NXM01</b>	THE HORDER CENTRE - ST JOHNS ROAD	0	0	16	11
<b>NXM04</b>	THE MCINDOE CENTRE	0	0	0	25
<b>NXP04</b>	HATHAWAY MEDICAL CENTRE	0	0	0	0
<b>NXP17</b>	WHITE HORSE HEALTH CENTRE - IHG	0	0	0	0
<b>NXP20</b>	LAWN MEDICAL CENTRE	0	0	0	0
<b>NXP37</b>	ASPEN CENTRE	0	0	0	0
<b>NXP40</b>	LITFIELD HOUSE MEDICAL CENTRE	0	0	0	0
<b>NXP41</b>	IHG CRAVEN ROAD	0	0	0	0
<b>NYW01</b>	ASPEN - THE HOLLY	#	0	9	16
<b>NYW02</b>	ASPEN - PARKSIDE HOSPITAL	0	0	0	6
<b>NYW03</b>	ASPEN - HIGHGATE HOSPITAL	0	0	#	20
<b>NYW04</b>	ASPEN - CLAREMONT HOSPITAL	0	0	#	7
<b>PHS</b>	PHOENIX HEALTH SOLUTIONS LTD	0	0	0	0
<b>R0A</b>	MANCHESTER UNIVERSITY NHS FOUNDATION TRUST	#	#	13	27
<b>R1D</b>	SHROPSHIRE COMMUNITY HEALTH NHS TRUST	0	0	0	0
<b>R1F</b>	ISLE OF WIGHT NHS TRUST	#	0	6	14
<b>R1H</b>	BARTS HEALTH NHS TRUST	12	#	17	77
<b>R1K</b>	LONDON NORTH WEST UNIVERSITY HEALTHCARE NHS TRUST	8	5	19	0
<b>RA2</b>	ROYAL SURREY COUNTY HOSPITAL NHS FOUNDATION TRUST	#	#	5	67
<b>RA3</b>	WESTON AREA HEALTH NHS TRUST	0	0	14	0
<b>RA4</b>	YEOVIL DISTRICT HOSPITAL NHS FOUNDATION TRUST	0	#	12	0
<b>RA7</b>	UNIVERSITY HOSPITALS BRISTOL NHS FOUNDATION TRUST	5	0	#	#
<b>RA9</b>	TORBAY AND SOUTH DEVON NHS FOUNDATION TRUST	6	0	13	179

OFFICIAL

<b>RAE</b>	BRADFORD TEACHING HOSPITALS NHS FOUNDATION TRUST	6	#	53	0
<b>RAJ</b>	SOUTHEND UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	10	0	12	271
<b>RAL</b>	ROYAL FREE LONDON NHS FOUNDATION TRUST	8	#	14	169
<b>RAN</b>	ROYAL NATIONAL ORTHOPAEDIC HOSPITAL NHS TRUST	0	0	8	35
<b>RAP</b>	NORTH MIDDLESEX UNIVERSITY HOSPITAL NHS TRUST	0	#	13	26
<b>RAS</b>	THE HILLINGDON HOSPITALS NHS FOUNDATION TRUST	0	#	41	74
<b>RAX</b>	KINGSTON HOSPITAL NHS FOUNDATION TRUST	#	0	#	25
<b>RBA</b>	TAUNTON AND SOMERSET NHS FOUNDATION TRUST	#	#	8	#
<b>RBD</b>	DORSET COUNTY HOSPITAL NHS FOUNDATION TRUST	#	0	#	0
<b>RBK</b>	WALSALL HEALTHCARE NHS TRUST	#	#	25	6
<b>RBL</b>	WIRRAL UNIVERSITY TEACHING HOSPITAL NHS FOUNDATION TRUST	#	0	7	37
<b>RBN</b>	ST HELENS AND KNOWSLEY HOSPITAL SERVICES NHS TRUST	9	#	6	31
<b>RBQ</b>	LIVERPOOL HEART AND CHEST HOSPITAL NHS FOUNDATION TRUST	0	0	0	0
<b>RBS</b>	ALDER HEY CHILDREN'S NHS FOUNDATION TRUST	0	0	0	#
<b>RBT</b>	MID CHESHIRE HOSPITALS NHS FOUNDATION TRUST	0	0	14	63
<b>RBV</b>	THE CHRISTIE NHS FOUNDATION TRUST	0	#	0	#
<b>RBZ</b>	NORTHERN DEVON HEALTHCARE NHS TRUST	0	#	#	14
<b>RC1</b>	BEDFORD HOSPITAL NHS TRUST	0	0	19	0
<b>RC9</b>	LUTON AND DUNSTABLE UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	6	#	10	86
<b>RCB</b>	YORK TEACHING HOSPITAL NHS FOUNDATION TRUST	#	#	8	7
<b>RCD</b>	HARROGATE AND DISTRICT NHS FOUNDATION TRUST	#	0	11	0

OFFICIAL

<b>RCF</b>	AIREDALE NHS FOUNDATION TRUST	0	0	#	#
<b>RCU</b>	SHEFFIELD CHILDREN'S NHS FOUNDATION TRUST	0	0	0	0
<b>RCX</b>	THE QUEEN ELIZABETH HOSPITAL, KING'S LYNN, NHS FOUNDATION TRUST	5	#	#	64
<b>RD1</b>	ROYAL UNITED HOSPITALS BATH NHS FOUNDATION TRUST	#	#	9	20
<b>RD3</b>	POOLE HOSPITAL NHS FOUNDATION TRUST	6	#	0	29
<b>RD8</b>	MILTON KEYNES UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	0	#	19	54
<b>RDD</b>	BASILDON AND THURROCK UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	5	0	10	14
<b>RDE</b>	EAST SUFFOLK AND NORTH ESSEX NHS FOUNDATION TRUST	#	#	12	#
<b>RDU</b>	FRIMLEY HEALTH NHS FOUNDATION TRUST	5	#	16	86
<b>RDY</b>	DORSET HEALTHCARE UNIVERSITY NHS FOUNDATION TRUST	0	0	#	0
<b>RDZ</b>	THE ROYAL BOURNEMOUTH AND CHRISTCHURCH HOSPITALS NHS FOUNDATION TRUST	0	#	8	5
<b>RE9</b>	SOUTH TYNESIDE NHS FOUNDATION TRUST	0	0	#	0
<b>REF</b>	ROYAL CORNWALL HOSPITALS NHS TRUST	#	#	10	22
<b>REM</b>	AINTREE UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	7	0	#	98
<b>REP</b>	LIVERPOOL WOMEN'S NHS FOUNDATION TRUST	0	#	0	0
<b>RET</b>	THE WALTON CENTRE NHS FOUNDATION TRUST	0	0	0	104
<b>RF4</b>	BARKING, HAVERING AND REDBRIDGE UNIVERSITY HOSPITALS NHS TRUST	7	#	16	27
<b>RFF</b>	BARNSLEY HOSPITAL NHS FOUNDATION TRUST	5	#	48	0
<b>RFR</b>	THE ROTHERHAM NHS FOUNDATION TRUST	7	#	28	0

OFFICIAL

<b>RFS</b>	CHESTERFIELD ROYAL HOSPITAL NHS FOUNDATION TRUST	12	0	20	205
<b>RGN</b>	NORTH WEST ANGLIA NHS FOUNDATION TRUST	6	#	29	181
<b>RGP</b>	JAMES PAGET UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	15	#	15	#
<b>RGQ</b>	IPSWICH HOSPITAL NHS TRUST	#	#	13	227
<b>RGR</b>	WEST SUFFOLK NHS FOUNDATION TRUST	#	0	30	94
<b>RGT</b>	CAMBRIDGE UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	#	#	#	#
<b>RH8</b>	ROYAL DEVON AND EXETER NHS FOUNDATION TRUST	9	#	#	61
<b>RHM</b>	UNIVERSITY HOSPITAL SOUTHAMPTON NHS FOUNDATION TRUST	#	0	7	#
<b>RHQ</b>	SHEFFIELD TEACHING HOSPITALS NHS FOUNDATION TRUST	22	0	7	58
<b>RHU</b>	PORTSMOUTH HOSPITALS NHS TRUST	9	0	28	34
<b>RHW</b>	ROYAL BERKSHIRE NHS FOUNDATION TRUST	#	0	29	#
<b>RJ1</b>	GUY'S AND ST THOMAS' NHS FOUNDATION TRUST	27	#	14	230
<b>RJ2</b>	LEWISHAM AND GREENWICH NHS TRUST	7	0	12	11
<b>RJ6</b>	CROYDON HEALTH SERVICES NHS TRUST	#	#	6	5
<b>RJ7</b>	ST GEORGE'S UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	#	#	#	159
<b>RJC</b>	SOUTH WARWICKSHIRE NHS FOUNDATION TRUST	9	5	16	#
<b>RJE</b>	UNIVERSITY HOSPITALS OF NORTH MIDLANDS NHS TRUST	5	0	25	34
<b>RJF</b>	BURTON HOSPITALS NHS FOUNDATION TRUST	0	#	25	0
<b>RJL</b>	NORTHERN LINCOLNSHIRE AND GOOLE NHS FOUNDATION TRUST	#	0	31	23
<b>RJN</b>	EAST CHESHIRE NHS TRUST	#	0	#	0

OFFICIAL

<b>RJR</b>	COUNTESS OF CHESTER HOSPITAL NHS FOUNDATION TRUST	#	0	5	131
<b>RJZ</b>	KING'S COLLEGE HOSPITAL NHS FOUNDATION TRUST	#	0	43	260
<b>RK5</b>	SHERWOOD FOREST HOSPITALS NHS FOUNDATION TRUST	#	#	18	145
<b>RK9</b>	UNIVERSITY HOSPITALS PLYMOUTH NHS TRUST	6	0	17	#
<b>RKB</b>	UNIVERSITY HOSPITALS COVENTRY AND WARWICKSHIRE NHS TRUST	20	6	28	125
<b>RKE</b>	WHITTINGTON HEALTH NHS TRUST	0	0	8	#
<b>RL1</b>	THE ROBERT JONES AND AGNES HUNT ORTHOPAEDIC HOSPITAL NHS FOUNDATION TRUST	0	0	11	9
<b>RL4</b>	THE ROYAL WOLVERHAMPTON NHS TRUST	#	#	6	73
<b>RLN</b>	CITY HOSPITALS SUNDERLAND NHS FOUNDATION TRUST	23	0	14	134
<b>RLQ</b>	WYE VALLEY NHS TRUST	#	0	9	0
<b>RLT</b>	GEORGE ELIOT HOSPITAL NHS TRUST	#	#	45	#
<b>RM1</b>	NORFOLK AND NORWICH UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	35	5	#	134
<b>RM2</b>	UNIVERSITY HOSPITAL OF SOUTH MANCHESTER NHS FOUNDATION TRUST	#	0	#	47
<b>RM3</b>	SALFORD ROYAL NHS FOUNDATION TRUST	#	0	18	142
<b>RMC</b>	BOLTON NHS FOUNDATION TRUST	7	#	9	5
<b>RMP</b>	TAMESIDE AND GLOSSOP INTEGRATED CARE NHS FOUNDATION TRUST	#	0	29	71
<b>RN3</b>	GREAT WESTERN HOSPITALS NHS FOUNDATION TRUST	#	#	50	5
<b>RN5</b>	HAMPSHIRE HOSPITALS NHS FOUNDATION TRUST	#	#	12	29
<b>RN7</b>	DARTFORD AND GRAVESHAM NHS TRUST	0	0	16	103
<b>RNA</b>	THE DUDLEY GROUP NHS FOUNDATION TRUST	#	#	45	#

OFFICIAL

<b>RNH</b>	NEWHAM UNIVERSITY HOSPITAL NHS TRUST	0	0	0	0
<b>RNL</b>	NORTH CUMBRIA UNIVERSITY HOSPITALS NHS TRUST	#	0	59	20
<b>RNQ</b>	KETTERING GENERAL HOSPITAL NHS FOUNDATION TRUST	#	#	32	110
<b>RNS</b>	NORTHAMPTON GENERAL HOSPITAL NHS TRUST	#	#	7	0
<b>RNZ</b>	SALISBURY NHS FOUNDATION TRUST	5	0	21	14
<b>RP4</b>	GREAT ORMOND STREET HOSPITAL FOR CHILDREN NHS FOUNDATION TRUST	0	0	0	0
<b>RP5</b>	DONCASTER AND BASSETLAW TEACHING HOSPITALS NHS FOUNDATION TRUST	7	#	117	678
<b>RP6</b>	MOORFIELDS EYE HOSPITAL NHS FOUNDATION TRUST	0	0	0	0
<b>RPA</b>	MEDWAY NHS FOUNDATION TRUST	9	#	29	37
<b>RPC</b>	QUEEN VICTORIA HOSPITAL NHS FOUNDATION TRUST	#	0	0	0
<b>RPY</b>	THE ROYAL MARSDEN NHS FOUNDATION TRUST	0	0	0	0
<b>RQ3</b>	BIRMINGHAM WOMEN'S AND CHILDREN'S NHS FOUNDATION TRUST	0	#	0	0
<b>RQ6</b>	ROYAL LIVERPOOL AND BROADGREEN UNIVERSITY HOSPITALS NHS TRUST	14	0	#	#
<b>RQ8</b>	MID ESSEX HOSPITAL SERVICES NHS TRUST	#	#	45	#
<b>RQM</b>	CHELSEA AND WESTMINSTER HOSPITAL NHS FOUNDATION TRUST	#	0	16	59
<b>RQW</b>	THE PRINCESS ALEXANDRA HOSPITAL NHS TRUST	#	#	#	#
<b>RQX</b>	HOMERTON UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	#	0	#	139
<b>RR1</b>	HEART OF ENGLAND NHS FOUNDATION TRUST	#	7	48	308
<b>RR7</b>	GATESHEAD HEALTH NHS FOUNDATION TRUST	0	#	53	15

OFFICIAL

<b>RR8</b>	LEEDS TEACHING HOSPITALS NHS TRUST	17	#	19	52
<b>RRF</b>	WRIGHTINGTON, WIGAN AND LEIGH NHS FOUNDATION TRUST	0	0	10	#
<b>RRJ</b>	THE ROYAL ORTHOPAEDIC HOSPITAL NHS FOUNDATION TRUST	0	0	#	33
<b>RRK</b>	UNIVERSITY HOSPITALS BIRMINGHAM NHS FOUNDATION TRUST	#	0	0	11
<b>RRV</b>	UNIVERSITY COLLEGE LONDON HOSPITALS NHS FOUNDATION TRUST	17	#	#	163
<b>RT3</b>	ROYAL BROMPTON & HAREFIELD NHS FOUNDATION TRUST	0	0	0	0
<b>RTD</b>	THE NEWCASTLE UPON TYNE HOSPITALS NHS FOUNDATION TRUST	19	#	8	269
<b>RTE</b>	GLOUCESTERSHIRE HOSPITALS NHS FOUNDATION TRUST	6	#	21	9
<b>RTF</b>	NORTHUMBRIA HEALTHCARE NHS FOUNDATION TRUST	0	#	15	15
<b>RTG</b>	UNIVERSITY HOSPITALS OF DERBY AND BURTON NHS FOUNDATION TRUST	6	#	14	44
<b>RTH</b>	OXFORD UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	#	0	#	#
<b>RTK</b>	ASHFORD AND ST PETER'S HOSPITALS NHS FOUNDATION TRUST	#	#	#	157
<b>RTP</b>	SURREY AND SUSSEX HEALTHCARE NHS TRUST	0	#	23	232
<b>RTR</b>	SOUTH TEES HOSPITALS NHS FOUNDATION TRUST	8	0	9	62
<b>RTX</b>	UNIVERSITY HOSPITALS OF MORECAMBE BAY NHS FOUNDATION TRUST	8	6	33	400
<b>RVJ</b>	NORTH BRISTOL NHS TRUST	0	#	15	8
<b>RVR</b>	EPSOM AND ST HELIER UNIVERSITY HOSPITALS NHS TRUST	#	#	18	383
<b>RVV</b>	EAST KENT HOSPITALS UNIVERSITY NHS FOUNDATION TRUST	7	#	37	454

OFFICIAL

<b>RWW</b>	NORTH TEES AND HARTLEPOOL NHS FOUNDATION TRUST	0	#	#	12
<b>RVY</b>	SOUTHPORT AND ORMSKIRK HOSPITAL NHS TRUST	#	0	5	#
<b>RW1</b>	SOUTHERN HEALTH NHS FOUNDATION TRUST	#	0	0	0
<b>RW3</b>	CENTRAL MANCHESTER UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	#	#	12	11
<b>RW6</b>	PENNINE ACUTE HOSPITALS NHS TRUST	5	#	23	177
<b>RWA</b>	HULL AND EAST YORKSHIRE HOSPITALS NHS TRUST	27	#	14	25
<b>RWD</b>	UNITED LINCOLNSHIRE HOSPITALS NHS TRUST	6	#	40	220
<b>RWE</b>	UNIVERSITY HOSPITALS OF LEICESTER NHS TRUST	#	0	28	184
<b>RWF</b>	MAIDSTONE AND TUNBRIDGE WELLS NHS TRUST	5	0	8	72
<b>RWG</b>	WEST HERTFORDSHIRE HOSPITALS NHS TRUST	10	#	14	33
<b>RWH</b>	EAST AND NORTH HERTFORDSHIRE NHS TRUST	#	#	5	#
<b>RWJ</b>	STOCKPORT NHS FOUNDATION TRUST	#	0	#	7
<b>RWP</b>	WORCESTERSHIRE ACUTE HOSPITALS NHS TRUST	0	6	12	92
<b>RWW</b>	WARRINGTON AND HALTON HOSPITALS NHS FOUNDATION TRUST	#	8	14	12
<b>RWY</b>	CALDERDALE AND HUDDERSFIELD NHS FOUNDATION TRUST	6	#	22	86
<b>RX1</b>	NOTTINGHAM UNIVERSITY HOSPITALS NHS TRUST	#	0	15	385
<b>RXC</b>	EAST SUSSEX HEALTHCARE NHS TRUST	#	#	6	#
<b>RXF</b>	MID YORKSHIRE HOSPITALS NHS TRUST	0	#	33	8
<b>RXH</b>	BRIGHTON AND SUSSEX UNIVERSITY HOSPITALS NHS TRUST	7	5	7	99
<b>RXK</b>	SANDWELL AND WEST BIRMINGHAM HOSPITALS NHS TRUST	#	0	19	65

OFFICIAL

<b>RXL</b>	BLACKPOOL TEACHING HOSPITALS NHS FOUNDATION TRUST	#	0	17	255
<b>RXN</b>	LANCASHIRE TEACHING HOSPITALS NHS FOUNDATION TRUST	35	0	#	632
<b>RXP</b>	COUNTY DURHAM AND DARLINGTON NHS FOUNDATION TRUST	5	#	#	128
<b>RXQ</b>	BUCKINGHAMSHIRE HEALTHCARE NHS TRUST	#	0	8	22
<b>RXR</b>	EAST LANCASHIRE HOSPITALS NHS TRUST	14	#	23	#
<b>RXW</b>	SHREWSBURY AND TELFORD HOSPITAL NHS TRUST	#	#	9	19
<b>RY8</b>	DERBYSHIRE COMMUNITY HEALTH SERVICES NHS FOUNDATION TRUST	0	0	#	0
<b>RYJ</b>	IMPERIAL COLLEGE HEALTHCARE NHS TRUST	8	18	30	54
<b>RYR</b>	WESTERN SUSSEX HOSPITALS NHS FOUNDATION TRUST	9	#	27	143
<b>RYV</b>	CAMBRIDGESHIRE COMMUNITY SERVICES NHS TRUST	0	0	0	0
<b>Total</b>		819	237	3,455	13,180

Table 9<sup>29</sup>: 2017/18 provider activity baseline estimate for Category 2 interventions (E-Q) and total Category 2 activity <sup>30</sup>

		E Breast reduction	F Removal of benign skin	G Grommets	H Tonsillectomy	I Haemorrhoid	J Hysterectomy for	K Chalazia removal	L Shoulder decompression	M Carpal tunnel syndrome	N Dupuytren's	O Ganglion excision	P Trigger finger release	Q Varicose vein
		No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells	No of spells
Provider code	Provider name	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity	2017/18 activity
8G301	RYALLS ANN (COUNSELLOR)	0	0	0	0	0	0	0	0	0	0	0	0	0
8G326	THE SPENCER WING	0	21	0	#	#	31	0	8	0	#	0	0	0
AAH01	TETBURY HOSPITAL TRUST	0	32	0	0	0	0	0	0	18	0	0	0	0
ADP02	KIMS HOSPITAL (NEWNHAM COURT)	0	109	0	8	19	50	0	68	132	39	18	43	0
AHH01	FOSCOTE COURT (BANBURY) TRUST	0	0	0	0	0	#	0	0	18	#	#	0	0
AJX01	SUSSEX MSK PARTNERSHIP 2 LTD	0	27	0	0	0	0	0	48	169	31	30	20	0
AVQ01	ONE ASHFORD HOSPITAL	0	#	0	0	#	#	0	0	#	17	9	#	0
RXG00	WILTSHIRE HEATH AND CARE LLP	0	#	0	0	0	0	0	0	0	0	0	0	0
NAM01	PROBUS SURGICAL CENTRE	0	#	0	0	0	0	0	0	307	#	0	#	0
NAM02	MENEAGE STREET SURGERY	0	0	0	0	0	0	0	0	#	0	0	0	0

<sup>29</sup> Figures based on 'all non-emergency spells' which includes day cases and inpatient activity and also non-emergency non-elective admissions. Due to limitations in what can be reliably measured nationally, outpatient activity is not included.

<sup>30</sup> Provider activity figures include activity that could not be attributed to a CCG.

OFFICIAL

<b>NAM04</b>	LISKEARD COMMUNITY HOSPITAL	0	0	0	0	0	0	0	0	#	0	0	0	0
<b>NAM06</b>	MORRAB SURGERY	0	0	0	0	0	0	0	0	#	0	0	0	0
<b>NEQ01</b>	PHOENIX HEALTH SOLUTIONS LIMITED	0	#	0	0	0	0	11	0	20	0	#	0	10
<b>NFH01</b>	SOMERSET SURGICAL SERVICES	0	108	0	0	0	0	0	0	34	22	5	12	0
<b>NHW03</b>	WIMBOURNE COMMUNITY HOSPITAL (STANDARD HEALTH)	0	0	0	0	0	0	0	0	19	7	0	0	0
<b>NN401</b>	TYNESIDE SURGICAL SERVICES	5	36	0	0	18	14	0	14	8	8	#	6	8
<b>NN501</b>	TRAFFORD GENERAL HOSPITAL	0	24	0	26	#	0	0	11	72	28	30	10	0
<b>NN801</b>	THE SPENCER WING (RAMSGATE ROAD)	0	36	0	#	7	53	0	15	#	5	#	#	0
<b>NN802</b>	THE SPENCER WING (WILLIAM HARVEY HOSPITAL)	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>NNE02</b>	DORKING GENERAL HOSPITAL	0	5	0	#	8	5	0	21	38	6	#	#	27
<b>NQM01</b>	ORTHOPAEDICS & SPINE SPECIALIST HOSPITAL SITE	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>NT202</b>	NUFFIELD HEALTH, BOURNEMOUTH HOSPITAL	0	46	0	0	0	26	0	0	39	9	5	#	0
<b>NT204</b>	NUFFIELD HEALTH, BRENTWOOD HOSPITAL	0	#	#	32	0	49	0	7	12	#	0	0	0
<b>NT206</b>	NUFFIELD HEALTH, BRISTOL HOSPITAL (CHESTERFIELD)	0	0	0	0	6	0	0	#	21	#	#	#	0

OFFICIAL

<b>NT209</b>	NUFFIELD HEALTH, CAMBRIDGE HOSPITAL	0	0	0	0	0	0	0	#	12	9	#	#	0
<b>NT210</b>	NUFFIELD HEALTH, THE GROSVENOR HOSPITAL, CHESTER	0	0	0	0	0	0	0	6	0	0	0	0	0
<b>NT211</b>	NUFFIELD HEALTH, CHELTENHAM HOSPITAL	0	#	0	0	0	0	0	5	0	0	0	0	0
<b>NT212</b>	NUFFIELD HEALTH, CHICHESTER HOSPITAL	0	10	0	0	0	25	0	39	54	22	#	7	0
<b>NT213</b>	NUFFIELD HEALTH, DERBY HOSPITAL	0	#	0	25	0	0	0	11	#	0	0	0	0
<b>NT214</b>	NUFFIELD HEALTH, WESSEX HOSPITAL	0	#	0	0	0	5	0	48	74	32	12	12	0
<b>NT215</b>	NUFFIELD HEALTH, EXETER HOSPITAL	0	0	0	0	6	0	0	5	0	0	0	0	0
<b>NT218</b>	NUFFIELD HEALTH, HAYWARDS HEATH HOSPITAL	0	0	0	0	0	0	0	13	#	#	0	0	0
<b>NT219</b>	NUFFIELD HEALTH, HEREFORD HOSPITAL	0	5	0	#	13	#	0	#	#	11	#	#	0
<b>NT224</b>	NUFFIELD HEALTH, WARWICKSHIRE HOSPITAL	0	#	0	0	0	0	0	16	32	12	8	12	0
<b>NT225</b>	NUFFIELD HEALTH, LEEDS HOSPITAL	0	9	0	114	78	79	0	69	143	47	20	44	0
<b>NT226</b>	NUFFIELD HEALTH, LEICESTER HOSPITAL	0	#	0	134	0	5	0	0	33	6	#	6	0
<b>NT229</b>	NUFFIELD HEALTH, NEWCASTLE UPON TYNE HOSPITAL	0	0	0	0	0	0	0	#	0	0	0	0	0
<b>NT230</b>	NUFFIELD HEALTH, NORTH STAFFORDSHIRE HOSPITAL	0	8	0	0	0	0	0	124	129	70	21	34	0
<b>NT233</b>	NUFFIELD HEALTH, PLYMOUTH HOSPITAL	0	19	0	0	5	0	0	22	#	37	7	12	0

OFFICIAL

<b>NT235</b>	NUFFIELD HEALTH, SHREWSBURY HOSPITAL	0	10	0	0	0	0	0	44	135	32	25	14	0
<b>NT237</b>	NUFFIELD HEALTH, TEES HOSPITAL	0	6	0	0	0	0	0	18	62	31	12	19	0
<b>NT238</b>	NUFFIELD HEALTH, TAUNTON HOSPITAL	0	#	0	31	0	9	0	20	123	43	0	30	0
<b>NT239</b>	NUFFIELD HEALTH, TUNBRIDGE WELLS HOSPITAL	0	#	0	12	#	0	0	9	10	#	5	#	0
<b>NT241</b>	NUFFIELD HEALTH, WOKING HOSPITAL	0	#	0	0	0	0	0	9	24	5	#	5	0
<b>NT242</b>	NUFFIELD HEALTH, WOLVERHAMPTON HOSPITAL	0	#	0	0	0	0	0	25	65	18	#	20	#
<b>NT245</b>	NUFFIELD HEALTH, YORK HOSPITAL	0	#	0	0	0	0	0	32	16	21	0	#	0
<b>NT301</b>	SPIRE SOUTH BANK HOSPITAL	0	12	0	15	20	34	0	22	186	43	42	28	0
<b>NT302</b>	SPIRE BRISTOL HOSPITAL	0	0	0	0	0	0	0	139	175	100	9	45	0
<b>NT304</b>	SPIRE SOUTHAMPTON HOSPITAL	0	6	0	0	0	5	0	51	47	7	10	10	0
<b>NT305</b>	SPIRE PORTSMOUTH HOSPITAL	0	#	0	0	0	0	0	36	34	6	#	5	0
<b>NT308</b>	SPIRE GATWICK PARK HOSPITAL	0	#	0	0	0	#	0	9	8	#	#	0	12
<b>NT309</b>	SPIRE SUSSEX HOSPITAL	0	14	0	5	#	28	0	22	65	17	20	17	0
<b>NT310</b>	SPIRE TUNBRIDGE WELLS HOSPITAL	0	12	0	8	#	14	0	15	20	15	11	5	0
<b>NT312</b>	SPIRE ALEXANDRA HOSPITAL	0	137	0	0	12	42	0	15	17	6	5	9	0
<b>NT313</b>	SPIRE WELLESLEY HOSPITAL	0	25	0	0	0	0	0	58	108	52	36	11	0
<b>NT314</b>	SPIRE LONDON EAST	0	134	0	27	19	25	0	20	26	9	12	#	0
<b>NT315</b>	SPIRE BUSHEY HOSPITAL	0	41	0	0	8	0	0	23	17	7	5	5	0

OFFICIAL

<b>NT316</b>	SPIRE HARPENDEN HOSPITAL	0	34	0	27	0	0	0	13	34	13	7	10	0
<b>NT317</b>	SPIRE CAMBRIDGE LEA HOSPITAL	0	5	0	0	0	0	0	21	0	0	0	#	0
<b>NT318</b>	SPIRE NORWICH HOSPITAL	0	0	0	0	0	0	0	32	0	0	0	0	0
<b>NT319</b>	SPIRE HARTSWOOD HOSPITAL	0	6	0	19	0	0	0	13	37	16	10	9	0
<b>NT320</b>	SPIRE PARKWAY HOSPITAL	0	5	0	0	18	53	0	77	99	33	16	21	0
<b>NT321</b>	SPIRE LITTLE ASTON HOSPITAL	0	26	0	0	0	56	0	58	51	26	9	#	0
<b>NT322</b>	SPIRE LEICESTER HOSPITAL	0	22	0	0	0	0	0	51	52	16	16	15	0
<b>NT324</b>	SPIRE CHESHIRE HOSPITAL	#	127	0	32	26	12	0	18	63	43	11	14	83
<b>NT325</b>	SPIRE MURRAYFIELD HOSPITAL	0	22	0	16	5	#	0	48	108	23	6	17	94
<b>NT327</b>	SPIRE MANCHESTER HOSPITAL	0	27	0	#	#	0	0	11	37	17	13	11	64
<b>NT332</b>	SPIRE LEEDS HOSPITAL	0	154	9	123	63	0	0	85	230	55	59	42	0
<b>NT333</b>	SPIRE WASHINGTON HOSPITAL	0	32	0	#	18	18	0	14	62	27	21	15	229
<b>NT337</b>	SPIRE LIVERPOOL HOSPITAL	0	14	0	61	41	16	0	71	200	40	16	37	0
<b>NT338</b>	SPIRE YALE HOSPITAL	0	0	0	0	0	0	0	0	#	0	#	0	0
<b>NT339</b>	SPIRE REGENCY HOSPITAL	0	10	0	18	19	32	0	35	66	38	31	13	0
<b>NT343</b>	SPIRE THAMES VALLEY HOSPITAL	0	#	0	#	8	17	0	8	#	#	0	0	0
<b>NT344</b>	SPIRE DUNEDIN HOSPITAL	0	#	0	0	5	0	0	32	28	8	6	13	0
<b>NT345</b>	SPIRE CLARE PARK HOSPITAL	0	#	0	5	#	0	0	12	19	5	#	#	0
<b>NT347</b>	SPIRE FYLDE COAST HOSPITAL	0	25	0	46	5	22	0	18	38	26	6	11	117

OFFICIAL

<b>NT348</b>	SPIRE ELLAND HOSPITAL	0	6	0	0	35	57	0	62	98	31	17	19	34
<b>NT350</b>	SPIRE METHLEY PARK HOSPITAL	0	19	0	28	#	32	0	46	72	13	9	13	100
<b>NT351</b>	SPIRE HULL AND EAST RIDING HOSPITAL	0	356	0	14	#	48	0	121	70	35	10	66	0
<b>NT364</b>	SPIRE MONTEFIORE HOSPITAL	0	#	0	0	46	7	0	7	21	#	#	#	0
<b>NT401</b>	BMI - THE ALEXANDRA HOSPITAL	0	79	0	0	8	26	0	69	99	39	15	25	#
<b>NT402</b>	BMI - BATH CLINIC	0	7	0	#	13	22	0	12	#	7	0	0	0
<b>NT403</b>	BMI - THE BEARDWOOD HOSPITAL	0	7	0	9	43	7	0	17	60	20	10	21	0
<b>NT404</b>	BMI - THE BEAUMONT HOSPITAL	0	16	0	38	25	38	0	39	62	11	#	27	0
<b>NT405</b>	BMI - BISHOPS WOOD	0	39	0	12	10	19	0	#	20	16	11	#	0
<b>NT406</b>	BMI - THE BLACKHEATH HOSPITAL	0	29	0	8	6	26	0	#	7	#	#	#	0
<b>NT408</b>	BMI - THE CHAUCER HOSPITAL	0	55	0	12	38	33	0	5	24	52	35	11	0
<b>NT409</b>	BMI - CHELSFIELD PARK HOSPITAL	0	15	0	21	#	27	0	9	16	#	#	#	0
<b>NT410</b>	BMI - THE CHILTERN HOSPITAL	0	0	0	#	5	11	0	7	0	0	0	0	0
<b>NT411</b>	BMI - THE CLEMENTINE CHURCHILL HOSPITAL	0	33	0	14	35	18	0	13	56	7	10	6	0
<b>NT412</b>	BMI - THE DROITWICH SPA HOSPITAL	0	18	0	11	#	43	#	39	130	31	22	10	33
<b>NT413</b>	BMI - THE ESPERANCE HOSPITAL	0	28	0	13	10	19	0	6	#	#	0	#	0
<b>NT414</b>	BMI - FAWKHAM MANOR HOSPITAL	0	32	0	#	18	16	0	8	10	5	#	#	0

OFFICIAL

<b>NT416</b>	BMI - HENDON HOSPITAL	0	10	0	0	#	22	0	#	16	#	6	5	19
<b>NT417</b>	BMI - GORING HALL HOSPITAL	0	#	0	0	0	58	0	8	30	14	5	12	0
<b>NT418</b>	BMI - THE HAMPSHIRE CLINIC	0	23	0	11	7	7	0	19	17	5	#	#	0
<b>NT419</b>	BMI - THE HARBOUR HOSPITAL	0	0	0	#	#	0	0	#	81	0	0	0	0
<b>NT420</b>	BMI - THE HIGHFIELD HOSPITAL	0	38	0	33	16	56	0	71	62	70	42	48	0
<b>NT421</b>	BMI - THE KINGS OAK HOSPITAL	0	21	0	0	#	11	0	40	53	11	5	7	0
<b>NT422</b>	BMI - THE LONDON INDEPENDENT HOSPITAL	0	156	0	10	67	#	0	14	60	7	9	8	#
<b>NT423</b>	BMI - THE MANOR HOSPITAL	0	#	0	0	0	7	0	#	#	0	0	#	0
<b>NT424</b>	BMI - THE MERIDEN HOSPITAL	0	6	0	18	#	37	0	52	68	7	10	6	0
<b>NT427</b>	BMI - THE PARK HOSPITAL	0	62	0	#	9	86	0	34	0	0	0	0	0
<b>NT428</b>	BMI - THE PRINCESS MARGARET HOSPITAL	0	#	0	0	8	14	0	6	5	#	0	#	0
<b>NT429</b>	BMI - THE PRIORY HOSPITAL	0	#	0	0	0	32	0	21	43	#	8	6	0
<b>NT430</b>	BMI - THE RIDGEWAY HOSPITAL	0	#	0	12	38	15	0	14	10	8	#	#	0
<b>NT431</b>	BMI - THE RUNNYMEDE HOSPITAL	0	24	0	0	9	0	0	0	#	0	0	#	0
<b>NT432</b>	BMI - THE SANDRINGHAM HOSPITAL	0	#	0	0	14	0	0	0	0	0	0	0	0
<b>NT433</b>	BMI - SARUM ROAD HOSPITAL	0	5	0	#	0	6	0	9	38	24	9	8	0
<b>NT434</b>	BMI - THE SAXON CLINIC	0	18	0	0	#	5	0	26	46	7	5	11	0

OFFICIAL

<b>NT435</b>	BMI - THE SHELBURNE HOSPITAL	0	6	0	0	0	0	0	14	0	0	0	0	0
<b>NT436</b>	BMI - SHIRLEY OAKS HOSPITAL	0	8	0	7	13	41	0	19	89	16	22	13	15
<b>NT437</b>	BMI - THE SLOANE HOSPITAL	0	#	0	25	8	9	0	#	10	#	0	#	0
<b>NT438</b>	BMI - THE SOMERFIELD HOSPITAL	0	32	0	7	#	40	0	5	14	8	5	9	0
<b>NT439</b>	BMI - THE SOUTH CHESHIRE PRIVATE HOSPITAL	0	16	0	22	6	27	0	41	80	21	6	20	0
<b>NT440</b>	BMI - THORNBURY HOSPITAL	0	#	0	0	33	71	0	10	21	10	7	#	0
<b>NT441</b>	BMI - THREE SHIRES HOSPITAL	0	45	0	0	35	37	0	42	158	35	26	21	0
<b>NT443</b>	BMI - THE WINTERBOURNE HOSPITAL	0	7	0	0	0	0	0	113	43	18	#	11	0
<b>NT445</b>	BMI THE EDGBASTON HOSPITAL	0	12	0	0	8	43	0	23	62	14	22	10	0
<b>NT446</b>	BMI ST EDMUNDS HOSPITAL	0	0	0	0	#	11	0	0	21	0	0	0	0
<b>NT447</b>	BMI THE DUCHY HOSPITAL	0	#	0	7	0	0	0	9	14	#	#	#	24
<b>NT448</b>	BMI THE HUDDERSFIELD HOSPITAL	0	8	0	51	8	12	0	40	47	6	0	9	0
<b>NT449</b>	BMI THE LANCASTER HOSPITAL	0	30	0	0	#	#	0	17	18	21	12	9	0
<b>NT450</b>	BMI THE LINCOLN HOSPITAL	0	#	0	0	0	0	0	7	0	0	0	0	0
<b>NT451</b>	BMI THE CAVELL HOSPITAL	0	36	0	13	0	47	0	17	11	5	0	#	0
<b>NT455</b>	BMI MOUNT ALVERNIA HOSPITAL	0	8	0	15	0	13	0	0	#	6	#	#	0
<b>NT457</b>	BMI WOODLANDS HOSPITAL	0	24	0	6	32	52	0	#	106	53	26	18	74

OFFICIAL

<b>NT490</b>	BMI SOUTHEND PRIVATE HOSPITAL	0	70	0	0	0	0	#	0	56	0	#	#	0
<b>NT497</b>	BMI GISBURNE PARK HOSPITAL	0	14	0	38	#	10	#	34	57	22	7	15	0
<b>NTE02</b>	ST HUGH'S HOSPITAL	0	14	0	#	#	20	0	56	87	13	#	#	0
<b>NTP11</b>	SOUTHAMPTON NHS TREATMENT CENTRE	0	213	0	96	14	45	0	#	165	10	47	25	8
<b>NTP13</b>	BARLBOROUGH NHS TREATMENT CENTRE	0	#	0	0	0	0	0	122	0	18	26	26	0
<b>NTP15</b>	NORTH EAST LONDON TREATMENT CENTRE CARE UK	0	298	0	0	78	0	25	7	190	45	82	43	26
<b>NTP16</b>	WILL ADAMS NHS TREATMENT CENTRE	0	180	0	0	0	0	18	0	52	11	26	9	45
<b>NTPAD</b>	ST MARY'S NHS TREATMENT CENTRE	0	123	0	0	5	0	0	0	240	44	17	46	0
<b>NTPAE</b>	THE CROFT SHIFA HEALTH CENTRE	0	0	0	0	0	0	12	0	0	0	0	0	0
<b>NTPH1</b>	SHEPTON MALLET NHS TREATMENT CENTRE	0	22	0	5	#	5	#	18	277	49	5	54	0
<b>NTPH2</b>	EMERSONS GREEN NHS TREATMENT CENTRE	0	59	0	45	#	57	36	0	251	98	31	29	79
<b>NTPH3</b>	DEVIZES NHS TREATMENT CENTRE	0	15	0	10	#	0	8	0	#	42	7	13	11
<b>NTPH5</b>	PENINSULA NHS TREATMENT CENTRE	0	8	0	0	0	0	0	69	202	59	30	40	0
<b>NTX01</b>	ONE HEALTH GROUP LTD	0	40	0	0	#	0	0	#	5	0	0	0	0
<b>NTX06</b>	ONE HEALTH GROUP CLINIC - OSSET	0	0	0	0	0	0	0	0	0	0	0	0	0

OFFICIAL

<b>NTX09</b>	ONE HEALTH GROUP CLINIC - GAINSBOROUGH	0	0	0	0	0	0	0	0	#	0	0	0	0
<b>NTX11</b>	ONE HEALTH GROUP CLINIC - THORNBURY	0	79	0	0	8	18	0	48	115	33	15	21	0
<b>NTX12</b>	ONE HEALTH GROUP CLINIC - CLAREMONT	0	33	0	0	#	9	0	16	22	8	#	#	0
<b>NV302</b>	CIRCLE BATH HOSPITAL	0	40	0	18	16	80	0	165	90	73	11	26	120
<b>NV313</b>	CIRCLE - NOTTINGHAM NHS TREATMENT CENTRE	0	1,444	0	0	58	99	#	116	656	131	64	110	117
<b>NV323</b>	CIRCLE READING HOSPITAL	0	14	0	16	22	24	0	67	66	19	5	20	0
<b>NVC01</b>	ASHTEAD HOSPITAL	0	33	0	6	15	7	0	15	40	10	10	#	24
<b>NVC02</b>	THE BERKSHIRE INDEPENDENT HOSPITAL	0	5	0	0	#	#	0	34	66	15	8	10	0
<b>NVC04</b>	DUCHY HOSPITAL	0	41	0	0	27	0	0	17	119	123	27	43	104
<b>NVC05</b>	EUXTON HALL HOSPITAL	0	175	0	15	35	40	0	64	95	68	24	11	0
<b>NVC06</b>	FITZWILLIAM HOSPITAL	0	9	0	29	13	55	0	99	47	14	8	#	24
<b>NVC07</b>	FULWOOD HALL HOSPITAL	0	92	0	15	27	36	0	72	122	61	39	20	131
<b>NVC08</b>	MOUNT STUART HOSPITAL	0	84	0	11	21	68	0	39	99	59	5	21	0
<b>NVC09</b>	NEW HALL HOSPITAL	0	13	0	13	5	33	0	165	49	41	11	18	0
<b>NVC0M</b>	RAMSAY CROYDON DAY HOSPITAL	0	0	0	0	0	0	0	0	#	0	0	0	0
<b>NVC0R</b>	TEES VALLEY HOSPITAL	0	63	0	0	#	0	0	0	#	#	#	#	0
<b>NVC11</b>	NORTH DOWNS HOSPITAL	0	34	0	0	11	15	0	22	45	#	6	#	17
<b>NVC12</b>	OAKLANDS HOSPITAL	0	46	0	30	#	22	0	48	155	44	48	51	0
<b>NVC13</b>	OAKS HOSPITAL	0	79	0	9	16	25	0	28	14	26	14	7	126

OFFICIAL

<b>NVC14</b>	PARK HILL HOSPITAL	0	7	0	#	0	0	0	#	21	9	9	6	0
<b>NVC15</b>	PINEHILL HOSPITAL	0	16	0	22	37	32	0	#	61	9	7	17	0
<b>NVC16</b>	RENACRES HOSPITAL	0	37	0	12	13	16	5	40	52	26	#	9	0
<b>NVC17</b>	ROWLEY HALL HOSPITAL	0	15	0	0	16	46	0	60	44	29	0	18	0
<b>NVC18</b>	SPRINGFIELD HOSPITAL	0	17	0	78	49	74	15	23	66	15	10	11	0
<b>NVC19</b>	RIVERS HOSPITAL	0	7	0	25	99	65	0	0	111	14	#	16	#
<b>NVC20</b>	THE YORKSHIRE CLINIC	0	262	0	69	55	21	#	117	333	154	85	60	0
<b>NVC21</b>	WEST MIDLANDS HOSPITAL	0	8	0	14	15	102	0	16	104	26	7	15	0
<b>NVC22</b>	WINFIELD HOSPITAL	0	8	0	6	8	8	0	8	98	20	5	15	0
<b>NVC23</b>	WOODLAND HOSPITAL	0	41	0	8	27	55	9	16	70	52	36	25	0
<b>NVC25</b>	HORTON NHS TREATMENT CENTRE	0	13	0	0	0	0	0	154	325	58	16	46	0
<b>NVC27</b>	BOSTON WEST HOSPITAL	0	6	0	0	0	0	0	97	#	18	#	0	30
<b>NVC28</b>	CLIFTON PARK HOSPITAL	0	#	0	0	0	0	0	58	69	43	#	11	0
<b>NVC29</b>	COBALT HOSPITAL	0	87	0	0	9	0	0	#	35	19	27	13	23
<b>NVC31</b>	BLAKELANDS HOSPITAL	0	61	0	0	21	0	14	70	125	21	20	27	0
<b>NVC35</b>	TEES VALLEY TREATMENT CENTRE	0	664	0	0	25	0	0	6	104	41	27	19	9
<b>NVC40</b>	WOODTHORPE HOSPITAL	0	16	0	0	#	25	0	36	105	47	12	28	0
<b>NVC44</b>	THE WESTBOURNE CENTRE	0	72	0	0	0	0	39	0	52	0	#	5	48
<b>NVG01</b>	FAIRFIELD HOSPITAL	0	166	0	23	18	53	#	9	63	13	5	5	162
<b>NVM01</b>	COBHAM DAY SURGERY HOSPITAL	0	394	0	0	0	0	0	0	27	18	10	6	8

OFFICIAL

<b>NVM02</b>	EPSOM DAY SURGERY LIMITED	0	243	0	0	8	0	0	0	38	11	#	9	25
<b>NW612</b>	HCA - 52 ALDERLEY ROAD	0	12	0	#	0	0	7	0	#	7	#	6	0
<b>NWF01</b>	BENENDEN HOSPITAL	0	19	0	15	9	21	0	29	23	17	9	6	0
<b>NWX11</b>	BICS-COMMUNITY	0	13	0	0	0	0	0	32	153	27	55	40	0
<b>NXM01</b>	THE HORDER CENTRE - ST JOHNS ROAD	0	13	0	0	0	0	0	145	37	12	11	8	0
<b>NXM04</b>	THE MCINDOE CENTRE	0	12	0	0	0	0	0	24	49	36	15	20	0
<b>NXP04</b>	HATHAWAY MEDICAL CENTRE	0	#	0	0	0	0	0	0	8	0	0	0	0
<b>NXP17</b>	WHITE HORSE HEALTH CENTRE - IHG	0	#	0	0	0	0	0	0	#	0	0	0	0
<b>NXP20</b>	LAWN MEDICAL CENTRE	0	0	0	0	0	0	0	0	5	0	0	0	0
<b>NXP37</b>	ASPEN CENTRE	0	0	0	0	0	0	0	0	25	0	0	0	0
<b>NXP40</b>	LITFIELD HOUSE MEDICAL CENTRE	0	0	0	0	0	0	0	0	9	0	0	0	0
<b>NXP41</b>	IHG CRAVEN ROAD	0	0	0	0	0	0	0	0	13	0	0	0	0
<b>NYW01</b>	ASPEN - THE HOLLY	0	94	0	44	#	26	0	23	86	25	21	17	0
<b>NYW02</b>	ASPEN - PARKSIDE HOSPITAL	0	7	0	0	0	#	0	12	17	#	8	7	0
<b>NYW03</b>	ASPEN - HIGHGATE HOSPITAL	0	67	0	11	#	0	0	7	15	5	#	7	0
<b>NYW04</b>	ASPEN - CLAREMONT HOSPITAL	0	8	0	0	12	26	0	17	33	#	#	#	0
<b>PHS</b>	PHOENIX HEALTH SOLUTIONS LTD	0	44	0	0	0	0	7	0	35	0	#	0	22
<b>R0A</b>	MANCHESTER UNIVERSITY NHS FOUNDATION TRUST	29	904	82	344	73	160	91	56	192	75	30	28	267
<b>R1D</b>	SHROPSHIRE COMMUNITY HEALTH NHS TRUST	0	25	0	0	0	0	0	0	49	5	9	#	0

OFFICIAL

<b>R1F</b>	ISLE OF WIGHT NHS TRUST	0	123	7	64	35	63	5	19	127	30	8	14	0
<b>R1H</b>	BARTS HEALTH NHS TRUST	20	796	127	705	139	239	49	48	303	42	60	50	726
<b>R1K</b>	LONDON NORTH WEST UNIVERSITY HEALTHCARE NHS TRUST	#	713	57	267	121	113	8	17	153	25	27	25	760
<b>RA2</b>	ROYAL SURREY COUNTY HOSPITAL NHS FOUNDATION TRUST	5	547	32	118	50	194	#	55	252	94	23	53	0
<b>RA3</b>	WESTON AREA HEALTH NHS TRUST	0	29	0	0	17	16	0	49	94	61	5	25	0
<b>RA4</b>	YEOVIL DISTRICT HOSPITAL NHS FOUNDATION TRUST	0	343	13	0	19	96	9	40	139	32	#	37	0
<b>RA7</b>	UNIVERSITY HOSPITALS BRISTOL NHS FOUNDATION TRUST	0	1,395	106	202	35	95	43	0	86	#	0	9	0
<b>RA9</b>	TORBAY AND SOUTH DEVON NHS FOUNDATION TRUST	#	188	65	100	12	134	0	20	172	56	#	17	105
<b>RAE</b>	BRADFORD TEACHING HOSPITALS NHS FOUNDATION TRUST	34	552	109	216	25	162	43	25	214	37	9	21	304
<b>RAJ</b>	SOUTHEND UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	0	408	49	272	57	181	111	74	151	38	20	17	328
<b>RAL</b>	ROYAL FREE LONDON NHS FOUNDATION TRUST	50	2,490	45	424	84	141	25	35	282	123	33	62	568
<b>RAN</b>	ROYAL NATIONAL ORTHOPAEDIC	0	114	0	0	0	0	0	13	11	0	#	0	0

OFFICIAL

	HOSPITAL NHS TRUST													
<b>RAP</b>	NORTH MIDDLESEX UNIVERSITY HOSPITAL NHS TRUST	#	126	0	0	23	71	25	40	137	11	17	22	0
<b>RAS</b>	THE HILLINGDON HOSPITALS NHS FOUNDATION TRUST	0	510	0	0	73	135	9	57	193	72	43	28	191
<b>RAX</b>	KINGSTON HOSPITAL NHS FOUNDATION TRUST	13	392	38	201	92	109	12	185	218	85	24	33	119
<b>RBA</b>	TAUNTON AND SOMERSET NHS FOUNDATION TRUST	#	348	75	133	39	172	5	32	179	57	13	38	196
<b>RBD</b>	DORSET COUNTY HOSPITAL NHS FOUNDATION TRUST	8	337	20	92	22	142	10	136	251	94	14	32	69
<b>RBK</b>	WALSALL HEALTHCARE NHS TRUST	10	443	21	168	57	127	0	125	103	22	18	16	133
<b>RBL</b>	WIRRAL UNIVERSITY TEACHING HOSPITAL NHS FOUNDATION TRUST	27	1,563	75	180	21	110	#	82	427	105	15	23	261
<b>RBN</b>	ST HELENS AND KNOWSLEY HOSPITAL SERVICES NHS TRUST	57	2,478	46	223	59	253	86	31	292	104	36	66	121
<b>RBQ</b>	LIVERPOOL HEART AND CHEST HOSPITAL NHS FOUNDATION TRUST	0	#	0	0	0	0	0	0	0	0	0	0	0
<b>RBS</b>	ALDER HEY CHILDREN'S NHS	0	471	128	303	0	0	20	0	0	0	6	0	0

OFFICIAL

	FOUNDATION TRUST													
<b>RBT</b>	MID CHESHIRE HOSPITALS NHS FOUNDATION TRUST	#	474	28	169	22	93	78	116	252	85	6	30	87
<b>RBV</b>	THE CHRISTIE NHS FOUNDATION TRUST	#	279	0	0	#	52	0	0	0	0	0	0	0
<b>RBZ</b>	NORTHERN DEVON HEALTHCARE NHS TRUST	#	456	5	0	#	73	7	40	121	39	18	40	109
<b>RC1</b>	BEDFORD HOSPITAL NHS TRUST	#	1,027	50	70	30	113	5	15	131	36	5	15	114
<b>RC9</b>	LUTON AND DUNSTABLE UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	5	310	44	336	52	100	20	24	167	45	16	21	102
<b>RCB</b>	YORK TEACHING HOSPITAL NHS FOUNDATION TRUST	29	426	67	281	48	193	20	166	403	113	34	45	545
<b>RCD</b>	HARROGATE AND DISTRICT NHS FOUNDATION TRUST	15	182	50	111	41	101	27	65	196	78	16	32	0
<b>RCF</b>	AIREDALE NHS FOUNDATION TRUST	#	88	48	0	19	118	28	32	127	37	16	9	130
<b>RCU</b>	SHEFFIELD CHILDREN'S NHS FOUNDATION TRUST	0	275	96	114	#	0	19	0	#	0	11	0	0
<b>RCX</b>	THE QUEEN ELIZABETH HOSPITAL, KING'S LYNN, NHS FOUNDATION TRUST	#	775	35	166	28	107	58	30	131	53	8	20	135

OFFICIAL

<b>RD1</b>	ROYAL UNITED HOSPITALS BATH NHS FOUNDATION TRUST	19	104	39	111	19	90	#	57	84	58	6	21	#
<b>RD3</b>	POOLE HOSPITAL NHS FOUNDATION TRUST	#	1,439	22	150	10	171	0	0	#	0	#	#	0
<b>RD8</b>	MILTON KEYNES UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	17	371	81	193	16	120	35	59	150	44	9	9	43
<b>RDD</b>	BASILDON AND THURROCK UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	#	130	60	272	51	146	0	62	128	43	#	6	108
<b>RDE</b>	EAST SUFFOLK AND NORTH ESSEX NHS FOUNDATION TRUST	5	393	78	350	49	117	7	49	85	39	11	18	100
<b>RDU</b>	FRIMLEY HEALTH NHS FOUNDATION TRUST	25	2,220	117	338	105	378	97	283	472	160	34	66	334
<b>RDY</b>	DORSET HEALTHCARE UNIVERSITY NHS FOUNDATION TRUST	0	93	0	0	#	0	0	0	70	26	#	#	0
<b>RDZ</b>	THE ROYAL BOURNEMOUTH AND CHRISTCHURCH HOSPITALS NHS FOUNDATION TRUST	#	1,413	0	0	59	149	124	184	505	173	52	78	261
<b>RE9</b>	SOUTH TYNESIDE NHS FOUNDATION TRUST	0	48	0	0	12	86	0	17	58	34	11	25	0
<b>REF</b>	ROYAL CORNWALL HOSPITALS NHS TRUST	30	926	85	164	30	150	7	163	99	73	25	33	244

OFFICIAL

<b>REM</b>	AINTREE UNIVERSITY HOSPITAL NHS FOUNDATION TRUST	13	507	0	134	28	0	46	84	104	51	8	10	195
<b>REP</b>	LIVERPOOL WOMEN'S NHS FOUNDATION TRUST	0	5	0	0	0	391	0	0	0	0	0	0	0
<b>RET</b>	THE WALTON CENTRE NHS FOUNDATION TRUST	0	#	0	0	0	0	0	0	#	0	0	0	0
<b>RF4</b>	BARKING, HAVERING AND REDBRIDGE UNIVERSITY HOSPITALS NHS TRUST	5	1,542	57	391	44	178	137	48	189	27	17	18	201
<b>RFF</b>	BARNSELEY HOSPITAL NHS FOUNDATION TRUST	#	1,070	38	164	33	140	34	85	128	51	32	26	109
<b>RFR</b>	THE ROTHERHAM NHS FOUNDATION TRUST	9	407	35	132	58	168	125	58	219	71	29	30	0
<b>RFS</b>	CHESTERFIELD ROYAL HOSPITAL NHS FOUNDATION TRUST	8	538	29	128	19	191	24	80	228	75	42	44	66
<b>RGN</b>	NORTH WEST ANGLIA NHS FOUNDATION TRUST	17	1,288	102	366	54	311	59	77	532	138	46	76	94
<b>RGP</b>	JAMES PAGET UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	6	790	67	201	32	84	31	63	147	46	10	13	37
<b>RGQ</b>	IPSWICH HOSPITAL NHS TRUST	15	288	39	204	43	197	59	13	478	70	17	35	248

OFFICIAL

<b>RGR</b>	WEST SUFFOLK NHS FOUNDATION TRUST	7	1,185	73	174	30	138	0	34	233	74	45	30	253
<b>RGT</b>	CAMBRIDGE UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	10	353	84	400	23	173	23	35	329	94	56	64	47
<b>RH8</b>	ROYAL DEVON AND EXETER NHS FOUNDATION TRUST	19	1,733	133	230	61	291	15	65	195	171	37	96	128
<b>RHM</b>	UNIVERSITY HOSPITAL SOUTHAMPTON NHS FOUNDATION TRUST	0	401	36	175	11	223	11	29	221	38	38	24	23
<b>RHQ</b>	SHEFFIELD TEACHING HOSPITALS NHS FOUNDATION TRUST	51	2,374	0	116	109	338	252	119	677	168	133	143	207
<b>RHU</b>	PORTSMOUTH HOSPITALS NHS TRUST	25	554	94	348	52	368	9	51	313	129	34	77	29
<b>RHW</b>	ROYAL BERKSHIRE NHS FOUNDATION TRUST	20	312	44	182	8	97	19	43	183	44	9	27	118
<b>RJ1</b>	GUY'S AND ST THOMAS' NHS FOUNDATION TRUST	73	903	48	343	110	160	19	70	255	63	80	60	416
<b>RJ2</b>	LEWISHAM AND GREENWICH NHS TRUST	0	1,062	68	539	61	222	#	54	333	52	40	60	82
<b>RJ6</b>	CROYDON HEALTH SERVICES NHS TRUST	7	266	18	108	40	135	0	49	163	37	33	29	138
<b>RJ7</b>	ST GEORGE'S UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	45	586	29	241	97	79	0	#	209	47	47	57	396

OFFICIAL

<b>RJC</b>	SOUTH WARWICKSHIRE NHS FOUNDATION TRUST	15	391	48	173	22	126	#	84	330	86	40	53	62
<b>RJE</b>	UNIVERSITY HOSPITALS OF NORTH MIDLANDS NHS TRUST	34	1,148	97	279	70	434	10	52	180	101	16	31	419
<b>RJF</b>	BURTON HOSPITALS NHS FOUNDATION TRUST	#	317	61	251	28	180	8	6	186	51	15	15	15
<b>RJL</b>	NORTHERN LINCOLNSHIRE AND GOOLE NHS FOUNDATION TRUST	20	262	15	223	45	278	8	75	161	71	22	28	0
<b>RJN</b>	EAST CHESHIRE NHS TRUST	7	140	7	101	14	77	34	24	77	55	17	12	0
<b>RJR</b>	COUNTESS OF CHESTER HOSPITAL NHS FOUNDATION TRUST	17	655	44	124	#	119	8	20	141	56	17	26	58
<b>RJZ</b>	KING'S COLLEGE HOSPITAL NHS FOUNDATION TRUST	8	2,307	60	188	139	278	26	131	385	129	51	97	434
<b>RK5</b>	SHERWOOD FOREST HOSPITALS NHS FOUNDATION TRUST	#	1,001	103	271	33	153	9	124	248	99	22	59	96
<b>RK9</b>	UNIVERSITY HOSPITALS PLYMOUTH NHS TRUST	15	1,007	111	294	34	274	11	93	306	70	18	27	408
<b>RKB</b>	UNIVERSITY HOSPITALS COVENTRY AND WARWICKSHIRE NHS TRUST	36	2,092	66	596	39	235	140	80	442	65	26	34	300
<b>RKE</b>	WHITTINGTON HEALTH NHS TRUST	0	671	0	0	67	34	0	21	119	28	22	26	32

OFFICIAL

<b>RL1</b>	THE ROBERT JONES AND AGNES HUNT ORTHOPAEDIC HOSPITAL NHS FOUNDATION TRUST	0	84	0	0	0	0	0	127	284	94	33	51	0
<b>RL4</b>	THE ROYAL WOLVERHAMPTON NHS TRUST	12	452	48	234	66	293	9	91	503	130	38	62	364
<b>RLN</b>	CITY HOSPITALS SUNDERLAND NHS FOUNDATION TRUST	0	655	175	447	27	203	231	29	233	68	42	41	624
<b>RLQ</b>	WYE VALLEY NHS TRUST	5	848	92	70	15	54	#	16	309	49	25	26	100
<b>RLT</b>	GEORGE ELIOT HOSPITAL NHS TRUST	0	290	0	0	20	76	0	36	205	29	10	32	38
<b>RM1</b>	NORFOLK AND NORWICH UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	12	2,069	94	290	80	336	21	55	150	230	39	25	474
<b>RM2</b>	UNIVERSITY HOSPITAL OF SOUTH MANCHESTER NHS FOUNDATION TRUST	19	414	9	77	33	63	0	29	111	43	#	19	216
<b>RM3</b>	SALFORD ROYAL NHS FOUNDATION TRUST	0	246	50	118	30	106	0	21	321	73	33	51	0
<b>RMC</b>	BOLTON NHS FOUNDATION TRUST	11	420	55	193	9	128	11	34	158	24	14	23	73
<b>RMP</b>	TAMESIDE AND GLOSSOP INTEGRATED CARE NHS FOUNDATION TRUST	#	236	51	189	46	132	#	105	127	41	16	20	88

OFFICIAL

<b>RN3</b>	GREAT WESTERN HOSPITALS NHS FOUNDATION TRUST	12	231	50	181	52	182	#	22	149	81	33	48	91
<b>RN5</b>	HAMPSHIRE HOSPITALS NHS FOUNDATION TRUST	14	555	73	231	36	195	25	58	367	89	36	54	14
<b>RN7</b>	DARTFORD AND GRAVESHAM NHS TRUST	8	398	#	0	166	203	0	81	233	45	42	37	162
<b>RNA</b>	THE DUDLEY GROUP NHS FOUNDATION TRUST	14	1,506	63	204	16	208	23	56	298	51	22	27	449
<b>RNH</b>	NEWHAM UNIVERSITY HOSPITAL NHS TRUST	0	#	0	0	0	0	0	0	0	0	0	0	0
<b>RNL</b>	NORTH CUMBRIA UNIVERSITY HOSPITALS NHS TRUST	14	118	26	221	11	241	#	40	274	76	15	37	176
<b>RNQ</b>	KETTERING GENERAL HOSPITAL NHS FOUNDATION TRUST	#	344	59	126	59	181	7	41	68	44	13	29	52
<b>RNS</b>	NORTHAMPTON GENERAL HOSPITAL NHS TRUST	10	467	34	86	27	208	50	58	141	46	22	22	119
<b>RNZ</b>	SALISBURY NHS FOUNDATION TRUST	26	717	37	121	15	84	8	135	31	33	9	29	62
<b>RP4</b>	GREAT ORMOND STREET HOSPITAL FOR CHILDREN NHS FOUNDATION TRUST	0	952	91	88	0	0	0	0	#	0	0	0	0
<b>RP5</b>	DONCASTER AND BASSETLAW TEACHING	17	366	98	180	52	205	5	123	301	124	98	70	84

OFFICIAL

	HOSPITALS NHS FOUNDATION TRUST													
<b>RP6</b>	MOORFIELDS EYE HOSPITAL NHS FOUNDATION TRUST	0	#	0	0	0	0	1,098	0	0	0	0	0	0
<b>RPA</b>	MEDWAY NHS FOUNDATION TRUST	10	267	83	465	31	249	0	82	95	13	16	11	102
<b>RPC</b>	QUEEN VICTORIA HOSPITAL NHS FOUNDATION TRUST	64	942	0	18	0	0	#	0	208	96	48	39	#
<b>RPY</b>	THE ROYAL MARSDEN NHS FOUNDATION TRUST	56	203	0	#	#	7	0	0	0	0	0	0	0
<b>RQ3</b>	BIRMINGHAM WOMEN'S AND CHILDREN'S NHS FOUNDATION TRUST	0	458	232	246	0	279	7	0	0	0	6	0	0
<b>RQ6</b>	ROYAL LIVERPOOL AND BROADGREEN UNIVERSITY HOSPITALS NHS TRUST	32	110	0	136	46	#	42	109	160	41	26	47	11
<b>RQ8</b>	MID ESSEX HOSPITAL SERVICES NHS TRUST	47	1,470	112	249	25	193	6	39	221	114	41	67	164
<b>RQM</b>	CHELSEA AND WESTMINSTER HOSPITAL NHS FOUNDATION TRUST	11	664	79	288	100	144	5	46	244	72	51	50	142
<b>RQW</b>	THE PRINCESS ALEXANDRA HOSPITAL NHS TRUST	5	151	42	205	44	68	7	32	121	39	14	13	159
<b>RQX</b>	HOMERTON UNIVERSITY	#	322	0	107	60	33	0	#	91	32	81	39	0

OFFICIAL

	HOSPITAL NHS FOUNDATION TRUST													
<b>RR1</b>	HEART OF ENGLAND NHS FOUNDATION TRUST	14	1,178	45	292	56	303	#	62	347	109	10	42	632
<b>RR7</b>	GATESHEAD HEALTH NHS FOUNDATION TRUST	18	61	0	0	29	211	0	25	167	98	78	71	114
<b>RR8</b>	LEEDS TEACHING HOSPITALS NHS TRUST	51	1,543	143	418	107	220	43	144	370	125	97	66	561
<b>RRF</b>	WRIGHTINGTON, WIGAN AND LEIGH NHS FOUNDATION TRUST	15	395	39	165	27	159	56	183	569	184	70	113	178
<b>RRJ</b>	THE ROYAL ORTHOPAEDIC HOSPITAL NHS FOUNDATION TRUST	0	94	0	0	0	0	0	50	258	69	10	40	0
<b>RRK</b>	UNIVERSITY HOSPITALS BIRMINGHAM NHS FOUNDATION TRUST	71	769	0	170	31	#	5	26	157	10	16	16	650
<b>RRV</b>	UNIVERSITY COLLEGE LONDON HOSPITALS NHS FOUNDATION TRUST	#	1,423	60	474	83	89	0	44	130	22	20	28	228
<b>RT3</b>	ROYAL BROMPTON & HAREFIELD NHS FOUNDATION TRUST	0	#	0	#	0	0	0	0	0	0	0	0	0
<b>RTD</b>	THE NEWCASTLE UPON TYNE HOSPITALS NHS FOUNDATION TRUST	68	2,018	188	403	55	318	22	#	290	107	56	34	448

OFFICIAL

<b>RTE</b>	GLOUCESTERSHIRE HOSPITALS NHS FOUNDATION TRUST	22	402	60	240	35	210	21	50	558	175	44	76	122
<b>RTF</b>	NORTHUMBRIA HEALTHCARE NHS FOUNDATION TRUST	40	547	0	0	22	242	0	103	449	180	52	93	0
<b>RTG</b>	UNIVERSITY HOSPITALS OF DERBY AND BURTON NHS FOUNDATION TRUST	11	706	80	369	45	365	14	87	686	232	57	121	293
<b>RTH</b>	OXFORD UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	56	2,289	75	322	28	183	7	35	385	130	27	65	260
<b>RTK</b>	ASHFORD AND ST PETER'S HOSPITALS NHS FOUNDATION TRUST	18	355	32	240	66	132	73	78	288	41	14	46	236
<b>RTP</b>	SURREY AND SUSSEX HEALTHCARE NHS TRUST	#	424	60	300	86	209	#	96	312	65	13	33	147
<b>RTR</b>	SOUTH TEES HOSPITALS NHS FOUNDATION TRUST	34	1,878	113	296	63	264	55	21	411	106	18	42	606
<b>RTX</b>	UNIVERSITY HOSPITALS OF MORECAMBE BAY NHS FOUNDATION TRUST	8	300	59	213	62	204	10	30	383	122	29	55	0
<b>RVJ</b>	NORTH BRISTOL NHS TRUST	36	1,688	0	0	71	175	#	150	182	81	16	27	133
<b>RVR</b>	EPSOM AND ST HELIER UNIVERSITY HOSPITALS NHS TRUST	0	211	46	218	52	242	11	53	281	61	51	80	141

OFFICIAL

<b>RVV</b>	EAST KENT HOSPITALS UNIVERSITY NHS FOUNDATION TRUST	0	1,844	120	552	120	290	39	108	243	128	78	48	172
<b>RVW</b>	NORTH TEES AND HARTLEPOOL NHS FOUNDATION TRUST	21	148	0	0	27	166	0	12	124	48	46	20	0
<b>RVY</b>	SOUTHPORT AND ORMSKIRK HOSPITAL NHS TRUST	0	273	20	0	9	124	45	22	103	34	7	9	177
<b>RW1</b>	SOUTHERN HEALTH NHS FOUNDATION TRUST	0	17	0	17	#	0	#	0	36	6	#	#	0
<b>RW3</b>	CENTRAL MANCHESTER UNIVERSITY HOSPITALS NHS FOUNDATION TRUST	0	363	92	242	56	108	151	12	62	27	16	7	47
<b>RW6</b>	PENNINE ACUTE HOSPITALS NHS TRUST	12	597	46	394	51	244	103	148	409	68	50	68	656
<b>RWA</b>	HULL AND EAST YORKSHIRE HOSPITALS NHS TRUST	54	1,253	191	497	101	236	21	97	519	196	33	97	470
<b>RWD</b>	UNITED LINCOLNSHIRE HOSPITALS NHS TRUST	37	813	90	324	68	224	11	90	27	119	19	5	210
<b>RWE</b>	UNIVERSITY HOSPITALS OF LEICESTER NHS TRUST	51	1,452	166	497	126	504	196	76	478	97	29	61	157
<b>RWF</b>	MAIDSTONE AND TUNBRIDGE WELLS NHS TRUST	24	147	48	328	47	231	7	35	113	42	19	23	85
<b>RWG</b>	WEST HERTFORDSHIRE	16	426	96	338	58	166	68	102	196	40	22	31	253

OFFICIAL

	HOSPITALS NHS TRUST													
<b>RWH</b>	EAST AND NORTH HERTFORDSHIRE NHS TRUST	9	427	131	285	53	146	117	14	315	82	48	53	129
<b>RWJ</b>	STOCKPORT NHS FOUNDATION TRUST	#	287	53	195	39	140	7	75	342	115	38	37	#
<b>RWP</b>	WORCESTERSHIRE ACUTE HOSPITALS NHS TRUST	9	484	101	225	19	284	6	57	484	80	43	32	345
<b>RWW</b>	WARRINGTON AND HALTON HOSPITALS NHS FOUNDATION TRUST	5	207	28	110	45	57	54	35	153	37	11	30	205
<b>RWY</b>	CALDERDALE AND HUDDERSFIELD NHS FOUNDATION TRUST	9	470	158	364	27	153	11	70	335	66	34	15	289
<b>RX1</b>	NOTTINGHAM UNIVERSITY HOSPITALS NHS TRUST	60	789	107	302	61	239	81	145	43	53	14	5	0
<b>RXC</b>	EAST SUSSEX HEALTHCARE NHS TRUST	6	605	35	273	58	194	38	60	143	55	36	22	128
<b>RXF</b>	MID YORKSHIRE HOSPITALS NHS TRUST	35	1,010	120	410	66	232	120	73	148	217	126	86	420
<b>RXH</b>	BRIGHTON AND SUSSEX UNIVERSITY HOSPITALS NHS TRUST	14	1,226	58	333	64	177	9	98	287	96	42	65	316
<b>RXK</b>	SANDWELL AND WEST BIRMINGHAM HOSPITALS NHS TRUST	27	758	33	292	77	211	184	37	27	41	8	7	229
<b>RXL</b>	BLACKPOOL TEACHING HOSPITALS NHS FOUNDATION TRUST	14	1,468	51	211	12	119	17	13	107	90	13	26	0

OFFICIAL

<b>RXN</b>	LANCASHIRE TEACHING HOSPITALS NHS FOUNDATION TRUST	25	1,796	47	230	15	137	40	43	113	62	49	15	531
<b>RXP</b>	COUNTY DURHAM AND DARLINGTON NHS FOUNDATION TRUST	28	2,331	216	217	59	298	46	60	237	80	55	40	109
<b>RXQ</b>	BUCKINGHAMSHIRE HEALTHCARE NHS TRUST	19	351	119	35	30	113	5	18	274	132	16	43	138
<b>RXR</b>	EAST LANCASHIRE HOSPITALS NHS TRUST	13	592	118	323	64	210	124	50	350	81	55	54	552
<b>RXW</b>	SHREWSBURY AND TELFORD HOSPITAL NHS TRUST	#	405	52	145	42	202	6	98	166	46	25	25	274
<b>RY8</b>	DERBYSHIRE COMMUNITY HEALTH SERVICES NHS FOUNDATION TRUST	0	267	0	0	12	0	61	18	103	57	10	16	52
<b>RYJ</b>	IMPERIAL COLLEGE HEALTHCARE NHS TRUST	84	557	26	275	116	245	139	28	174	26	18	16	1,064
<b>RYR</b>	WESTERN SUSSEX HOSPITALS NHS FOUNDATION TRUST	10	1,608	85	205	30	297	#	125	422	106	39	80	205
<b>RYV</b>	CAMBRIDGESHIRE COMMUNITY SERVICES NHS TRUST	0	0	#	0	0	0	0	0	0	0	0	0	0
<b>Total</b>		2,392	116,676	8,678	32,345	8,493	27,683	6,042	14,020	44,545	14,394	6,239	7,798	28,908

# **Appendix 1: Equality and Health Inequalities Analysis Form**



# **Evidence-Based Interventions Policy: Equality and Health Inequalities – Full Analysis Form**

## **Evidence-Based Interventions Policy: Equalities and Health Inequalities Full Analysis Form**

Version number: Final

First published: 28 November 2018

To be read in conjunction with the Evidence-Based Interventions Policy: Response to public consultation and next steps document

Classification: OFFICIAL

Gateway reference: 08659

This information can be made available in alternative formats, such as easy read or large print, and may be available in alternative languages, upon request. Please email: [england.EBinterventions@nhs.net](mailto:england.EBinterventions@nhs.net)

PART A: General Information
<p><b>1. Title of project, programme or work:</b> Evidence-Based Interventions Programme</p>
<p><b>2. What are the intended outcomes?</b></p> <p>In July 2018, we launched a consultation on the design and implementation of a new programme to ensure interventions routinely available on the NHS are evidence-based and appropriate. The aim of the programme is to prevent avoidable harm to patients and to free up clinical time. Any savings arising from the reduction in interventions will be recycled back into local patient care.</p> <p>Our research has shown that some interventions are not appropriate in certain circumstances, and on occasion, a less invasive but appropriate alternative is available. 17 interventions that fall under this category formed the basis of our consultation and were grouped in to 2 categories. Category 1 interventions which should not be routinely commissioned or performed, and Category 2 interventions which should only be routinely commissioned or performed when specific criteria are met (see appendix C).</p> <p>We believe that our proposals are consistent with National Institute for Health and Care Excellence (NICE), NICE-accredited and specialist society guidance which reflects the most current clinical evidence available. Therefore, we have decided to issue the criteria for the 17 interventions under Section 14Z8 of the NHS Act 2006 as commissioning guidance. This means that CCGs should by April 2019, have 'regard to' the commissioning guidance, in accordance with the Health and Social Care Act. It is for individual CCGs to determine how they do this.</p> <p>However, none of these interventions will be subject to a blanket ban. Category 1 interventions, which are appropriate in exceptional circumstances, will be available via the Individual Funding Request (IFR) process and Category 2 interventions will be available where patients meet the agreed clinical criteria set out in the guidance.</p>
<p><b>3. Who will be affected by this project, programme or work? Please summarise in a few sentences which of the groups below are very likely to be affected by this work.</b></p> <ul style="list-style-type: none"> <li>• Patients – who already receive these interventions or have conditions that would result in a referral for one of these interventions.</li> <li>• Staff: <ul style="list-style-type: none"> <li>○ commissioners who make decisions about their commissioning policies, payment proposals and local systems such as prior approval and IFR processes</li> <li>○ primary care staff, in particular, General Practitioners, as they will need to take account of this guidance when assessing and referring patients as well as offer the alternatives recommended</li> </ul> </li> </ul>

- secondary care clinicians who also need to take account of this guidance when treating patients
- other staff groups (e.g. physiotherapy, nutritionists) who will have a role in offering patients' alternative treatments.
- Partner organisations - (NICE, NHS Clinical Commissioners (NHSCC), Academy of Medical Royal Colleges (AoMRC) and NHS Improvement (NHSI)) have played a key role in finalising the guidance and will support implementation of the changes.

#### **4. Which groups protected by the Equality Act 2010 and/ or groups that face health inequalities are very likely to be affected by this work?**

##### **Proposals for clinical guidance**

The key consideration of this programme is about equitable access to appropriate, evidence-based interventions. We must also ensure patients are not referred for inappropriate interventions that do not meet their needs. Any savings arising from a reduction in referrals for the 17 interventions will be reinvested to provide appropriate interventions to better meet patient's needs.

Current commissioning guidance for these interventions varies between CCGs across England, which could result in inequalities to the wider population through inappropriate referrals and ineffective use of NHS resources. Resources used on these interventions may reduce the availability of resources on more evidence-based and appropriate treatments. By undertaking this work, we aim to reduce variation of inequalities in health outcomes for the wider population by systematically offering the most up-to-date clinically proven treatments and making the most effective use of NHS resources.

The profile of people who are currently being referred for these interventions has been interrogated by age, sex and ethnicity (Source: SUS), no data is available in respect of the other protected characteristics, but comments from consultees in relation to these groups have been considered. The results show that these interventions are accessed by all age groups, gender and ethnicity. However, some interventions are accessed more (or solely) by a specific group, such as grommets for glue ear in children (children) and hysterectomy for menstrual bleeding (women), but overall, as this guidance applies to the whole population all groups protected by the Equality Act 2010 and/or groups that face health inequalities will be affected by this work.

##### **Consultation**

A 12-week consultation was carried out between July 4<sup>th</sup> and September 28<sup>th</sup>, 2018. This offered an opportunity for views to be sought from people representing many of the equality groups referred to in this equality and health inequalities impact assessment. Therefore, we included a specific question about the impact on equality and health inequality groups in the Evidence-Based Interventions consultation, see appendix B as well as working directly with individuals from equality groups.

We received 707 online responses and 97 individual submissions. We also spoke to 397 individuals by hosting or attending a number of events, including:

- Patient and public face to face events in Birmingham, London and Leeds

- Workshops with individuals with learning disabilities in Leeds and London
- NHS Expo conference in Manchester
- NHS Improvement costing forums in Leeds, Birmingham and London
- Guidelines International Network conference in Manchester
- Seven online webinars with; Health and Wellbeing Alliance; Healthwatch; NHS Clinical Commissioners; NHS Youth Forum; and Voluntary Sector and Community Enterprises.

Key themes from the analysis of the responses relevant to equality and health inequalities impact assessment have been reflected throughout this document. They have also been taken into account in the Evidence-Based Interventions Policy: Response to the public consultation and next steps document.

## PART B: Equalities Groups and Health Inequalities Groups

### 5. Impact of this work for the equality groups listed below.

Focusing on each equality group listed below (sections 5.1. to 5.9), please answer the following questions:

- Does the equality group face discrimination in this work area?
- Could the work tackle this discrimination and/or advance equality or good relations?
- Has due regard been paid to the need to:
  - eliminate discrimination, harassment, victimisation and any other conduct that is prohibited by or under this Act;
  - advance equality of opportunity between persons who share a relevant protected characteristic and persons who do not share it (in particular, by removing or minimising disadvantages arising from that characteristic, meeting particular needs of persons with a protected characteristic, and encouraging people with a protected characteristic to participate in public life or other activity where participation is disproportionately low);
  - foster good relations between persons who share a relevant protected characteristic and persons who do not share it (in particular, by tackling prejudice and promoting understanding).
- Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?
- If you cannot answer these questions what action will be taken and when?

#### 5.1. Age


##### Does the equality group face discrimination in this work area?

Looking at the age profiles of patients referred in 2017/18, the prevalence for these interventions vary across children/young people, adults and older people although the majority are within the 18-64 group which is in line with all elective care. Overall the data demonstrates that some interventions have a similar age profile to all elective interventions and where this differs, such as for grommets, haemorrhoids and varicose veins they are consistent with the age groups at which the underlying problem is most prevalent.

Following extensive consultation with clinical specialists and CCGs, we have removed any restriction on children from the criteria for trigger finger release, Dupuytren’s contracture release and snoring surgery. This is so our clinical criteria is based on clinical evidence developed by NICE, NICE-accredited or specialist society guidance.

In addition, some respondents stated children and young people should have access to information that supports them to make an informed decision about their care and treatment where necessary.

**Chart 1: Percentage of patients receiving each intervention in 2017/18 by age**



**Chart 1: Children and young people are excluded from snoring surgery, knee arthroscopy for patients with osteoarthritis, Dupuytren’s contracture release and trigger finger release and grommets for glue ear is specific to children and young people only**

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

The clinical guidance has been reviewed and amended to take account of children and young people. For example, the impact on children and young people's mental health has been added as a criterion for removal of benign skin lesions and children and young people have been removed from the clinical criteria for trigger finger release, Dupuytren's contracture release and snoring surgery.

We will continue to use the NHS Youth Forum in an advisory capacity to seek the views of children and young people, to help co-produce materials and information that is accessible.

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

**5.2. Disability**

**Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and disability so we cannot definitively assess, at a national level, how many people with a disability will be affected.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. A number of people raised issues that vulnerable groups, such as people with a learning disability may be disadvantaged from these proposals. They may not understand why an intervention is not being offered and they may not have the ability to voice their opinion to challenge a decision.

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less

invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

There are no expected adverse impacts on the clinically based decisions. However, this protected group as with children and young people may need extra support in understanding the decisions taken, the alternative options and how to access the IFR process.

We will use our national steering group (membership includes patient representatives, The Patients Association and National Voices) and the existing patient networks our steering group partners have access to help co-produce materials and information to support implementation, in particular this equality group.

We will emphasise the need for an advocate to support vulnerable groups, such as individuals with a learning disability, when attending a doctor's appointment to support discussions about what the most appropriate treatment is for the individual.

We will produce easy read pamphlets on the 17 interventions to describe the changes we are implementing.

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

**5.3. Gender reassignment**

**Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and gender reassignment so we cannot definitively assess, at a national level, how many people will be affected. None of the interventions included in the proposed guidance are used for the purposes of gender reassignment as it is specific to breast hyperplasia.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. There were no results from the consultation that indicated this for gender reassignment.

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less

invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

#### **5.4. Marriage and civil partnership**

**Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and marriage/civil partnership so we cannot definitively assess, at a national level, how many people in a marriage/civil partnership will be affected.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. There were no results from the consultation that indicated this for people in a marriage/civil partnership.

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

#### **5.5. Pregnancy and maternity**

**Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and pregnancy/maternity so we cannot definitively assess, at a national level, how many people will be affected. None of the interventions in the guidance are used for conditions that are closely related to pregnancy or maternity.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. There were no results from the consultation that indicated this for pregnancy or maternity.

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.


**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

**5.6. Race****Does the equality group face discrimination in this work area?**

Looking at the ethnicity profiles of patients referred in 2017/18, the prevalence for these interventions are similar to all elective care. The majority of the analysis demonstrated no substantial difference between the proportion of these interventions that are accessed by ethnic groups compared to the white British group when you take account of the different age groups. The exceptions are for chalazia removal which is less common in the white British group (48%) and higher in the Asian group (12%) and unknown groups (22%) and Dupuytren's contracture release in adults which is more common in the white British group (80%) compared to the others, although this is expected due to the increased occurrence in people of white European descent.

*Chart 2: Percentage of patients receiving each intervention by ethnicity*



**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

The chalazia removal and Dupuytren's contracture release clinical criteria has been reviewed to ensure it is based on NICE, NICE-accredited and specialist society guidance

and that the interventions will still be available to people who meet the criteria and in exceptional cases through an individual funding review where appropriate.

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

### **5.7. Religion or belief**

#### **Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and religion or belief, so we cannot definitively assess, at a national level, how many people will be affected. We have not identified any religious beliefs that would make an individual more or less likely to receive the interventions included in the guidance.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. There were no results from the consultation that indicated this for religion or belief.

#### **Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

#### **Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

#### **Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**


CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve

### **5.8. Sex or gender**

#### **Does the equality group face discrimination in this work area?**

Overall the data demonstrates that on average slightly more women are referred for both the category 1 (60%) and the category 2 (56%) interventions than males. This is because there are two interventions that are provided only to women (menstrual dilatation and curettage and hysterectomy), and one which is predominantly women (breast reduction). Because of this a number of the consultation responses referred to gender as the equality group and women as the equality characteristic that was most likely to be disproportionately affected by this work.

Chart 3: Percentage of patients receiving each intervention by gender



**Could the work tackle this discrimination and/or advance equality or good relations?**

As a result of this concern we engaged directly with organisations representing women by inviting them to respond to the consultation. We have worked with the Royal College of Gynaecologists and Obstetricians and used NICE guidance to ensure our clinical criteria for women-specific conditions are based on the most up-to-date research, evidence and professional opinion.

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

Taking into account the consultation results we are continuing to engage with organisations that advocate for women and inviting them to contribute to the co-production of materials and information to support implementation.

CCGs will be required to assess the impact on their population with regard to the particular demographics of the population they serve.

### **5.9. Sexual orientation**

**Does the equality group face discrimination in this work area?**

There is no routinely collected data on access to the 17 interventions and sexual orientation so we cannot definitively assess, at a national level, how many people will be affected. There is no established link between the interventions proposed in this guidance and sexual orientation.

During the consultation, responses were monitored to ascertain if there were likely unintended consequences on the equality groups. There were no results from the consultation that indicated this for sexual orientation.

**Could the work tackle this discrimination and/or advance equality or good relations?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment. The most appropriate alternative may be less invasive reducing risk and offering further health benefits where it involves a lifestyle change.

**Could the work assist or undermine compliance with the Public Sector Equality Duty (PSED)?**

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient. The work aims to reduce health inequalities in access and outcomes for all patient groups as well as compliance with the other public sector equality duties by ensuring the offer of appropriate treatment.

**Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?**

CCGs will also be required to assess the impact on their population with regard to the particular demographics of the population they serve.

### **6. Implications of our work for the health inclusion groups listed below.**

Focusing on the work described in sections 1 and 2, in relation to each health inclusion group listed below (Sections 6.1. To 6.12), and any others relevant to your work<sup>31</sup>, please answer the following questions:

- f) Does the health inclusion group experience inequalities in access to healthcare?
- g) Does the health inclusion group experience inequalities in health outcomes?
- h) Could the work be used to tackle any identified inequalities in access to healthcare or health outcomes?
- i) Could the work assist or undermine compliance with the duties to reduce health inequalities?
- j) Does any action need to be taken to address any important adverse impact? If yes, what action should be taken?
- k) As some of the health inclusion groups overlap with equalities groups you may prefer to also respond to these questions about a health inclusion group when responding to 5.1 to 5.9. That is fine; please just say below if that is what you have done.
- l) If you cannot answer these questions what action will be taken and when?

**6.1. Alcohol and / or drug misusers**

There is no data available on the prevalence of alcohol and / or drug misuse with regards to who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

**6.2. Asylum seekers and /or refugees**

There is no data available on the prevalence of asylum seekers and/or refugees who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

**6.3. Carers**

There is no data available on the prevalence of carers who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

**6.4. Ex-service personnel / veterans**

There is no data available on the prevalence of ex-service personnel / veterans who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

**6.5. Those who have experienced Female Genital Mutilation (FGM)**

There is no data available on the prevalence of those who have experienced Female Genital Mutilation (FGM) who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in

this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

#### **6.6. Gypsies, Roma and travellers**

There is no data available on the prevalence of Gypsies, Roma and travellers who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

#### **6.7. Homeless people and rough sleepers**

There is no data available on the prevalence of homeless people and rough sleepers who are currently accessing the interventions in the review.

A number of consultation responses highlighted this guidance could impact on individuals that do not have a fixed address from accessing the necessary treatments.

As part of the delivery actions to support implementation, we have considered what needs to be in place to support referrals for any of these interventions via accident and emergency which will be worked through with demonstrator sites and CCGs / providers going forward. Also, treatment may include less invasive alternatives where appropriate following implementation which would be beneficial for a homeless person or someone who sleeps rough.

#### **6.8. Those who have experienced human trafficking or modern slavery**

There is no data available on the prevalence of those who have experienced human trafficking or modern slavery who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

#### **6.9. Those living with mental health issues**

The interventions are not specific to individuals with mental health issues. However, the inclusion of mental health issues as criterion for why some of these interventions should be offered was highlighted in a number of consultation responses.

This was recognised as an appropriate criterion, resulting in amendments to the clinical criteria for benign skin lesions.

Beyond the need to include mental health as selection criterion, there was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

#### **6.10. Sex workers**

There is no data available on the prevalence of sex workers who are currently accessing these interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.

<p><b>6.11. Trans people or other members of the non-binary community</b></p> <p>There is no data available on trans people or other members of the non-binary community who are currently accessing the interventions in the review. There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.</p>		
<p><b>6.12. The overlapping impact on different groups who face health inequalities</b></p> <p>There is no data available on different groups who face health inequalities who are currently accessing the interventions in the review.</p> <p>There was no indication from the consultation results that the proposals would result in this health inclusion group experiencing inequalities in access to healthcare or health outcomes.</p>		
<p><b>7. Other groups that face health inequalities that we have identified.</b></p> <p><b>Have you have identified other groups that face inequalities in access to healthcare?</b></p> <p><b>Does the group experience inequalities in access to healthcare and/or inequalities in health outcomes?</b></p> <p><b>Short explanatory notes</b> - other groups that face health exclusion. As we research and gather more data, we learn more about which groups are facing health inequalities. If your work has identified more groups that face important health inequalities please answer questions 7 and 8. Please circle as appropriate.</p> <p>If you have not identified additional groups, that face health inequalities, just say not applicable or N/A in the box below.</p>		
<p>Yes Complete section 8</p>	<p>No Go to section 9</p>	<p><b>N/A</b></p>
<p><b>8. Other groups that face health inequalities that we have identified.</b></p> <p>Could the work be used to tackle any identified inequalities in access to healthcare or health outcomes in relation to these other groups that face health inequalities? Could the work undermine compliance with the duties to reduce health inequalities and, if so, what action should be taken to reduce any adverse impact? Is the work going to help NHS England to comply with the duties to reduce health inequalities?</p>		

OFFICIAL

If you have identified other groups that face health inequalities please answer the questions below. You will only answer this question if you have identified additional groups facing important health inequalities.

<b>PART C: Promoting integrated services and working with partners</b>		
Short explanatory notes: Integrated services and reducing health inequalities.		
Our detailed guidance explains the duties in relation to integrated services and reducing health inequalities. Please answer the questions listed below.		
<b>9. Opportunities to reduce health inequalities through integrated services.</b>		
Does the work offer opportunities to encourage integrated services that could reduce health inequalities? If yes please also answer 10.		
Yes Go to section 10	<b>No</b> Go to section 11	Do not know
<b>10. How can this work increase integrated services and reduce health inequalities?</b>		
Please explain below, in a few short sentences, how the work will encourage more integrated services that reduce health inequalities and which partners we will be working with.		
<b>PART D: Engagement and involvement</b>		
<b>11. Engagement and involvement activities already undertaken.</b>		
<b>How were stakeholders, who could comment on equalities and health inequalities engaged, or involved with this work? For example in gathering evidence, commenting on evidence, commenting on proposals or in other ways? And what were the key outputs?</b>		
NHS England has established a programme board with its partner organisations that are all signatories on the consultation and a steering group with all the key stakeholders for the programme. The programme board includes; NHS Clinical Commissioners (NHSCC), Academy of Medical Royal Colleges (AoMRC), NICE and NHS Improvement. The steering group includes representatives from:		
<ul style="list-style-type: none"> <li>• NHSCC</li> <li>• NHSI</li> <li>• NICE</li> <li>• AoMRC</li> <li>• National Voices</li> <li>• Patients Association</li> <li>• Patient representatives</li> <li>• NHS Providers</li> </ul>		

- British Medical Association
- CQC.

A 12-week consultation was carried out between July 4<sup>th</sup> and September 28<sup>th</sup>, 2018. This offered an opportunity for views to be sought from people representing many of the equality groups referred to in this equality and health inequalities impact assessment. Therefore, we included a specific question about the impact on equality and health inequality groups in the evidence-based interventions consultation.

We received 707 online responses and 97 individual submissions. We also spoke to 397 individuals by hosting or attending a number of events, including:

- Patient and public face to face events in Birmingham, London and Leeds
- Workshops with individuals with learning disabilities in Leeds and London
- NHS Expo conference in Manchester
- NHS Improvement costing forums in Leeds, Birmingham and London
- Guidelines International Network conference in Manchester
- Seven online webinars with; Health and Wellbeing Alliance; Healthwatch; NHS Clinical Commissioners; NHS Youth Forum; and Voluntary Sector and Community Enterprises.

Key themes from the analysis of the responses relevant to the equality and health inequalities impact assessment have been reflected throughout this document. They have also been taken account of in the Evidence-Based Interventions Policy: Response to the public consultation and next steps document.

**12. Which stakeholders and equalities and health inclusion groups were involved?**

NHSCC, NHSI, NICE, AoMRC, National Voices, The Patients Association, patient representatives, NHS Providers, NHS Confederation, NHS Partners, British Medical Association and CQC.

The consultation had involvement of a number of stakeholders and equalities and health inclusion groups (see response 11 above).

**13. Key information from the engagement and involvement activities undertaken.**

**Were key issues, concerns or questions expressed by stakeholders and if so what were these and how were they addressed? Were stakeholders broadly supportive of this work?**

Stakeholders are broadly supportive of the work on the proposals for the 17 interventions and concerns relating to the equalities and health inequalities raised by stakeholders are reflected throughout this review.

**14. Stakeholders were not broadly supportive but we need to go ahead.**

**If stakeholders were not broadly supportive of the work but you are recommending progressing with the work anyway, why are you making this recommendation?**

For some of the 17 interventions and implementation mechanisms there are groups that are not broadly supportive of the specific recommendations. Further details can be found in the Evidence-Based Interventions Policy: Response to the public consultation and next steps document (Nov 2018).

**15. Further engagement and involvement activities planned.**

**Are further engagement and involvement activities planned? If so what is planned, when and why?**

We plan to hold a number of further engagement and involvement activities, including:

- Publication of the Evidence-Based Interventions Policy: Response to the public consultation and next steps document that includes the clinical criteria for the 17 interventions end of 2018
- Ongoing engagement throughout January – April 2019 with all sectors (primary care, commissioners, providers and patients and the public) to raise awareness, understanding and embed change to support implementation.
- National steering group meetings with individual patient representatives as well as organisations that represent patients.

In addition, we will we will use existing patient networks from our steering group partners, to help co-produce and advise on materials and information to support implementation.

## PART E: Monitoring and Evaluation

**16. In relation to equalities and reducing health inequalities, please summarise the most important monitoring and evaluation activities undertaken in relation to this work**

Analysis, reporting and consideration of the SUS data and consultation responses.

**17. Please identify the main data sets and sources that you have drawn on in relation to this work. Which key reports or data sets have you drawn on?**

SUS data sources.

Responses to the evidence-based intervention consultation.

<p><b>18. Important equalities or health inequalities data gaps or gaps in relation to evaluation.</b></p> <p><b>In relation to this work have you identified any:</b></p> <ul style="list-style-type: none"> <li>• important equalities or health inequalities data gaps or</li> <li>• gaps in relation to monitoring and evaluation?</li> </ul>	
Yes	No
<p>There is currently no nationally collected data for 6 of the 9 equality groups and additional health improvement groups for the interventions in this review.</p>	
<p><b>19. Planned action to address important equalities or health inequalities data gaps or gaps in relation to evaluation.</b></p> <p>If you have identified important gaps and you have identified action to be taken, what action are you planning to take, when and why?</p> <p>We think that individual CCGs may have more insight on this when looking at their local population data and we will encourage them to consider this as part of local consultation and impact assessments.</p>	

PART F: Summary analysis and recommended action		
<b>20. Contributing to the first PSED equality aim.</b>		
Can this work contribute to eliminating discrimination, harassment or victimisation?		
Yes	No	Do not know
If yes please explain how, in a few short sentences		
<b>21. Contributing to the second PSED equality aim.</b>		
Can this policy or piece of work contribute to advancing equality of opportunity? Please circle as appropriate.		
Yes	No	Do not know
Currently patients could be receiving interventions that are not appropriate for their needs. By setting national direction on when certain interventions should be commissioned this programme encourages NHS commissioners and providers to implement policy about reviewing patients' needs with the doctor to identify the most appropriate treatment for that individual. This enables patients to have access to the most effective treatment to achieve the best outcome, which may be less invasive and offer further health benefits where it is a lifestyle change. Through ensuring effective use of NHS resources, the programme will enable local systems to provide appropriate treatments to optimise wider population benefit and outcomes.		
<b>22. Contributing to the third PSED equality aim.</b>		
Can this policy or piece of work contribute to fostering good relations between groups? Please circle as appropriate.		
Yes	No	Do not know
If yes please explain how, in a few short sentences		
<p>The evidence-based interventions programme is a partnership with NHS Clinical Commissioners, NICE, Academy of Medical Royal Colleges and NHS Improvement. The approach is based on working collaboratively with our partner organisations. An example of this is that all our partners are joint signatories on the consultation and the Evidence-Based Interventions Policy: Response to the public consultation and next steps document which includes the clinical criteria for the 17 interventions.</p> <p>Fostering of good relationships was also enhanced through engagement with a number of other key stakeholders including charities and patient groups prior, during and post consultation. The consultation also provided an opportunity for organisations,</p>		

health professionals, patients and the public to contribute to the development of the guidance and all other outputs and decisions regarding the delivery actions (implementation mechanisms).

We will continue this work through our ongoing engagement programme to support implementation with our national steering group and we will use existing patient networks from our steering group partners.

**23. Contributing to reducing inequalities in access to health services.**

Can this policy or piece of work contribute to reducing inequalities in access to health services?

Yes	No	Do not know
-----	----	-------------

If yes which groups should benefit and how and/or might any group lose out?

When implemented, this guidance should prompt consideration of what is the most appropriate treatment in discussion between the doctor and their patient, meaning patients will receive the most appropriate treatment.

There are also wider population gains for those patients who will receive treatments supported by the resource saved from stopping doing interventions that are not appropriate in some cases and re-directed in to providing treatments that are.

An additional benefit is where an alternative treatment involves a lifestyle change that has an added health benefit for the individual.

**24. Contributing to reducing inequalities in health outcomes.**

Can this work contribute to reducing inequalities in health outcomes?

Yes	No	Do not know
-----	----	-------------

If yes which groups should benefit and how and/or might any group lose out?

As section 23

**25. Contributing to the PSED and reducing health inequalities.**

**How will the policy or piece of work contribute to the achieving the PSED and reducing health inequalities in access and outcomes? Please describe below in a few short sentences.**

As section 23

**26. Agreed or recommended actions.**

What actions are proposed to address any key concerns identified in this Equality and Health Inequalities Analysis (EHIA) and / or to ensure that the work contributes to the reducing unlawful discrimination / acts, advancing equality of opportunity, fostering good relations and / or reducing health inequalities? Is there a need to review the EHI analysis at a later stage?

Action	Public Sector Equality Duty	Health Inequality	By when	By whom
Ensure the opportunity to challenge any decision about accessing these interventions remains through an IFR process and that prior approval is applied appropriately. Processes should be open, transparent and understood by the local population.	Yes	Yes	April 2019	CCGs, clinicians (primary & secondary care)
Produce easy read pamphlets on the 17 interventions to describe the changes we are implementing by December 2018.	Yes	Yes	Support publication of guidance	NHSE, NHSCC, AoMRC, NICE, NHSI
National steering group meetings with individual patient representatives as well as organisations that represent patients. We will use existing patient networks from our steering group partners, to help co-produce and advise on materials and information to support implementation.	Yes	Yes	Post consultation	NHSE, NHSCC, AoMRC, NICE, NHSI
Emphasise the need for an advocate to support vulnerable groups, such as individuals with a learning disability, when attending a doctor's appointment. Include in our supporting tools.	Yes	Yes	Post consultation	CCGs, clinicians (primary & secondary care)

OFFICIAL

Continue to use the NHS Youth Forum in an advisory capacity to seek the views of children and young people, to help co-produce materials and information that is accessible.	Yes	Yes	Post consultation	NHSE, NHSCC, AoMRC, NICE, NHSI
--	-----	-----	-------------------	--------------------------------

## Appendix A: Activity for each intervention by equality group in 2017/2018

Table 1: Number of patients receiving each intervention by **age**

Row Labels	00to17	18to64	65plus	Unknown / missing
<b>All category 1</b>	<b>0.0%</b>	<b>65.0%</b>	<b>34.8%</b>	<b>0.2%</b>
Snoring surgery	n/a	93.3%	6.7%	0.0%
Dilatation and curettage	0.0%	83.1%	14.4%	2.5%
Knee arthroscopy for patients with osteoarthritis	n/a	82.7%	17.3%	0.0%
Injections for non-specific back pain	0.0%	58.3%	41.5%	0.2%
<b>All category 2</b>	<b>11.4%</b>	<b>59.6%</b>	<b>28.6%</b>	<b>0.3%</b>
Breast reduction	0.6%	85.1%	13.6%	0.7%
Removal of benign skin lesions	7.3%	58.1%	34.1%	0.4%
Grommets for Glue Ear in children	100.0%	n/a	n/a	0.0%
Tonsillectomy for recurrent tonsillitis	55.1%	43.8%	0.9%	0.2%
Haemorrhoid surgery	0.1%	79.2%	19.9%	0.8%
Hysterectomy for heavy menstrual bleeding	0.0%	79.4%	20.0%	0.6%
Chalazia removal	16.3%	72.8%	10.6%	0.3%
Arthroscopic shoulder decompression for subacromial pain	0.0%	78.9%	21.0%	0.1%
Carpal tunnel syndrome release	0.1%	58.8%	41.0%	0.2%
Dupuytren's contracture release in adults	n/a	40.2%	59.8%	0.0%
Ganglion excision	4.9%	75.1%	19.8%	0.3%
Trigger finger release	n/a	61.1%	38.9%	0.0%
Varicose veins	0.1%	69.5%	30.1%	0.3%
<b>Other non-emergency spells</b>	<b>11.6%</b>	<b>52.1%</b>	<b>35.1%</b>	<b>1.2%</b>
<b>All non-emergency spells</b>	<b>11.6%</b>	<b>52.3%</b>	<b>34.9%</b>	<b>1.2%</b>

Table 1: Children and young people are excluded from snoring surgery, knee arthroscopy for patients with osteoarthritis, Dupuytren's contracture release and trigger finger release and grommets for glue ear is specific to children and young people only

OFFICIAL

**Appendix A:** Activity for each intervention by equality group in 2017/2018

*Table 2: Number of patients receiving each intervention by ethnicity*

Row Labels	White British	Other White (not-British)	Black	Asian (including Chinese)	Mixed	Other	Unknown / missing
<b>All category 1</b>	<b>78.5%</b>	<b>3.7%</b>	<b>2.0%</b>	<b>3.3%</b>	<b>0.5%</b>	<b>0.9%</b>	<b>11.1%</b>
Snoring surgery	70.6%	4.5%	2.8%	3.9%	1.2%	1.6%	15.4%
Dilatation and curettage	73.7%	4.9%	5.8%	4.9%	1.2%	0.8%	8.6%
Knee arthroscopy for patients with osteoarthritis	76.8%	3.4%	1.9%	3.6%	0.6%	0.9%	12.9%
Injections for non-specific back pain	79.5%	3.7%	1.9%	3.2%	0.5%	0.9%	10.4%
<b>All category 2</b>	<b>71.1%</b>	<b>4.5%</b>	<b>1.8%</b>	<b>4.1%</b>	<b>0.9%</b>	<b>1.4%</b>	<b>16.2%</b>
Breast reduction	73.7%	3.3%	3.3%	2.6%	1.6%	1.2%	14.4%
Removal of benign skin lesions	70.4%	4.6%	1.6%	3.3%	0.7%	1.2%	18.2%
Grommets for Glue Ear in children	75.6%	3.5%	1.4%	6.0%	2.4%	1.3%	9.7%
Tonsillectomy for recurrent tonsillitis	67.7%	4.8%	2.1%	6.4%	2.3%	2.1%	14.7%
Haemorrhoid surgery	61.2%	6.7%	4.6%	7.1%	1.5%	2.6%	16.2%
Hysterectomy for heavy menstrual bleeding	75.0%	3.5%	2.6%	4.3%	0.8%	1.2%	12.6%
Chalazia removal	48.0%	4.6%	5.2%	13.4%	1.6%	4.8%	22.4%
Arthroscopic shoulder decompression for subacromial pain	77.4%	3.2%	1.4%	2.7%	0.6%	0.8%	14.0%
Carpal tunnel syndrome release	74.8%	4.1%	1.6%	3.2%	0.5%	1.1%	14.8%
Dupuytren's contracture release in adults	80.6%	2.4%	0.3%	0.4%	0.2%	0.3%	15.8%
Ganglion excision	67.7%	4.3%	3.0%	4.1%	1.1%	1.5%	18.3%
Trigger finger release	75.3%	3.2%	1.9%	3.3%	0.5%	1.1%	14.7%
Varicose veins	65.8%	7.1%	1.1%	5.6%	0.6%	2.1%	17.7%
<b>Other non-emergency spells</b>	<b>70.7%</b>	<b>5.3%</b>	<b>2.6%</b>	<b>5.7%</b>	<b>1.2%</b>	<b>1.7%</b>	<b>12.7%</b>
<b>All non-emergency spells</b>	<b>70.8%</b>	<b>5.2%</b>	<b>2.6%</b>	<b>5.7%</b>	<b>1.2%</b>	<b>1.7%</b>	<b>12.8%</b>

**Appendix A:** Activity for each intervention by equality group in 2017/2018*Table 3: Number of patients receiving each intervention by gender*

Row Labels	Male	Female	Unknown / missing
<b>All category 1</b>	<b>37.9%</b>	<b>62.1%</b>	<b>0.0%</b>
Snoring surgery	73.6%	26.4%	0.0%
Dilatation and curettage	0.0%	100.0%	0.0%
Knee arthroscopy for patients with osteoarthritis	48.5%	51.5%	0.0%
Injections for non-specific back pain	33.6%	66.4%	0.0%
<b>All category 2</b>	<b>41.9%</b>	<b>58.1%</b>	<b>0.0%</b>
Breast reduction	1.2%	98.8%	0.0%
Removal of benign skin lesions	48.8%	51.2%	0.0%
Grommets for Glue Ear in children	58.1%	41.9%	0.0%
Tonsillectomy for recurrent tonsillitis	38.3%	61.7%	0.0%
Haemorrhoid surgery	51.4%	48.6%	0.0%
Hysterectomy for heavy menstrual bleeding	0.1%	99.9%	0.0%
Chalazia removal	52.5%	47.5%	0.0%
Arthroscopic shoulder decompression for subacromial pain	46.8%	53.2%	0.0%
Carpal tunnel syndrome release	35.2%	64.8%	0.0%
Dupuytren's contracture release in adults	77.1%	22.9%	0.0%
Ganglion excision	31.5%	68.5%	0.0%
Trigger finger release	43.4%	56.6%	0.0%
Varicose veins	44.0%	56.0%	0.0%
<b>Other non-emergency spells</b>	<b>42.8%</b>	<b>57.2%</b>	<b>0.0%</b>
<b>All non-emergency spells</b>	<b>42.8%</b>	<b>57.2%</b>	<b>0.0%</b>

**Appendix B:** Evidence-Based Interventions consultation question and key themes from the analysis of responses

**Q14. What positive and negative impact will these changes make to improving access, experience and outcomes for the following groups and how can any risks be mitigated to ensure the changes do not worsen health inequalities for:**

- **groups protected under the Equality Act 2010?**
- **those individuals who experience health inequalities such as homeless people/rough sleepers, vulnerable migrants, gypsy traveller groups and carers?**

<b>Respondent type</b>	<b>Key themes - summary</b>
Clinician	<ul style="list-style-type: none"> <li>• 2 respondents stated there would be a positive impact</li> <li>• The equality groups respondents stated as having a potential negative impact on included; vulnerable groups (3) or women (2)</li> </ul>
CCG	<ul style="list-style-type: none"> <li>• 1 respondent stated there would be no negative impact</li> <li>• The equality group respondents stated as having a potential negative impact on was women (2)</li> </ul>
National body	<ul style="list-style-type: none"> <li>• 4 respondents stated there would be no negative impact and 2 respondents stated there would be a positive impact</li> <li>• The equality groups respondents stated as having a potential negative impact on included; access in general (1) or vulnerable groups (2)</li> </ul>
NHS provider organisation	<ul style="list-style-type: none"> <li>• 2 respondents stated there would be a positive impact</li> <li>• The equality groups respondents stated as having a potential negative impact on included; access in general (3) or vulnerable groups (1)</li> </ul>
Other / unknown	<ul style="list-style-type: none"> <li>• 3 respondents stated there would be no negative impact and 2 respondents stated there would be a positive impact</li> <li>• The equality groups respondents stated as having a potential negative impact on included; vulnerable groups (12), individuals without a permanent address (1) or women (1)</li> </ul>
Patient / member of the public	<ul style="list-style-type: none"> <li>• 13 respondents stated there would be no negative impact</li> <li>• The equality groups respondents stated as having a potential negative impact on included; access in general (103), women (76), vulnerable groups (27), individuals without a permanent address (5) or travellers (1)</li> </ul>

OFFICIAL

Patient representative organisation	<ul style="list-style-type: none"><li>• One respondent stated there would be no negative impact and 1 respondent stated there would be a positive impact</li><li>• The equality groups respondents stated as having a potential negative impact on included; access in general (6) or individuals without a permanent address (1)</li></ul>
VSO / Charity	<ul style="list-style-type: none"><li>• The equality groups respondents stated as having a potential negative impact on included; access in general (6) or women (4)</li></ul>

OFFICIAL

**Appendix C:** The list of 17 interventions grouped into **Category 1** and **Category 2**

<b>Category 1:</b> Interventions which should not be routinely commissioned or performed	
<i>Intervention</i>	<i>Summary of intervention</i>
<b>ENT</b>	
Snoring surgery	<p>Snoring is a noise that occurs during sleep that can be caused by vibration of tissues of the throat and palate. It is very common and as many as one in four adults snore, as long as it is not complicated by periods of apnoea (temporarily stopping breathing) it is not usually harmful to health, but can be disruptive, especially to a person's partner.</p> <p>This guidance relates to surgical procedures in adults to remove, refashion or stiffen the tissues of the soft palate (Uvulopalatopharyngoplasty, Laser assisted Uvulopalatoplasty &amp; Radiofrequency ablation of the palate) in an attempt to improve the symptom of snoring. Please note this guidance only relates to patients with snoring in the absence of Obstructive Sleep Apnoea (OSA) and should not be applied to the surgical treatment of patients who snore and have proven OSA who may benefit from surgical intervention as part of the treatment of the OSA.</p> <p>It is important to note that snoring can be associated with multiple other causes such as being overweight, smoking, alcohol or blockage elsewhere in the upper airways (e.g. nose or tonsils) and often these other causes can contribute to the noise alongside vibration of the tissues of the throat and palate.</p>
<b>Gynaecology</b>	
Dilatation and curettage for heavy menstrual bleeding	Dilation and curettage (D&C) is a minor surgical procedure where the opening of the womb (cervix) is widened (dilatation) and the lining of the womb is scraped out (curettage).
<b>Orthopaedics</b>	
Knee arthroscopy for patients with osteoarthritis	Arthroscopic washout of the knee is an operation where an arthroscope (camera) is inserted in to the knee along with fluid. Occasionally loose debris drains out with the fluid, or debridement, (surgical removal of damaged cartilage) is performed, but the procedure does not improve symptoms or function of the knee joint.
Injections for non-specific low back pain	Spinal injections of local anaesthetic and steroid in people with non-specific low back pain without sciatica.

OFFICIAL

<b>Category 2:</b> Interventions which should only be routinely commissioned or performed when specific criteria are met	
<i>Intervention</i>	<i>Summary of intervention</i>
<b>General surgery</b>	
Breast reduction	Breast reduction surgery is a procedure used to treat women with breast hyperplasia (enlargement), where breasts are large enough to cause problems like shoulder girdle dysfunction, intertrigo and adverse effects to quality of life.
<b>Dermatology</b>	
Removal of benign skin lesions	Removal of benign skin lesions means treating asymptomatic lumps, bumps or tags on the skin that are not suspicious of cancer. Treatment carries a small risk of infection, bleeding or scarring and is not usually offered by the NHS if it is just to improve appearance. In certain cases, treatment (surgical excision or cryotherapy) may be offered if certain criteria are met. A patient with a skin or subcutaneous lesion that has features suspicious of malignancy must be treated or referred according to NICE skin cancer guidelines. This policy does not refer to pre-malignant lesions and other lesions with potential to cause harm.
<b>ENT</b>	
Grommets for Glue Ear in children	<p>This is a surgical procedure to insert tiny tubes (grommets) into the eardrum as a treatment for fluid build up (glue ear) when it is affecting hearing in children.</p> <p>Glue ear is a very common childhood problem (4 out of 5 children will have had an episode by age 10), and in most cases it clears up without treatment within a few weeks. Common symptoms can include earache and a reduction in hearing. Often, when the hearing loss is affecting both ears it can cause language, educational and behavioural problems.</p> <p>Please note this guidance only relates to children with Glue Ear (Otitis Media with Effusion) and SHOULD NOT be applied to other clinical conditions where grommet insertion should continue to be normally funded, these include:</p> <ul style="list-style-type: none"> <li>• Recurrent acute otitis media</li> <li>• Atrophic tympanic membranes</li> <li>• Access to middle ear for transtympanic instillation of medication</li> </ul> <p>Investigation of unilateral glue ear in adults</p>

OFFICIAL

Tonsillectomy or recurrent tonsillitis	<p>This guidance relates to surgical procedures to remove the tonsils as a treatment for recurrent sore throats in adults and children.</p> <p>Recurring sore throats are a very common condition that presents a large burden on healthcare; they can also impact on a person's ability to work or attend school. It must be recognised however, that not all sore throats are due to tonsillitis and they can be caused by other infections of the throat. In these cases, removing the tonsils will not improve symptoms.</p>
<b>General surgery</b>	
Haemorrhoid surgery	This procedure involves surgery for haemorrhoids (piles).
<b>Gynaecology</b>	
Hysterectomy for heavy menstrual bleeding	Hysterectomy is the surgical removal of the uterus.
<b>Ophthalmology</b>	
Chalazia removal	This procedure involves incision and curettage (scraping away) of the contents of the chalazion. Chalazia (meibomian cysts) are benign lesions on the eyelids due to blockage and swelling of an oil gland that normally change size over a few weeks. Many but not all resolve within six months with regular application of warm compresses and massage.
<b>Orthopaedics</b>	
Arthroscopic shoulder decompression for subacromial shoulder pain	Arthroscopic sub-acromial decompression is a surgical procedure that involves decompressing the sub-acromial space by removing bone spurs and soft tissue arthroscopically.
Carpal tunnel syndrome release	Open or endoscopic surgical procedure to release median nerve from carpal tunnel.
Dupuytren's contracture release in adults	<p>Dupuytren's contracture is caused by fibrous bands in the palm of the hand which draw the finger(s) (and sometimes the thumb) into the palm and prevent them from straightening fully. If not treated the finger(s) may bend so far into the palm that they cannot be straightened. All treatments aim to straighten the finger(s) to restore and retain hand function for the rest of the patient's life. However none cure the condition which can recur after any intervention so that further interventions are required.</p> <p>Splinting and radiotherapy have not been shown be effective treatments of established Dupuytren's contractures.</p>

OFFICIAL

	<p>Several treatments are available: collagenase injections, needle fasciotomy, fasciectomy and dermofasciectomy. None is entirely satisfactory with some having slower recovery periods, higher complication rates or higher reoperation rates (for recurrence) than others. The need for, and choice of, intervention should be made on an individual basis and should be a shared decision between the patient and a practitioner with expertise in the various treatments of Dupuytren's contractures.</p> <p>No-one knows which interventions are best for restoring and maintaining hand function throughout the rest of the patient's life, and which are the cheapest and most cost-effective in the long term. Ongoing and planned National Institute for Health Research studies aim to answer these conditions.</p>
Ganglion excision	<p>Ganglia are cystic swellings containing jelly-like fluid which form around the wrists or in the hand. In most cases wrist ganglia cause only mild symptoms which do not restrict function, and many resolve without treatment within a year. Wrist ganglion rarely press on a nerve or other structure, causing pain and reduced hand function.</p> <p>Ganglia in the palm of the hand (seed ganglia) can cause pain when carrying objects.</p> <p>Ganglia which form just below the nail (mucous cysts) can deform the nail bed and discharge fluid, but occasionally become infected and can result in aseptic arthritis of the distal finger joint.</p>
Trigger finger release in adults	<p>Trigger digit occurs when the tendons which bend the thumb/finger into the palm intermittently jam in a tight tunnel (flexor sheath) through which they run. It may occur in one or several fingers and causes the finger to "lock" in the palm of the hand. Mild triggering is a nuisance and causes infrequent locking episodes. Other cases cause pain and loss and unreliability of hand function. Mild cases require no treatment and may resolve spontaneously.</p>
<b>Vascular Vein Intervention</b>	
Varicose veins interventions	<p>There are various interventional procedures for treating varicose veins. These include endothermal ablation, ultrasound guided foam sclerotherapy and traditional surgery (this is a surgical procedure that involves ligation and stripping of varicose veins) all of which have been shown to be clinically and cost effective compared to no treatment or treatment with compression hosiery. Varicose veins are common and can markedly affect patients quality of life, can be associated with complications such as eczema, skin changes, thrombophlebitis, bleeding, leg ulceration, deep vein thrombosis and pulmonary embolism that can be life threatening.</p>

# Appendix 2: Clinical glossary

**Acromio-clavicular joint:** a joint at the top of the shoulder between the clavicle and the scapula

**Amenorrhoea:** not having periods (bleeding from the womb)

**Analgesia:** medication to get rid of pain

**Apnoea:** Temporary pausing / stopping of breathing

**Arthroscope:** small camera that is inserted into a joint to examine the inside of the joint

**Arthroscopic shoulder decompression:** surgery to take out small pieces of bone and soft tissue (like tendons) from inside the shoulder by keyhole surgery

**Arthroscopic washout:** operation where an arthroscope (camera) is inserted in to a joint along with fluid that is drained out again.

**Asymptomatic:** not causing any symptoms (problems), for example not causing pain

**Atrophic tympanic membrane:** Thinned, collapsing or retracting ear drum that can affect hearing or lead to erosion of hearing bones

**Benign skin lesions:** lumps or bumps on the skin that are not suspicious for skin cancer

**Biopsy:** small sample of tissue, for example the lining of the womb, is taken out for examination under a microscope

**Breast hyperplasia:** enlargement of the breasts

**Breast reduction:** surgery to reduce the size of the breast by removing fat, breast tissue and skin

**Calcific tendinopathy:** a condition where small particles or crystals collect in the tendons that connect muscle to bone. It occurs most commonly in the shoulder.

**Carpal tunnel syndrome:** pressure on a nerve in the wrist causing pain, tingling or numbness in the fingers

## OFFICIAL

**Cervix:** opening of the womb

**Chalazia (meibomian cyst):** small lump in the eyelid caused by a blocked and swollen oil gland

**Chronic venous insufficiency:** a condition where the veins are not working properly and blood pools or collects in the vein and is not returned to the heart

**Complex regional pain syndrome:** severe pain and swelling in the hand that sometimes occurs following surgery

**Deep vein thrombosis:** blood clot that develops in one of the large veins in the body for example in the lower leg

**D&C:** dilatation and curettage, a procedure where the opening to the womb (the cervix) is widened (dilated) and the lining of the womb is scraped out (curettage)

**Digital artery:** blood vessel in a finger

**Distal interphalangeal joint mucous cysts:** ganglions or fluid filled sacks that occur near the tip of the finger at the joint near the nailbed

**Dupuytren's contracture:** small nodules or thickening on the tendons in the hand that prevent the fingers from straightening completely

**Endothermal ablation:** radio waves or lasers are used to seal off the varicose vein

**Fasciectomy:** removing thickened tissue by surgery

**Fasciotomy:** cutting or dividing thickened tissue

**Fibroids:** growths in the uterus (womb) that are not cancer but can cause heavy periods and pain

**Ganglion:** small cyst or fluid filled sac that arises near a joint or a tendon, for example at the wrist, the ganglion can press on a nerve causing pain or tingling.

**Ganglion excision:** surgery to remove a ganglion and the stalk from the tendon it is attached to.

**Globus:** Persistent feeling of something in the throat when there is nothing there

**Glue Ear:** Build up of fluid in the middle part of the ear, behind the ear drum.

**Grommet:** Tiny plastic tube inserted through ear drum during a surgical procedure

**Gynaecomastia:** enlargement breast tissue in men

**Haemarthrosis:** bleeding inside a joint, for example the knee joint

## OFFICIAL

**Haemorrhoids (piles):** swellings containing blood vessels that come from inside the bottom

**Heavy menstrual bleeding:** heavy bleeding from the womb during a woman's period

**Hypermastia:** excessively large breasts

**Hysterectomy:** surgery to remove the uterus (womb)

**Hysteroscopy :** camera test of the womb

**Ischaemic fissuring:** a cut or tear in the anus caused by a complication of a surgical intervention

**Incontinence:** lack of control over going to the toilet (urine or stool), so not being able to hold in stool.

**Intertrigo:** skin rash that develops in between skin folds

**Intrauterine system (IUS):** small plastic device that is inserted into the womb via the cervix

**Ligation:** tying off

**Locked finger:** the finger cannot be straightened

**Obstructive sleep apnoea (OSA):** Throat can partially or completely close whilst sleeping, temporarily stopping or reducing breathing which can disturb sleep and oxygen levels.

**Oophorectomy:** removal of the ovaries during surgery

**Osteoarthritis:** a degeneration of the joints, especially the knees and hips that affects people from middle age onward, causing stiffness and pain in the joints

**Osteotomy:** surgery where bone in a joint is shaved away to re-align a joint that has become crooked

**Otitis Media:** Infection in the middle part of the ear behind the ear drum

**Parapharyngeal abscess:** Collection of pus in deep spaces of neck that may have spread from a tonsil infection

**Pulmonary embolism:** a blocked blood vessel in the lung that can be life threatening if not treated quickly

**Radiofrequency denervation:** procedure where the nerves that are connected to the small joints in the spine (facet joints) are destroyed to numb pain

## OFFICIAL

**Rotator cuff tear:** a tear in the tendons that connect muscles to the top of the humerus (the bone in the upper arm bone). A tear can cause pain or weakness in the arm.

**Sciatica:** tingling and pain in the buttocks and travelling down the leg due to irritation of the sciatic nerve

**Sclerotherapy:** injection of a substance into the varicose vein to shrink it

**Shoulder girdle dysfunction:** pain and restricted movement of the shoulder

**Spinal injection:** using a needle to insert medication, for example steroid, into the back around the nerves near the spine

**Splinting:** a support is used to keep a body part from moving to allow it to heal

**Stenosis:** tightening of an opening in the body, for example the anus

**Subacromial pain or impingement:** the bones and tendons in the shoulder rub against each other when the arm is raised, causing pain.

**Subcutaneous lesion:** a lump or bump that lies underneath the skin  
**Trigger finger:** tightening of the tendons in a finger that prevent the finger from being completely straightened.

**Systematic Review:** Literature review of multiple existing research studies to answer defined research question

**Tendon bowstringing:** tendon comes away from its attachments and causes difficulty in bending the finger

**Therapeutic mammoplasty:** breast surgery to remove cancer and reshape the breast

**Thrombophlebitis:** inflammation that causes a blood clot in a vein causing redness and pain

**Transtympanic instillation of medication:** Injection of medication through the ear drum e.g. for the treatment of balance problems or sudden nerve related hearing loss.

**Trigger finger release:** surgery to cut the tendon sheath (the coat around the tendon) to release the tendon.

**Truncal reflux:** backflow of blood the wrong way through a vein

**Truncal vein:** superficial vein in the body, lying outside the muscles but underneath the skin

**Varicose veins:** veins that are swollen, enlarged, and twisted, usually in the legs

## OFFICIAL

**Venous disease:** a long term condition related to veins including varicose veins and chronic venous insufficiency

# Appendix 3: Technical glossary

**Prior Approval Schemes:** Are referred to in the NHS Standard Contract as a request from a clinician (or provider) to a commissioner to undertake a specific treatment. The Evidence-Based Interventions Programme refers to two schemes: Individual Funding Requests for Category 1 interventions and Prior Approval for Category 2 interventions.

**Individual Funding Request (IFR):** Is a request received from a clinician providing care to a patient, for:

- A specific treatment that is not covered by existing policy or for a service which is not routinely commissioned by a CCG, or
- Where the CCG is responsible for commissioning the service/treatment in question and/or a local policy is in place however the patient does not meet the criteria and is deemed to be clinically exceptional.

Arguments on the basis of exceptionality are requests where a patient is deemed to have exceptional clinical circumstances, i.e. a patient who has clinical circumstances which, taken as a whole, are outside the range of clinical circumstances presented by a patient within the normal population of patients with the same medical condition and at a similar stage of progression as the patient, exceptional to the cohort.

**Prior Approval (PA):** Is a process in which clinicians demonstrate how a patient meets set threshold criteria prior to referring to secondary care and/or by consultants prior to listing for surgery or performing a procedure for which a CCG routinely commissions and is within agreed contracts.

- Prior Approval means that a General Practitioner and/or provider must seek the agreement of the responsible commissioner to fund a treatment for an individual for an intervention which there is a CCG policy before that treatment is carried out.
- The Prior Approval process then compares requests for elective procedure against a set of threshold criteria for the Prior Approval process.
- On occasions patients may fall outside of the PA threshold criteria and clinicians may appeal by demonstrating how the patient is clinically exceptional. In these cases the request is then considered via the Individual Funding Request process.

# Appendix 4: Procedure and diagnostic codes

For each of the 17 interventions, the clinical definitions have been converted into combinations of one or more OPCS procedure codes and ICD-10 diagnosis codes. This process was informed by a combination of NICE, NICE-accredited and specialist society guidance alongside engagement with our demonstrator sites to identify the appropriate codes.

Our analysis is based on SUS+ data of spells completing in 2017/18. The following descriptors use Microsoft SQL Server structure but are easily adaptable to other systems. For reference:

- A “%” symbol represents a wildcard for zero or more characters.
- Values in square brackets mean “one of these characters”. E.g. [03] mean 0 or 3 and [0-3] means 0 or 1 or 2 or 3.
- The field “der\_diagnosis\_all” is a concatenation of all diagnosis fields in all episodes within the spell.

## Adult snoring surgery (not OSA)

when left(der.Spell\_Dominant\_Procedure,4) in ('F324','F325','F326') and der.Spell\_Primary\_Diagnosis not like '%G473%' and APCS.Age\_At\_Start\_of\_Spell\_SUS between 18 and 120 then 'A\_snoring'

## Dilatation and curettage for heavy menstrual bleeding

when left(der.Spell\_Dominant\_Procedure,4) in ('Q103') and apcs.der\_diagnosis\_all not like '%O0[0-8]%' and apcs.der\_diagnosis\_all not like '%O6[0-9]%' and apcs.der\_diagnosis\_all not like '%O7[0-5]%' then 'B\_menstr\_D&C'

## Knee arthroscopy with osteoarthritis

when der.Spell\_Dominant\_Procedure in ('W821','W822','W823','W828','W829','W851','W852','W853','W858','W859','W861+KNEE','W831+KNEE','W832+KNEE','W833+KNEE','W834+KNEE','W835+KNEE','W836+KNEE','W837+KNEE','W838+KNEE','W839+KNEE','W841+KNEE','W842+KNEE','W843+KNEE','W844+KNEE') and (APCS.Age\_At\_Start\_of\_Spell\_SUS between 18 and 120) and apcs.der\_diagnosis\_all not like '%C[0-9][0-9]%' and der.Spell\_Primary\_Diagnosis like 'M1[57]%' then 'C\_knee\_arth'

## Injections for nonspecific low back pain without sciatica

when left(der.Spell\_Dominant\_Procedure,4) in ('A521','A522','A528','A529','A577','A735','V363','V368','V369','V382','V383','V384','V385','V386','V388','V389','V544','W903') and left(der.spell\_primary\_diagnosis,4) in

OFFICIAL

('G834','G551','M518','M519','M545','M549') and apcs.der\_procedure\_all like '%Z67[67]%' then 'D\_low\_back\_pain\_inj'

**Breast reduction**

when left(der.Spell\_Dominant\_Procedure,4) in ('B311') and apcs.der\_diagnosis\_all not like '%C[0-9][0-9]%' then 'E\_breast\_red'

**Removal of benign skin lesions**

when left(der.Spell\_Dominant\_Procedure,4) in ('S063','S064','S065','S066','S067','S068','S069','S081','S082','S083','S088','S089','S091','S092','S093','S094','S095','S098','S099','S101','S102','S111','S112','D021','D022','D028','D029') and APCS.Der\_Diagnosis\_All not like '%C4[3469]%' then 'F\_skin\_lesions'

**Grommets for glue ear in children**

when left(der.Spell\_Dominant\_Procedure,4) in ('D151','D289') and (der.Spell\_Primary\_Diagnosis like 'H65[23]%' or der.Spell\_Primary\_Diagnosis like 'H66[1-9]%) and (apcs.age\_at\_start\_of\_Spell\_SUS between 1 and 17 or apcs.age\_at\_start\_of\_Spell\_SUS between 7001 and 7007 ) then 'G\_gromm'

**Tonsillectomy for recurrent tonsillitis**

when left(der.Spell\_Dominant\_Procedure,4) in ('F341','F342','F343','F344','F345','F346','F347','F348','F349','F361') and apcs.der\_diagnosis\_all not like '%C[0-9][0-9]%' and apcs.der\_diagnosis\_all not like '%G47%' and apcs.der\_diagnosis\_all not like '%J36%' then 'H\_tonsil'

**Haemorrhoid surgery**

when left(der.Spell\_Dominant\_Procedure,4) in ('H511','H512','H513','H518','H519') and apcs.der\_diagnosis\_all not like '%C[0-9][0-9]%' then 'I\_haemmor'

**Hysterectomy for heavy menstrual bleeding**

when left(der.Spell\_Dominant\_Procedure,4) in ('Q072','Q074','Q078','Q079','Q082','Q088','Q089') and apcs.der\_diagnosis\_all not like '%C[0-9][0-9]%' and apcs.der\_diagnosis\_all not like '%O0[0-8]%' and apcs.der\_diagnosis\_all not like '%O6[0-9]%' and apcs.der\_diagnosis\_all not like '%O7[0-5]%' then 'J\_hysterec'

**Chalazia removal**

when left(der.Spell\_Dominant\_Procedure,4) in ('C121','C122','C124','C191','C198') and left(der.Spell\_Primary\_Diagnosis,4) in ('H001') then 'K\_chalazia'

**Arthroscopic shoulder decompression for subacromial shoulder pain**

when (der.Spell\_Dominant\_Procedure ='W844+SHOULDER' or (der.Spell\_Dominant\_Procedure ='O291' and apcs.der\_procedure\_all like '%Y767%')) and (der.Spell\_Primary\_Diagnosis like 'M754%' or der.Spell\_Primary\_Diagnosis like 'M2551%') then 'L\_should\_decom'

**Carpal tunnel syndrome release**

when left(der.Spell\_Dominant\_Procedure,4) in ('A651','A659') and der.Spell\_Primary\_Diagnosis like '%G560%' then 'M\_carpal'

## OFFICIAL

### **Dupuytren's contracture release in adults**

when left(der.Spell\_Dominant\_Procedure,4) in ('T521','T522','T525','T526','T541') and (APCS.Age\_At\_Start\_of\_Spell\_SUS between 18 and 120) and left(der.Spell\_Primary\_Diagnosis,4)='M720' then 'N\_dupuytr'

### **Ganglion excision**

when left(der.Spell\_Dominant\_Procedure,4) in ('T591','T592','T598','T599','T601','T602','T608','T609') and der.Spell\_Primary\_Diagnosis like '%M674%' then 'O\_ganglion'

### **Trigger finger release in adults**

when der.Spell\_Dominant\_Procedure in ('T692+HAND','T691+HAND','T698+HAND','T699+HAND','T701+HAND','T702+HAND','T718+HAND','T719+HAND','T723+HAND','T728+HAND','T729+HAND','Z894+HAND','Z895+HAND','Z896+HAND','Z897+HAND') and (APCS.Age\_At\_Start\_of\_Spell\_SUS between 18 and 120) and der.Spell\_Primary\_Diagnosis like '%M653%' then 'P\_trigger\_fing'

### **Varicose vein interventions**

when left(der.Spell\_Dominant\_Procedure,4) in ('L832','L838','L839','L841','L842','L843','L844','L845','L846','L848','L849','L851','L852','L853','L858','L859','L861','L862','L863','L868','L869','L871','L872','L873','L874','L875','L876','L877','L878','L879','L881','L882','L883','L888','L889') and der.Spell\_Primary\_Diagnosis like ('%I8[03]%' ) then 'Q\_var\_veins'