

ID	Stage	Location	Short description updated	Numerator updated	Denominator updated
LL01	Immediate Care	Leg	Proportion of adult patients with a lower leg wound without NWCSP 'Red Flag Symptoms' being treated with first line mild graduated compression	Total number of patients aged 18+ years with a lower leg wound without NWCSP 'Red Flag Symptoms' identified by a health care practitioner within the reporting period with intention to treat with first line mild graduated compression	Total number of patients aged 18+ years with a lower leg wound without NWCSP 'Red Flag Symptoms' as identified by a health care practitioner assessed within the reporting period
LL02	Assessment	Leg	Proportion of adult patients with a lower leg wound receiving initial full assessment within 14 days of initial presentation	Total number of patients aged 18+ years with documentation of a lower leg wound assessment that meets the minimum requirements described in NWCSP Lower leg Assessment Essential Criteria (within 14 days of being identified by a health care practitioner) within the reporting period	Total number of patients aged 18+ years with a lower leg wound assessment within the reporting period
LL03	Referral	Both	Proportion of adult patients diagnosed with venous insufficiency and with Venous Leg Ulceration (VLU) referred to vascular service	Total number of patients aged 18+ years diagnosed with venous insufficiency and with Venous Leg Ulceration (VLU) within the reporting period referred to vascular service	Total number of patients aged 18+ years diagnosed with venous insufficiency and with Venous Leg Ulceration (VLU) within the reporting period
LL04	Referral	Both	Proportion of adult patients diagnosed with Peripheral Arterial Disease (PAD) with a lower limb wound referred to vascular service	Total number of patients aged 18+ years diagnosed with Peripheral Arterial Disease (PAD) and a lower limb wound within the reporting period referred to the vascular service	Total number of patients aged 18+ years diagnosed with Peripheral Arterial Disease (PAD) with a lower limb wound within the reporting period
LL05	Referral	Foot	Proportion of adult patients in hospital, with diabetes and a foot wound referred to the multidisciplinary foot care service within 24 hours of being seen by a health care practitioner	Total number of patients aged 18+ years in hospital with a type 1 or 2 diabetes diagnosis and a foot wound referred to the multidisciplinary foot care service or foot protection service within 24 hours of being seen by a health care practitioner within the reporting period	Total number of patients aged 18+ years in hospital with a type 1 or 2 diabetes diagnosis and a foot wound seen by a health care practitioner within the reporting period
LL06	Referral	Foot	Proportion of adult non-diabetic patients in hospital with a foot wound, referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner	Total number of non-diabetic patients aged 18+ years in hospital with a foot wound referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner within the reporting period	Total number of patients aged 18+ years in hospital with a type 1 or 2 diabetes diagnosis and a foot wound seen by a health care practitioner within the reporting period
LL07	Referral	Foot	Proportion of adult patients in the community with diabetes and a foot wound referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner	Total number of patients aged 18+ years in the community with type 1 or 2 diabetes diagnosis and a foot wound, referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner within the reporting period	Total number of patients aged 18+ years in the community with type 1 or 2 diabetes and a foot wound seen by a healthcare practitioner within the reporting period
LL08	Referral	Foot	Proportion of adult non-diabetic patients in the community with a foot wound, referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner	Total number of non-diabetic patients aged 18+ years in the community with a foot wound, referred to the multidisciplinary foot care service or foot protection service within 1 working day of being seen by a health care practitioner with the reporting period	Total number of non-diabetic patients aged 18+ years in the community with a foot wound seen by a health care practitioner within the reporting period
LL09	Treatment	Leg	Proportion of adult patients with a lower leg wound and an adequate arterial supply, where no aetiology other than venous insufficiency is suspected treated with strong compression	Total number of patients aged 18+ years with a lower leg wound with an adequate arterial supply (ABPI > 0.8-1.3) and where no other condition that contra-indicates compression therapy is suspected, treated with a minimum of 40mmHg compression therapy within the reporting period	Total number of patients aged 18+ years treated in the defined service with a lower leg wound with an adequate arterial supply (ABPI > 0.8-1.3) within the reporting period
LL10	Treatment	Leg	Proportion of adult patients with venous insufficiency and with Venous Leg Ulceration (VLU) referred to vascular service who undergo endovenous ablation	Total number of patients aged 18+ years diagnosed with venous insufficiency and with Venous Leg Ulceration (VLU) referred to a vascular service who undergo endovenous ablation within the reporting period	Total number of patients aged 18+ years diagnosed with venous insufficiency and with Venous Leg Ulceration (VLU) referred to a vascular service treated within the reporting period
LL11	Treatment	Both	Proportion of adult patients referred to vascular surgeons for assessment who undergo revascularisation	Total number of patients aged 18+ years referred to vascular surgeons for assessment within the reporting period who undergo revascularisation	Total number of patients aged 18+ years referred to vascular surgeons for assessment within the reporting period
LL12a	Outcomes - healing	Leg	Proportion of adult patients diagnosed with Venous Leg Ulceration (VLU) recorded as healed 0-12 weeks after identification by a health care practitioner	Total number of patients aged 18+ years diagnosed with Venous Leg Ulceration (VLU) recorded as healed 0-12 weeks after identification by a health care practitioner in the reporting period	Total number of patients aged 18+ years with Venous Leg Ulceration (VLU), recorded as healed within the reporting period

LL30	Outcomes - sub-optimal treatment	Foot	Proportion of adult non-diabetic patients and with a foot ulceration who have had a major amputation	Total number of non-diabetic patients aged 18+ years and treated for a foot ulceration, who have had a major amputation (amputation above the ankle) within the reporting period	Total number of patients aged 18+ years with type 1 or 2 diabetes treated for a foot ulceration within the reporting period
------	----------------------------------	------	--	--	---

